

Quiver D -Modules and Homology of Local Systems over an Arrangement of Hyperplanes

S. Khoroshkin[◦] and A. Varchenko^{*}

[◦]*Institute of Theoretical and Experimental Physics, 117259 Moscow, Russia*

^{*}*Department of Mathematics, University of North Carolina at Chapel Hill, NC 27599-3250, USA*

Abstract

Let $\mathcal{C}$ be an arrangement of hyperplanes in $\mathbb{C}^N$, D the ring of algebraic differential operators on $\mathbb{C}^N$. We define a category of quivers associated with $\mathcal{C}$. A quiver is a collection of vector spaces, attached to strata of the arrangement, and suitable linear maps between the spaces. To a quiver we assign a D -module on $\mathbb{C}^N$, called a quiver D -module. We describe basic operations for D -modules in terms of linear algebra of quivers. We give an explicit construction of a free resolution of a quiver D -module and use the construction to describe the associated perverse sheaf. As an application, we calculate the cohomology of $\mathbb{C}^N$ with coefficients in the quiver perverse sheaf (under certain assumptions).

CONTENTS

1. Introduction	2
2. Arrangements	9
2.1. Graph of an arrangement	9
2.2. Framings of an arrangement	10
2.3. Principal open subsets X^n	12
2.4. Truncated graphs	12
2.5. Orlik-Solomon algebra and flag complex	13
3. Quivers	15
3.1. Quivers of an admissible graph	15
3.2. Quivers of level n	18
3.3. Restrictions of quivers	19
3.4. Direct images of quivers	20
3.5. Example	23
3.6. Direct images of level zero quivers	24
3.7. Monodromy operators for quivers $\mathbb{J}_{0,!}(\mathcal{W}_{\Gamma^0})$ and $\mathbb{J}_{0,*}(\mathcal{W}_{\Gamma^0})$	26
3.8. Example	27
4. Quiver D -modules	29
4.1. Modules over rings of differential operators	29
4.2. Quiver $\mathcal{D}_X$ -modules	30
4.3. Example	32
4.4. Quiver $\mathcal{D}_{X^n}$ -modules	33
4.5. Direct and inverse images of quiver $\mathcal{D}$ -modules	36
4.6. Specialization of a quiver $\mathcal{D}_X$ -module to a stratum	37
4.7. Free resolution of quiver $\mathcal{D}_{X^0}$ -modules	40
4.8. Free resolution of quiver $\mathcal{D}_X$ -modules	41
4.9. Derived categories and derived functors	43

4.10.	Duality of quiver $\mathcal{D}_X$ -modules	44
4.11.	Fourier transform of quiver $\mathcal{D}_X$ -modules	46
5.	Quiver perverse sheaves	47
5.1.	Perverse sheaves over analytic varieties	47
5.2.	De Rham functor	49
5.3.	De Rham complex of a quiver $\mathcal{D}_X$ -module	50
5.4.	Quiver perverse sheaves	51
5.5.	Cohomology groups of $\mathbb{C}^N$ with coefficients in quiver perverse sheaves	52
6.	Equivariant structures	53
6.1.	Arrangement with a group action	53
6.2.	Equivariant quivers	54
6.3.	Equivariant quiver $\mathcal{D}$ -modules	54
6.4.	Equivariant quiver perverse sheaves	58
6.5.	Cohomology of $\mathbb{C}^N$ with coefficients in equivariant quiver perverse sheaves	59
6.6.	Application to discriminantal arrangements	60
7.	Proofs	63
7.1.	Proof of Propositions 3.1 and 3.2	63
7.2.	Proof of Propositions 3.9–3.11	65
7.3.	Proof of Proposition 4.1	66
7.4.	Proof of Proposition 4.2	70
7.5.	Proof of Proposition 4.4	70
7.6.	Proof of Proposition 4.3	71
7.7.	Proof of Theorem 4.6	79
7.8.	Proof of Theorem 4.11	83
7.9.	Proof of Theorem 4.13.	84
7.10.	Proof of Theorem 4.14	87
7.11.	Proof of Theorem 5.6	88
7.12.	Proof of Theorem 6.6	92
	References	92

1. INTRODUCTION

1. It is known that nontrivial homological characteristics of a space can be expressed in terms of linear algebra, if the space is an affine space with an arrangement of hyperplanes. An example is the description by Arnold, Brieskorn, Orlik and Solomon of the cohomology groups of the complement to the arrangement in terms of combinatorics of intersections of hyperplanes [Ar, Bs, OS]. On the other hand, holonomic D -modules with regular singularities and constructible perverse sheaves on algebraic manifolds also admit combinatorial description, according to Beilinson, Kashiwara, Malgrange [Be, K3, M]. In this paper we combine the two points of view and describe a certain category of holonomic D -modules and associated perverse sheaves, attached to an arrangement of hyperplanes in an affine space.

2. Let $\mathcal{C}$ be an arrangement of hyperplanes in a complex affine space X . The intersections of hyperplanes define in X the structure of a stratified space. The strata and their adjacencies are described by the graph Γ of the arrangement. The vertices $\alpha \in I(\Gamma)$

correspond to strata X_α . If the closure $\overline{X}_\alpha$ of a stratum X_α contains a stratum X_β and $\text{codim}_{\overline{X}_\alpha} X_\beta = 1$, then the vertices α and β are connected by an edge and we write $\alpha \succ \beta$. To each vertex $\alpha \in I(\Gamma)$ we attach a nonnegative integer $l(\alpha)$, equal to the codimension of X_α in X .

We assign to the graph Γ quivers with relations, which form an abelian category Qui_Γ . A quiver $\mathcal{V} \in Qui_\Gamma$ consists of a collection of finite-dimensional vector spaces V_α , $\alpha \in I(\Gamma)$, and linear maps $A_{\alpha,\beta} : V_\beta \rightarrow V_\alpha$, $\alpha, \beta \in I(\Gamma)$, such that

- (a) $A_{\alpha,\beta} = 0$ if α and β are not adjacent;
- (b) $\sum_\beta A_{\alpha,\beta} A_{\beta,\gamma} = 0$ if $|l(\alpha) - l(\gamma)| = 2$;
- (c) $\sum_\beta A_{\alpha,\beta} A_{\beta,\gamma} = 0$ if $l(\alpha) = l(\gamma)$, $\alpha \neq \gamma$, and there exists $\delta \in I(\Gamma)$, adjacent to α and γ , such that $l(\delta) = l(\alpha) + 1$.

Example 1.1. Let X be $\mathbb{C}$ with coordinate z . Let $\mathcal{C} = \{z = 0\}$. The space X has two strata: $X_\emptyset = \{z \neq 0\}$ and $X_\alpha = \{0\}$. The graph Γ has two vertices, $\emptyset$ and α , connected by an edge, $l(\alpha) = 1$, $l(\emptyset) = 0$. A quiver $\mathcal{V} \in Qui_\Gamma$ is a collection of two vector spaces, $V_\emptyset$ and V_α , and two linear maps, $A_{\emptyset,\alpha} : V_\alpha \rightarrow V_\emptyset$ and $A_{\alpha,\emptyset} : V_\emptyset \rightarrow V_\alpha$.

Let D_X denote the ring of algebraic differential operators on X and $\mathcal{D}_X$ the corresponding sheaf of rings on X . To a quiver $\mathcal{V} \in Qui_\Gamma$ we attach a holonomic D_X -module $E\mathcal{V}$. This D_X -module is given by generators and relations. The generators are elements of the vector space $\bigoplus_{\alpha \in I(\Gamma)} \overline{V}_\alpha$. Here $\overline{V}_\alpha = \overline{\Omega}_\alpha \otimes V_\alpha$, and $\overline{\Omega}_\alpha$ denotes the one-dimensional vector space of top exterior forms on $\overline{X}_\alpha$, invariant with respect to affine translations along $\overline{X}_\alpha$. The relations in $E\mathcal{V}$ are:

- (a) for any generator $\overline{\omega}_\alpha \otimes v_\alpha \in \overline{V}_\alpha$ and any vector field ξ with constant coefficients, which is parallel to X_α , the element $\xi(\overline{\omega}_\alpha \otimes v_\alpha)$ equals to a certain linear combination of the vectors $\overline{\omega}_\beta \otimes A_{\beta,\alpha}(v_\alpha)$, $\beta \in I(\Gamma)$, $\alpha \succ \beta$,
- (b) for any affine function f , vanishing on X_α , the element $f(\overline{\omega}_\alpha \otimes v_\alpha)$ equals to a certain linear combination of the vectors $\overline{\omega}_\beta \otimes A_{\beta,\alpha}(v_\alpha)$, $\beta \in I(\Gamma)$, $\beta \succ \alpha$,

see formulas (4.25), (4.26).

Example 1.2. In Example 1.1, the D_X -module $E\mathcal{V}$, assigned to the quiver $\mathcal{V} = \{V_\emptyset, V_\alpha, A_{\emptyset,\alpha}, A_{\alpha,\emptyset}\}$ is generated by vectors $dz \otimes v_\emptyset \in \overline{V}_\emptyset$, where $v_\emptyset \in V_\emptyset$, and $1 \otimes v_\alpha \in \overline{V}_\alpha$, where $v_\alpha \in V_\alpha$. The relations are: $\frac{d}{dz}(dz \otimes v_\emptyset) = 1 \otimes A_{\alpha,\emptyset}(v_\emptyset)$ for any $v_\emptyset \in V_\emptyset$ and $z(1 \otimes v_\alpha) = dz \otimes A_{\emptyset,\alpha}(v_\alpha)$ for any $v_\alpha \in V_\alpha$.

The assignment $\mathcal{V} \mapsto E\mathcal{V}$ defines a functor E from the category Qui_Γ to the category $\mathcal{M}_X^{hol}$ of holonomic $\mathcal{D}_X$ -modules. We describe a full subcategory in Qui_Γ of non-resonant quivers, such that the restriction of E to this subcategory is full and faithful.

For $n = 0, \dots, N$, let X^n denote the complement in X to the union of all strata of codimension greater than n . This set X^n is called the principal open subset. The principal subset X^0 is the complement to the union of hyperplanes of the arrangement. All X^n are stratified spaces. The adjacency structure of the stratification of X^n is described by the graph Γ^n , which is a suitable truncation of Γ . We introduce an abelian category Qui_{Γ^n} of quivers of Γ^n . The quivers of Qui_{Γ^n} are also called level n quivers. To each level n quiver $\mathcal{V}_{\Gamma^n} \in Qui_{\Gamma^n}$ we assign a sheaf of $\mathcal{D}_{X^n}$ -modules, denoted by $E^n \mathcal{V}_{\Gamma^n} \in \mathcal{M}_{X^n}^{hol}$.

Example 1.3. In Example 1.1, we have $X^0 = X_\emptyset = \{z \neq 0\}$, $X^1 = X = \mathbb{C}$, $\Gamma^1 = \Gamma$. The graph Γ^0 has one vertex $\emptyset$ and one edge, which is a loop. A quiver $\mathcal{V}_{\Gamma^0} \in \text{Qui}_{\Gamma^0}$ consists of a vector space $V_\emptyset$ and a linear map $A_\emptyset^\emptyset : V_\emptyset \rightarrow V_\emptyset$. The $\mathcal{D}_{X^0}$ -module $E^0 \mathcal{V}_{\Gamma^0}$, assigned to this quiver, is generated by the elements $dz \otimes v_\emptyset \in \overline{V}_\emptyset$, subject to the relations $z \frac{d}{dz} (dz \otimes v_\emptyset) = dz \otimes A_\emptyset^\emptyset(v_\emptyset)$.

For $k \leq l$ we have inclusions $j_{l,k} : X^k \rightarrow X^l$ of principal open subsets and related functors $j_{l,k}^*$ and $j_{l,k,*}$ of inverse and direct images of $\mathcal{D}$ -modules. We define quiver inverse and direct image functors $\mathbb{J}_{l,k}^* : \text{Qui}_{\Gamma^l} \rightarrow \text{Qui}_{\Gamma^k}$ and $\mathbb{J}_{l,k,*} : \text{Qui}_{\Gamma^k} \rightarrow \text{Qui}_{\Gamma^l}$, respectively. The functor $\mathbb{J}_{l,k}^*$ is left adjoint to the functor $\mathbb{J}_{l,k,*}$. We have an isomorphism of $\mathcal{D}_{X^k}$ -modules $E^k \mathbb{J}_{l,k}^*(\mathcal{V}_{\Gamma^l}) \approx j_{l,k}^* E^l(\mathcal{V}_{\Gamma^l})$ for any level l quiver $\mathcal{V}_{\Gamma^l}$. We construct an isomorphism of $\mathcal{D}_{X^l}$ -modules $j_{l,k,*} E^k(\mathcal{V}_{\Gamma^k})$ and $E^l \mathbb{J}_{l,k,*}(\mathcal{V}_{\Gamma^k})$ under certain non-resonance conditions on the level k quiver $\mathcal{V}_{\Gamma^k}$.

Example 1.4. In Examples 1.1 and 1.3, the level zero quiver $\mathbb{J}_{1,0}^*(\mathcal{V})$, associated to a level one quiver $\mathcal{V} = \{V_\emptyset, V_\alpha, A_{\emptyset,\alpha}, A_{\alpha,\emptyset}\}$, is $\{U_\emptyset, B_\emptyset^\emptyset\}$, where $U_\emptyset = V_\emptyset$, and $B_\emptyset^\emptyset = A_{\emptyset,\alpha} A_{\alpha,\emptyset}$. The level one quiver $\mathbb{J}_{1,0,*}(\mathcal{V}_{\Gamma^0})$, associated to a level zero quiver $\mathcal{V}_{\Gamma^0} = \{V_\emptyset, A_\emptyset^\emptyset\}$, is $\{W_\emptyset, W_\alpha, C_{\emptyset,\alpha}, C_{\alpha,\emptyset}\}$, where $W_\emptyset = V_\emptyset$, $W_\alpha = V_\emptyset$, $C_{\emptyset,\alpha} = \text{Id}_{V_\emptyset}$, $C_{\alpha,\emptyset} = A_\emptyset^\emptyset$. The quiver $\mathbb{J}_{1,0,*}(\mathcal{V}_{\Gamma^0})$ is non-resonant, if any two eigenvalues of the operator $A_\emptyset^\emptyset$ do not differ by a nonzero integer, and the only possible integer eigenvalue of $A_\emptyset^\emptyset$ is zero.

There is a *specialization* construction in the theory of $\mathcal{D}$ -modules due to Kashiwara [K3]. Let Y be a smooth complex algebraic variety, $Z \subset Y$ a subvariety, $T_Z Y \rightarrow Z$ the normal bundle. Given a $\mathcal{D}_Y$ -module M and some additional data, the specialization construction produces a $\mathcal{D}$ -module on the total space $T_Z Y$ of the normal bundle. This $\mathcal{D}_{T_Z Y}$ -module is called *the specialization* of M to Z and denoted by $Sp_Z(M)$.

The specialization construction gives a collection of functors between quiver $\mathcal{D}$ -modules.

We consider the case when Y is the affine space $X = \mathbb{C}^N$ with a central arrangement $\mathcal{C}$ and M is the quiver $\mathcal{D}$ -module $E\mathcal{V}$ associated with a quiver $\mathcal{V}$ of $\mathcal{C}$. Then it turns out that for any closed stratum $\overline{X}_\alpha$ of X , the specialization $Sp_{\overline{X}_\alpha}(E\mathcal{V})$ is isomorphic to a suitable quiver $\mathcal{D}$ -module on $T_{\overline{X}_\alpha} X$.

More precisely, we construct an arrangement of hyperplanes $\mathcal{C}_\alpha$ in $T_{\overline{X}_\alpha} X$ and a quiver $Sp_\alpha(\mathcal{V})$ of the arrangement $\mathcal{C}_\alpha$ such that the specialization $Sp_{\overline{X}_\alpha}(E\mathcal{V})$ is isomorphic to the quiver $\mathcal{D}$ -module $ESp_\alpha(\mathcal{V})$ associated with the quiver $Sp_\alpha(\mathcal{V})$, if the linear maps of the initial quiver $\mathcal{V}$ satisfy certain non-resonance conditions.

Example 1.5. In Examples 1.1 and 1.2, we identify the space $T_{\overline{X}_\alpha} X$ with $X = \mathbb{C}$. Then the arrangement $\mathcal{C}_\alpha$ is identified with $\mathcal{C}$, the quiver $Sp_\alpha(\mathcal{V})$ is identified with $\mathcal{V}$. Introduce the linear operator

$$S = A_{\emptyset,\alpha} A_{\alpha,\emptyset} + A_{\alpha,\emptyset} A_{\emptyset,\alpha} : V_\emptyset \oplus V_\alpha \rightarrow V_\emptyset \oplus V_\alpha.$$

If any two eigenvalues of S do not differ by a nonzero integer and if S has no nonzero integer eigenvalues, then the $\mathcal{D}_{T_{\overline{X}_\alpha}}$ -modules $Sp_{\overline{X}_\alpha}(E\mathcal{V})$ and $ESp_\alpha(\mathcal{V})$ are isomorphic.

For any quiver $\mathcal{V}_\Gamma \in \text{Qui}$, we construct a free $\mathcal{D}_X$ -module resolution $\mathcal{R}E\mathcal{V}_\Gamma$ of the quiver $\mathcal{D}_X$ -module $E\mathcal{V}_\Gamma$.

Example 1.6. In Example 1.2, the free $\mathcal{D}_X$ -module resolution $\mathcal{R}E\mathcal{V}$ of the quiver $\mathcal{D}_X$ -module $E\mathcal{V}$ is the complex

$$\mathcal{R}E\mathcal{V} : 0 \rightarrow \mathcal{R}^{-1}E\mathcal{V} \xrightarrow{d} \mathcal{R}^0E\mathcal{V} \xrightarrow{\nu} E\mathcal{V} \rightarrow 0.$$

Here $\mathcal{R}^0E\mathcal{V}$ is the free $\mathcal{D}_X$ -module

$$\mathcal{D}_X \otimes \overline{V}_\emptyset \oplus \mathcal{D}_X \otimes \overline{V}_\alpha.$$

The term $\mathcal{R}^{-1}E\mathcal{V}$ is the free $\mathcal{D}_X$ -module

$$\mathcal{D}_X \otimes T_\emptyset \otimes \overline{V}_\emptyset \oplus \mathcal{D}_X \otimes F_\alpha \otimes \overline{V}_\alpha,$$

where $T_\emptyset$ is the one-dimensional space $\mathbb{C} \cdot \frac{d}{dz}$, and F_α is the one-dimensional space $\mathbb{C} \cdot z$ of affine functions, vanishing on X_α . The differentials d, ν are given by the formulas:

$$\begin{aligned} d(D \otimes \frac{d}{dz} \otimes dz \otimes v_\emptyset) &= D \cdot \frac{d}{dz} \otimes dz \otimes v_\emptyset - D \otimes 1 \otimes A_{\alpha, \emptyset}(v_\emptyset), \\ d(D \otimes z \otimes 1 \otimes v_\alpha) &= D \cdot z \otimes 1 \otimes v_\alpha - D \otimes dz \otimes A_{\emptyset, \alpha}(v_\alpha), \\ \nu(D \otimes 1 \otimes v_\alpha) &= D \cdot (1 \otimes v_\alpha), \quad \nu(D \otimes dz \otimes v_\emptyset) = D \cdot (dz \otimes v_\emptyset), \end{aligned}$$

for any $D \in \mathcal{D}_X$, $v_\emptyset \in V_\emptyset$, $v_\alpha \in V_\alpha$. Here $D \cdot (1 \otimes v_\alpha)$ and $D \cdot (dz \otimes v_\emptyset)$ denote the application of D to elements of $E\mathcal{V}$.

The de Rham functor transforms holonomic $\mathcal{D}_X$ -modules to perverse sheaves [K2, Mb], see also [B]. The application of the de Rham functor to the complex $\mathcal{R}E\mathcal{V}_\Gamma$ gives an explicit description of the associated perverse sheaf $\mathcal{Q}\mathcal{V}$, as a complex of free $\mathcal{O}_X$ -modules.

Example 1.7. In Example 1.2, the perverse sheaf $\mathcal{Q}\mathcal{V}$, associated with $E\mathcal{V}$, is given by the complex

$$\mathcal{Q}\mathcal{V} : 0 \rightarrow \mathcal{Q}\mathcal{V}^{-1} \xrightarrow{d} \mathcal{Q}\mathcal{V}^0 \rightarrow 0.$$

Here

$$\mathcal{Q}\mathcal{V}^{-1} = \Omega_X^{an} \otimes T_\emptyset \otimes \overline{V}_\emptyset \oplus \Omega_X^{an} \otimes F_\alpha \otimes \overline{V}_\alpha,$$

and

$$\mathcal{Q}\mathcal{V}^0 = \Omega_X^{an} \otimes \overline{V}_\emptyset \oplus \Omega_X^{an} \otimes \overline{V}_\alpha,$$

where Ω_X^{an} denotes the space of holomorphic one-forms on X . The differential d is given by the formulas:

$$\begin{aligned} d(\omega \otimes \frac{d}{dz} \otimes dz \otimes v_\emptyset) &= -Lie_{\frac{d}{dz}}(\omega) \otimes dz \otimes v_\emptyset - \omega \otimes 1 \otimes A_{\alpha, \emptyset}(v_\emptyset), \\ d(\omega \otimes z \otimes 1 \otimes v_\alpha) &= z\omega \otimes 1 \otimes v_\alpha - \omega \otimes dz \otimes A_{\emptyset, \alpha}(v_\alpha), \end{aligned}$$

for any $\omega \in \Omega_X^{an}$, $v_\emptyset \in V_\emptyset$, $v_\alpha \in V_\alpha$. Here $Lie_{\frac{d}{dz}}(\omega)$ denotes the Lie derivative of ω in the direction of the vector field $\frac{d}{dz}$.

Let $\Delta_X : \mathcal{M}_X^{hol} \rightarrow \mathcal{M}_X^{hol}$ and $\Delta_{X^0} : \mathcal{M}_{X^0}^{hol} \rightarrow \mathcal{M}_{X^0}^{hol}$ be the duality functors on holonomic $\mathcal{D}$ -modules [K1, KK], see also [K4, B]. For any $k = 0, 1, \dots, N$, we define an anti-automorphism τ_k of the category Qui_{Γ^k} . We establish an isomorphism $\Delta_{X^0}(E^0(\mathcal{V}_{\Gamma^0})) \approx E^0(\tau_0(\mathcal{V}_{\Gamma^0}))$ of $\mathcal{D}_{X^0}$ -modules for any level zero quiver $\mathcal{V}_{\Gamma^0}$ and an isomorphism $\Delta_X(E(\mathcal{V}_\Gamma)) \approx E(\tau_N(\mathcal{V}_\Gamma))$ of $\mathcal{D}_X$ -modules for any level N quiver $\mathcal{V}_\Gamma$.

Example 1.8. In Example 1.3, for any level zero quiver $\mathcal{V}_{\Gamma^0} = \{V_\emptyset, A_\emptyset^\emptyset\}$, the quiver $\tau_0 \mathcal{V}_{\Gamma^0}$ is $\{U_\emptyset, B_\emptyset^\emptyset\}$, where $U_\emptyset = V_\emptyset^*$ is the dual space and $B_\emptyset^\emptyset = (A_\emptyset^\emptyset)^t$ is the transpose operator.

In the theory of $\mathcal{D}$ -modules, there are two functors of direct image, $j_{l,k,!} : \mathcal{M}_{X^k}^{hol} \rightarrow \mathcal{M}_{X^l}^{hol}$, and $j_{l,k,*} : \mathcal{M}_{X^k}^{hol} \rightarrow \mathcal{M}_{X^l}^{hol}$, associated to the inclusion $j_{l,k} : X^k \rightarrow X^l$, $k < l$. In addition to the functor $\mathbb{J}_{l,k,*} : Qui_{\Gamma^k} \rightarrow Qui_{\Gamma^l}$, we introduce the second functor of quiver direct image $\mathbb{J}_{l,k,!} : Qui_{\Gamma^k} \rightarrow Qui_{\Gamma^l}$, and construct an isomorphisms of $\mathcal{D}_X$ modules $\mathbb{J}_{N,0,!} \mathcal{V}_{\Gamma^0}$ and $\tau_N \mathbb{J}_{N,0,*} \tau_0 \mathcal{V}_{\Gamma^0}$ for a non-resonant quiver $\mathbb{J}_{N,0,!} \mathcal{V}_{\Gamma^0}$.

Example 1.9. In Example 1.3, for any level zero quiver $\mathcal{V}_{\Gamma^0} = \{V_\emptyset, A_\emptyset^\emptyset\}$, the quiver $\mathbb{J}_{1,0,!}(\mathcal{V}_{\Gamma^0})$, is $\{W'_\emptyset, W'_\alpha, C'_{\emptyset,\alpha}, C'_{\alpha,\emptyset}\}$, where $W'_\emptyset = V_\emptyset$, $W'_\alpha = V_\alpha$, $C'_{\alpha,\emptyset} = \text{Id}_{V_\emptyset}$, $C'_{\emptyset,\alpha} = A_\emptyset^\emptyset$. It is non-resonant if and only if the quiver $\mathbb{J}_{1,0,*}(\mathcal{V}_{\Gamma^0})$ is non-resonant.

Let $z_1, \dots, z_N$ be linear coordinates on $X = \mathbb{C}^N$ and $\xi_1, \dots, \xi_N$ the dual coordinates on the dual space X^* . The assignment $\xi_i \mapsto -\frac{\partial}{\partial z_i}$, $\frac{\partial}{\partial \xi_i} \mapsto z_i$ for $i = 1, \dots, N$, defines an isomorphism of the rings D_{X^*} and D_X called *the Fourier transform*. The Fourier transform defines a functor from the category $\mathcal{M}_X$ to the category $\mathcal{M}_{X^*}$.

We consider a central arrangement $\mathcal{C}$ of hyperplanes in X , a quiver $\mathcal{V}$ of $\mathcal{C}$ and the associated quiver $\mathcal{D}_X$ -module $E\mathcal{V}$. We describe the Fourier transform of $E\mathcal{V}$. It turns out that the Fourier transform of $E\mathcal{V}$ is the quiver $\mathcal{D}_{X^*}$ -module associated with a suitable arrangement in X^* .

Example 1.10. In Example 1.1, the space X^* has two strata: $X_\emptyset^* = \{\xi = 0\}$ and $X_\alpha^* = \{\xi \neq 0\}$. The Fourier transform of the $\mathcal{D}_X$ -module $E\mathcal{V}$ from Example 1.2 is generated by vectors $1 \otimes v_\emptyset$, where $v_\emptyset \in V_\emptyset$, and $d\xi \otimes v_\alpha$, where $v_\alpha \in V_\alpha$. The relations are: $\xi(1 \otimes v_\emptyset) = -d\xi \otimes A_{\alpha,\emptyset}(v_\emptyset)$ for any $v_\emptyset \in V_\emptyset$ and $\frac{d}{d\xi}(d\xi \otimes v_\alpha) = 1 \otimes A_{\emptyset,\alpha}(v_\alpha)$ for any $v_\alpha \in V_\alpha$.

In the theory of perverse sheaves there is a notion of the MacPherson extension [BBD, GMc]. In our situation it is a morphism from the category of perverse sheaves on X^0 to the category of perverse sheaves on X . A perverse sheaf on X^0 is a local system on X^0 . Thus the MacPherson extension is an extension of a local system on X^0 to a suitable perverse sheaf on X .

The MacPherson extension of perverse sheaves corresponds to a morphism $j_{N,0,!} : \mathcal{M}_{X^0}^{hol} \rightarrow \mathcal{M}_X^{hol}$ of holonomic $\mathcal{D}$ -modules. The morphism $j_{N,0,!}$ can be defined in two steps. First, one defines a morphism $s_0^{(D)} : j_{N,0,!}(M) \rightarrow j_{N,0,*}(M)$ for any $\mathcal{D}_X$ -module M and then one defines $j_{N,0,!}(M)$ as the image of $j_{N,0,!}(M)$ in $j_{N,0,*}(M)$ under the morphism $s_0^{(D)}$. The MacPherson extension of the perverse sheaf associated to the $\mathcal{D}_{X^0}$ -module M , is the perverse sheaf, associated to the $\mathcal{D}_X$ -module $j_{N,0,!}(M)$.

In this paper we construct a morphism $s_0 : \mathbb{J}_{N,0,!} \rightarrow \mathbb{J}_{N,0,*}$ of functors of quiver categories. We introduce the quiver MacPherson extension of a level zero quiver $\mathcal{V}_{\Gamma^0}$ as the image of $\mathbb{J}_{N,0,!}(\mathcal{V}_{\Gamma^0})$ under the map s_0 . We prove that for a non-resonant level zero quiver $\mathcal{V}_{\Gamma^0}$, the $\mathcal{D}_X$ -module morphisms $Es_0 : E\mathbb{J}_{N,0,!}(\mathcal{V}_{\Gamma^0}) \rightarrow E\mathbb{J}_{N,0,*}(\mathcal{V}_{\Gamma^0})$ and $s_0^{(D)} E^0 : j_{N,0,!}(E^0 \mathcal{V}_{\Gamma^0}) \rightarrow j_{N,0,*}(E^0 \mathcal{V}_{\Gamma^0})$ are equal under the identifications of $j_{0,N,!}(E^0 \mathcal{V}_{\Gamma^0})$ with $E(\mathbb{J}_{0,N,!} \mathcal{V}_{\Gamma^0})$, and $j_{0,N,*}(E^0 \mathcal{V}_{\Gamma^0})$ with $E(\mathbb{J}_{0,N,*} \mathcal{V}_{\Gamma^0})$, mentioned above. Thus we have the

isomorphism of $\mathcal{D}_X$ -modules: $j_{N,0,!}(E\mathcal{V}_{\Gamma_0}) \approx E_{\mathbb{J}_{N,0,!}}(\mathcal{V}_{\Gamma_0})$, which relates the geometric and quiver MacPherson extensions.

Example 1.11. In Examples 1.4 and 1.9, the morphism s_0 of the level one quiver $\mathbb{J}_{1,0,!}\mathcal{V}_{\Gamma^0}$ to the level one quiver $\mathbb{J}_{1,0,*}\mathcal{V}_{\Gamma^0}$ is the collection of two linear maps, $(s_0)_\emptyset : W'_\emptyset \rightarrow W_\emptyset$ and $(s_0)_\alpha : W'_\alpha \rightarrow W_\alpha$. They are $(s_0)_\emptyset = Id_{V_\emptyset}$, and $(s_0)_\alpha = A_\emptyset^\emptyset$. The level one quiver $\mathbb{J}_{1,0,!}\mathcal{V}_{\Gamma^0}$ is $\{W''_\emptyset, W''_\alpha, C''_{\emptyset,\alpha}, C''_{\alpha,\emptyset}\}$, where $W''_\emptyset = V_\emptyset$, $W''_\alpha = \text{Im } A_\emptyset^\emptyset \subset V_\alpha$, $C''_{\alpha,\emptyset} = A_\emptyset^\emptyset$, and $C''_{\emptyset,\alpha}$ is the inclusion of $\text{Im } A_\emptyset^\emptyset$ to $V_\emptyset$.

As an application of quiver constructions, we calculate the cohomology groups of X (and X^0) with coefficients in a quiver perverse sheaf if the arrangement $\mathcal{C}$ is central and all linear maps of the quiver are close to zero. Namely, for a quiver $\mathcal{V} = \{V_\alpha, A_{\alpha,\beta}\} \in \text{Qui}_\Gamma$, we construct a finite-dimensional complex in terms of the spaces V_α and linear maps $A_{\alpha,\beta}$ and show that this complex calculates the cohomology groups of X with coefficients in the quiver perverse sheaf associated with $\mathcal{V}$.

Example 1.12. In Examples 1.1 and 1.2, consider the perverse sheaf $\mathcal{QV}$ on X , associated with the $\mathcal{D}_X$ -module $E\mathcal{V}$. If the operators $A_{\alpha,\emptyset}$ and $A_{\emptyset,\alpha}$ are close to zero, then the cohomology groups of X with coefficients in the perverse sheaf $\mathcal{QV}$ are calculated by the complex

$$0 \rightarrow V_\emptyset \xrightarrow{A_{\alpha,\emptyset}} V_\alpha \rightarrow 0,$$

where $\deg V_\emptyset = -1$, and $\deg V_\alpha = 0$.

Example 1.13. In Example 1.3, consider the perverse sheaf $\mathcal{Q}^0\mathcal{V}^0$ on X^0 , associated with the $\mathcal{D}_{X^0}$ -module $E^0\mathcal{V}_{\Gamma^0}$. If the operator $A_\emptyset^\emptyset$ is close to zero, then the cohomology groups of X^0 with coefficients in $\mathcal{Q}^0\mathcal{V}^0$, are calculated by the complex

$$0 \rightarrow V_\emptyset \xrightarrow{A_\emptyset^\emptyset} V_\emptyset \rightarrow 0.$$

Example 1.14. In Example 1.11, consider the perverse sheaf $\mathcal{Q}^0\mathcal{V}^0$ on X^0 . If the operator $A_\emptyset^\emptyset$ is close to zero, then the cohomology groups of X with coefficients in the MacPherson extension of $\mathcal{Q}^0\mathcal{V}^0$ are calculated by the complex

$$0 \rightarrow V_\emptyset \xrightarrow{A_\emptyset^\emptyset} \text{Im } A_\emptyset^\emptyset \rightarrow 0.$$

3. All results of the paper admit an equivariant extension to the case when the arrangement $\mathcal{C}$ admits a finite symmetry group.

As an application of equivariant constructions, we consider in Section 6.6 the discriminantal arrangements and one-dimensional local systems appearing in hypergeometric solutions of the KZ equations [SV]. We calculate the equivariant intersection cohomology groups of the discriminantal arrangements and formulate the result in terms of representation theory in Corollary 6.11. The statement of Corollary 6.11 is one of the main motivations of this paper.

4. In some of the statements of the paper we impose two types of non-resonance conditions on a quiver $\mathcal{V}_\Gamma = \{V_\alpha, A_{\alpha,\beta}\}$. For every vertex $\alpha \in \Gamma$, we construct a finite

collection of auxiliary linear operators of the form $\sum A_{\alpha,\beta} A_{\beta,\alpha} : V_\alpha \rightarrow V_\alpha$ with suitable range of summation over β . In the first type of non-resonance conditions, we require that the auxiliary operators have no eigenvalues which differ by an integer (the weak non-resonance condition). In the second type of non-resonance conditions, we require that the auxiliary operators have no eigenvalues which differ by an integer and zero is the only admissible integer eigenvalue (the non-resonance condition). The importance of non-resonance conditions is illustrated in the Example 1.13 below.

Example 1.13. Under conditions of Example 1.3, let $\mathcal{V}_{\Gamma^0} = \{V_\emptyset, A_\emptyset^\emptyset\}$ and $\mathcal{W}_{\Gamma^0} = \{W_\emptyset, B_\emptyset^\emptyset\}$ be two level zero quivers and $E^0\mathcal{V}_{\Gamma^0}$ and $E^0\mathcal{W}_{\Gamma^0}$ the associated D_{X^0} -modules. Assume that $G \subset \mathbb{C}$ is a subset, such that $x - y \notin \mathbb{Z} \setminus \{0\}$ for all $x, y \in G$. Assume that eigenvalues of $A_\emptyset^\emptyset$ and $B_\emptyset^\emptyset$ belong to G . Under these assumptions we will construct an isomorphism of vector spaces $\text{Hom}_{D_{X^0}}(E^0\mathcal{V}_{\Gamma^0}, E^0\mathcal{W}_{\Gamma^0})$ and $\text{Hom}_{\text{Quiv}_{\Gamma^0}}(\mathcal{V}_{\Gamma^0}, \mathcal{W}_{\Gamma^0})$. Indeed, let $\Phi : E^0\mathcal{V}_{\Gamma^0} \rightarrow E^0\mathcal{W}_{\Gamma^0}$ be a morphism of D_{X^0} -modules. The morphism Φ is uniquely determined by its restriction $\varphi = \Phi|_{\overline{V}_\emptyset}$ to the space $\overline{V}_\emptyset \subset E^0\mathcal{V}_{\Gamma^0}$. The map φ is a regular rational function on X^0 with values in $\text{Hom}(\overline{V}_\emptyset, \overline{W}_\emptyset)$. Write its Laurent series,

$$\varphi = (1 \otimes \varphi_k)z^k + (1 \otimes \varphi_{k+1})z^{k+1} + \dots,$$

where $\varphi_l : V_\emptyset \rightarrow W_\emptyset$ are linear maps and $\varphi_k \neq 0$. The condition $[\Phi, \frac{d}{dz}] = 0$ implies $\varphi_k A_\emptyset^\alpha = (B_\emptyset^\alpha + k)\varphi_k$. Since all eigenvalues of $A_\emptyset^\alpha$ and $B_\emptyset^\alpha$ are in G , we conclude that $k = 0$. The leading term φ_0 is a morphism of quivers, $\mathcal{V}_{\Gamma^0} \rightarrow \mathcal{W}_{\Gamma^0}$. Let $E^0(\varphi_0) : E^0\mathcal{V}_{\Gamma^0} \rightarrow E^0\mathcal{W}_{\Gamma^0}$ be the D_{X^0} -module morphism associated with φ_0 . The restriction of $E^0(\varphi_0)$ to $\overline{V}_\emptyset$ is $1 \otimes \varphi_0$. The leading term $\tilde{\varphi}_k$ of the D_{X^0} -module morphism $\tilde{\Phi} = \Phi - E^0(\varphi_0)$ has $k > 0$. Hence $\tilde{\Phi} = 0$ and $\Phi = E^0(\varphi_0)$.

5. In the theory of arrangements, two complexes play an important role. The first is the Aomoto complex [A] on the space of the Orlik-Solomon algebra and the second is the flag complex on the dual space to the Orlik-Solomon algebra [SV]. The Shapovalov form of [SV] defines a morphism of the flag complex to the Aomoto complex. The image of the Shapovalov form is called the complex of flag forms. The cohomology groups of the flag complex give the cohomology groups of X^0 with compact support. The cohomology groups of the Aomoto complex give the cohomology groups of X^0 with coefficients in a suitable local system on X^0 . An important problem is to give a topological meaning to the complex of flag forms.

The main quiver constructions of this paper are the constructions of the two quiver direct images $\mathbb{J}_{N,0,*}$ and $\mathbb{J}_{N,0,!}$. The constructions of $\mathbb{J}_{N,0,*}$ and $\mathbb{J}_{N,0,!}$ are respective generalizations of the constructions of the Aomoto complex and the flag complex. The morphism of functors $s_0 : \mathbb{J}_{N,0,!} \rightarrow \mathbb{J}_{N,0,*}$ is a generalization of the Shapovalov morphism of the flag complex to the Aomoto complex, see Section 3.6. The quiver MacPherson extension $\mathbb{J}_{N,0,!,*}$ is a generalization of the complex of flag forms in [SV].

From this point of view, our calculation of the cohomology groups of X with coefficients in the MacPherson extension of a quiver perverse sheaf on X^0 gives a topological meaning to the complex of flag forms.

To find a topological meaning of the complex of flag forms was one of the main goals of this paper.

6. The plan of the paper is as follows. In Section 2 we introduce graphs and stratified spaces, associated with an arrangement of hyperplanes. In Section 3 we study quivers, assigned to the arrangement, and operations on quivers. Section 4 is devoted to quiver $\mathcal{D}$ -modules. In Section 5 we describe perverse sheaves related to quiver $\mathcal{D}$ -modules and calculate the cohomology groups of the affine space with coefficients in these sheaves. In Section 6 we study the equivariant version of quiver constructions. All proofs are collected in Section 7.

7. We thank V. Schechtman, who inspired and initiated the study of $\mathcal{D}$ -modules associated with an arrangement of hyperplanes [Kh, KhS1] and who suggested an idea of the construction of a free $\mathcal{D}$ -module resolution of a quiver $\mathcal{D}$ -module, see an announcement in [KhS2].

We thank the referee for many helpful suggestions.

The work of S.Kh. was supported by CRDF grant RM1-2334, the grants INTAS-OPEN-03-51-3350, RFBR grant 04-01-00642 and RFBR grant to support scientific schools NSh-1999.2003.2. The work of A.V. was supported by NSF grant DMS-0555327.

2. ARRANGEMENTS

2.1. Graph of an arrangement. Let $\mathcal{C} = \{H_j\}$, $j \in J(\mathcal{C})$, be an arrangement of finitely many hyperplanes in the complex affine space $X = \mathbb{C}^N$. Hyperplanes H_j define in X the structure of a stratified space which we denote by the same letter X or by $X^{\mathcal{C}}$. The closure of a stratum $X_\alpha \subset \mathbb{C}^N$ is the intersection of some hyperplanes H_j , $j \in J_\alpha \subset J(\mathcal{C})$, and the stratum X_α itself is the complement in $\overline{X}_\alpha$ to the union of the closures of all smaller strata contained in $\overline{X}_\alpha$:

$$\overline{X}_\alpha = \bigcap_{j \in J_\alpha \subset J(\mathcal{C})} H_j, \quad X_\alpha = \overline{X}_\alpha \setminus \left(\bigcup_{\overline{X}_\beta \subset \overline{X}_\alpha, \overline{X}_\beta \neq \overline{X}_\alpha} \overline{X}_\beta \right).$$

We denote by $X_\emptyset$ the complement to the union of all hyperplanes H_j which is the unique stratum of X open in X .

The combinatorial structure of the adjacency of strata can be described by a non-oriented graph $\Gamma = \Gamma(\mathcal{C})$. The vertices $\alpha \in I(\Gamma)$ of the graph are in one-to-one correspondence with the strata X_α of X . The edges $(\alpha, \beta) \in E(\Gamma)$ describe adjacency in codimension one:

$$(\alpha, \beta) \in E(\Gamma) \Leftrightarrow \begin{cases} \overline{X}_\alpha \supset \overline{X}_\beta, & \text{codim}_{\overline{X}_\alpha} \overline{X}_\beta = 1, \\ \overline{X}_\beta \supset \overline{X}_\alpha, & \text{codim}_{\overline{X}_\beta} \overline{X}_\alpha = 1. \end{cases} \quad \text{or}$$

Two vertices α and β will be called *adjacent*, if $(\alpha, \beta) \in E(\Gamma)$.

To any vertex α we attach a nonnegative integer $l(\alpha)$,

$$l(\alpha) = \text{codim}_{\mathbb{C}^N} X_\alpha.$$

We write $\alpha \succ \beta$ for vertices α, β if (α, β) is an edge and $l(\alpha) < l(\beta)$. In other words,

$$\alpha \succ \beta \Leftrightarrow \overline{X}_\alpha \supset \overline{X}_\beta, \quad \text{codim}_{\overline{X}_\alpha} \overline{X}_\beta = 1.$$

We set

$$\varepsilon(\alpha, \beta) = \begin{cases} 1 & \text{if } \alpha \succ \beta, \\ -1 & \text{if } \beta \succ \alpha, \\ 0 & \text{otherwise.} \end{cases}$$

For an integer $n \geq 0$ denote by $I_n(\Gamma) \subset I(\Gamma)$ the subset of all vertices α such that $l(\alpha) = n$. We have $I_0(\Gamma) = \emptyset$, $I_1(\Gamma) = J(\mathcal{C})$.

We write $\alpha > \beta$ for $\alpha, \beta \in I(\Gamma)$ if there exists a chain $\alpha_1 \succ \alpha_2 \succ \dots \succ \alpha_n$ with $\alpha_1 = \alpha$ and $\alpha_n = \beta$. We write $\alpha \geq \beta$ if $\alpha > \beta$ or $\alpha = \beta$. Clearly,

$$\alpha \geq \beta \iff \overline{X}_\alpha \supset \overline{X}_\beta.$$

The labeled graph $\Gamma = \{I(\Gamma), E(\Gamma), l : I(\Gamma) \rightarrow \mathbb{Z}_{\geq 0}\}$ has the following properties.

- (i) There exists the unique vertex $\emptyset \in I(\Gamma)$ such that $l(\emptyset) = 0$ and $\emptyset \geq \beta$ for any $\beta \in I(\Gamma)$;
- (ii) If $(\alpha, \beta) \in E(\Gamma)$, then $|l(\alpha) - l(\beta)| = 1$;
- (iii) For any $\alpha, \beta \in I(\Gamma)$ the set $\Lambda(\alpha, \beta) = \{\gamma \in I(\Gamma) : \alpha \geq \gamma, \beta \geq \gamma\}$ is either empty or has the unique element γ maximal with respect to the partial order $\geq$. Moreover, we have $l(\gamma) \leq l(\alpha) + l(\beta)$. This element is called *the intersection of α and β* and denoted $\alpha \wedge \beta$.
- (iv) For any $\alpha, \beta \in I(\Gamma)$ with non-empty $\Lambda(\alpha, \beta)$, denote $U(\alpha, \beta) = \{\delta \in I(\Gamma) : \delta \geq \alpha, \delta \geq \beta\}$. Then for any $\delta \in U(\alpha, \beta)$ we have $l(\delta) \leq l(\alpha) + l(\beta) - l(\alpha \wedge \beta)$. Moreover, if there is $\delta \in U(\alpha, \beta)$ such that $l(\delta) = l(\alpha) + l(\beta) - l(\alpha \wedge \beta)$, then δ is the unique minimal element of $U(\alpha, \beta)$ with respect to the partial order $\geq$.

One can see that if $\beta \wedge \gamma$ and $\alpha \wedge (\beta \wedge \gamma)$ are defined, then $\alpha \wedge \beta$ and $(\alpha \wedge \beta) \wedge \gamma$ are defined, and $\alpha \wedge (\beta \wedge \gamma) = (\alpha \wedge \beta) \wedge \gamma$.

Any labeled graph with properties (i-iv) will be called *admissible*.

2.2. Framings of an arrangement. Let $\mathcal{C} \subset \mathbb{C}^N$ be an arrangement. To each stratum $X_\alpha \subset X^{\mathcal{C}}$ we attach certain spaces of affine functions, constant vector fields, and the top exterior forms on $\overline{X}_\alpha$ with constant coefficients.

Denote by F the space of affine functions on $\mathbb{C}^N$,

$$f \in F \iff f = a_0 + \sum_{i=1}^N a_i z_i, \quad a_i \in \mathbb{C},$$

where $z_1, \dots, z_N$ are affine coordinates on $\mathbb{C}^N$.

Denote by T the space of vector fields ξ on $\mathbb{C}^N$ with constant coefficients,

$$\xi \in T \iff \xi = \sum b_i \frac{\partial}{\partial z_i}, \quad b_i \in \mathbb{C}.$$

For a stratum X_α denote by $F_\alpha \subset F$ the subspace of all functions vanishing on X_α and by $T_\alpha \subset T$ the subspace of all vector fields parallel to X_α , that is, $\xi \in T_\alpha$ if $\langle \xi, f \rangle = 0$ for any $f \in F_\alpha$. Here $\langle \xi, f \rangle = i_\xi(df)$ denotes the substitution of the vector field ξ to the differential form df .

A function $f \in F_\alpha$ is called *generic*, if for any stratum X_β the intersection

$$\{f = 0\} \cap X_\beta$$

has the minimal possible dimension among all $f \in F_\alpha$.

Denote by $\mathcal{T}_\alpha$ the N -dimensional complex vector space,

$$\mathcal{T}_\alpha = T_\alpha \oplus F_\alpha .$$

Let X_α and X_β be strata, such that $\overline{X}_\alpha \supset X_\beta$. Then there are tautological inclusions

$$(2.1) \quad \mu_{\beta,\alpha} : F_\alpha \hookrightarrow F_\beta, \quad \overline{\mu}_{\alpha,\beta} : T_\beta \hookrightarrow T_\alpha ,$$

We denote

$$\mu_{\beta,\alpha} : \bigwedge^p F_\alpha \hookrightarrow \bigwedge^p F_\beta, \quad \overline{\mu}_{\alpha,\beta} : \bigwedge^p T_\beta \hookrightarrow \bigwedge^p T_\alpha, \quad p > 0,$$

the induced inclusions.

For $\alpha \succ \beta$ and $f \in F_\beta$, introduce the linear maps

$$i_f^{\beta,\alpha} : \bigwedge^p T_\alpha \rightarrow \bigwedge^{p-1} T_\beta, \quad \vec{\xi} \mapsto i_{df}(\vec{\xi}) .$$

Here $i_{df}(\vec{\xi})$ is the substitution of df to the p -vector $\vec{\xi} \in \bigwedge^p T_\alpha$ with the result considered as a $(p-1)$ -vector in $\bigwedge^{p-1} T_\beta$. More precisely, if $\vec{\xi} = \xi_1 \wedge \dots \wedge \xi_p$ and $\xi_2, \dots, \xi_p \in T_\beta$, then

$$(2.2) \quad i_f^{\beta,\alpha}(\vec{\xi}) = \langle \xi_1, f \rangle \xi_2 \wedge \dots \wedge \xi_p .$$

For $\alpha \succ \beta$ and $\xi \in T_\alpha$ introduce the linear map

$$i_\xi^{\alpha,\beta} : \bigwedge^p F_\beta \rightarrow \bigwedge^{p-1} F_\alpha, \quad \vec{f} \mapsto i_\xi(\vec{f}) .$$

More precisely, if $\vec{f} = f_1 \wedge \dots \wedge f_p$ with $f_2, \dots, f_p \in F_\alpha$, then

$$(2.3) \quad i_\xi^{\alpha,\beta}(\vec{f}) = \langle \xi, f_1 \rangle f_2 \wedge \dots \wedge f_p .$$

Denote by $\overline{\Omega}^p$ the complex vector space of holomorphic exterior p -forms on $\mathbb{C}^N$, invariant with respect to affine translations. Each form in $\overline{\Omega}^p$ is a linear combination of the forms $df_1 \wedge \dots \wedge df_p$ with $f_1, \dots, f_p \in F$.

For a stratum X_α , denote by $\overline{\Omega}_\alpha$ the one-dimensional vector space of the top exterior holomorphic forms on X_α invariant with respect to translations along X_α . The space $\overline{\Omega}_\alpha$ is isomorphic to the quotient of the space $\overline{\Omega}^{N-l(\alpha)}$ over the subspace generated by the forms $df_1 \wedge \dots \wedge df_{N-l(\alpha)}$, where $f_1, \dots, f_{N-l(\alpha)} \in F$ and at least one of them belongs to F_α .

It is technically convenient to fix a vector space structure on $\mathbb{C}^N$ and fix a non-degenerate bilinear form $(,)$ in this vector space $\mathbb{C}^N$. In this setting to each stratum X_α we also attach the following spaces T'_α and F'_α .

The space T'_α consists of constant vector fields orthogonal to T_α . Let $z_1, \dots, z_N$ be linear coordinates in $\mathbb{C}^N$ with respect to an orthonormal basis. Let $\xi' = \sum b_i \frac{\partial}{\partial z_i}$ be a constant vector field. Then ξ' belongs to T'_α , if $b_1 a_1 + \dots + b_N a_N = 0$ for any $\xi = \sum a_i \frac{\partial}{\partial z_i} \in T_\alpha$. The space F'_α consists of all linear functions $f' = \sum c_i z_i$, which are annihilated by vector fields in T'_α , i.e. $\langle \xi', f' \rangle = 0$ for any $\xi' \in T'_\alpha$.

We use two types of framings of the arrangement $\mathcal{C}$: edge framings and vertex framings.

An *edge framing* is a choice for any edge $(\alpha, \beta) \in E(\Gamma)$, $l(\alpha) + 1 = l(\beta)$, of a function $f_{\beta, \alpha} \in F_\beta \setminus F_\alpha$ and a vector field $\xi_{\alpha, \beta} \in T_\alpha \setminus T_\beta$. For any edge $(\alpha, \beta) \in E(\Gamma)$, the edge framing determines the linear map $\pi_{\beta, \alpha} : \overline{\Omega}_\alpha \rightarrow \overline{\Omega}_\beta$, such that for $\overline{\omega}_\alpha \in \overline{\Omega}_\alpha$ and $\overline{\omega}_\beta \in \overline{\Omega}_\beta$ we have

$$(2.4) \quad \begin{aligned} \pi_{\beta, \alpha}(\overline{\omega}_\alpha) = \overline{\omega}_\beta &\Leftrightarrow df_{\beta, \alpha} \wedge \overline{\omega}_\beta = \overline{\omega}_\alpha, \quad \text{if } \alpha \succ \beta, \\ \pi_{\beta, \alpha}(\overline{\omega}_\alpha) = \overline{\omega}_\beta &\Leftrightarrow i_{\xi_{\beta, \alpha}}(\overline{\omega}_\beta) = \overline{\omega}_\alpha, \quad \text{if } \beta \succ \alpha. \end{aligned}$$

A *vertex framing* of $\mathcal{C}$ is a choice of a function $\overline{f}_\alpha \in F_\alpha$ for any stratum X_α . A vertex framing $\mathcal{F}$ is called *generic* if for any $\alpha \in I(\Gamma)$ the function $\overline{f}_\alpha$ is generic.

For a nonnegative integer n , a *level n vertex framing* is a choice of a function $\overline{f}_\alpha \in F_\alpha$ for any $\alpha \in I_{n+1}(\Gamma)$. It is generic if every chosen $\overline{f}_\alpha$ is generic.

2.3. Principal open subsets X^n . For a nonnegative integer n and an arrangement $\mathcal{C}$, define the open subset $X^n \subset X$ as the complement to the closure of the union of all codimension $n + 1$ strata:

$$X^n = \mathbb{C}^N \setminus \bigcup_{\alpha, l(\alpha)=n+1} \overline{X}_\alpha.$$

We call X^n the *principal level n open subset* of X . In particular, X^0 coincides with $X_\emptyset$, the open stratum in X , and $X^N = X$. The sets X^n determine a filtration of X :

$$X^0 \subset X^1 \subset \dots \subset X^N = X,$$

For $k \leq l$ denote by $j_{l,k}$ the inclusion

$$j_{l,k} : X^k \rightarrow X^l.$$

We have $j_{k_3, k_1} = j_{k_3, k_2} j_{k_2, k_1}$ for $k_1 \leq k_2 \leq k_3$ and we use the notation j_0 for the inclusion $j_{N,0} : X^0 \rightarrow X = X^N$.

Each level n vertex framing $\mathcal{F}_n = \{\overline{f}_\alpha \mid \alpha \in I_{n+1}(\Gamma)\}$ determines an open subset $U_{\mathcal{F}_n} \subset X^n$,

$$(2.5) \quad U_{\mathcal{F}_n} = X \setminus \bigcup_{\alpha \in I_{n+1}(\Gamma)} \{\overline{f}_\alpha = 0\}.$$

The open subsets $U_{\mathcal{F}_n}$ for different choices of framings $\mathcal{F}_n$ form a cover of the space X^n by open affine subsets.

2.4. Truncated graphs. The sets X^n are stratified spaces; their strata are the strata X_α of X with $l(\alpha) \leq n$. We define the graph (with loops) of the stratified space X^n as follows. The vertices of this graph Γ^n are the vertices $\alpha \in I(\Gamma)$ such that $l(\alpha) \leq n$. The edges $(\alpha, \beta) \in E(\Gamma^n)$, which are not loops, coincide with the edges (α, β) of Γ , connecting the vertices of Γ^n . For any pair $\alpha \succ \beta$ with $l(\alpha) = n$, the graph Γ^n has a loop at the vertex α , which we denote by $(\alpha, \alpha)^\beta$.

The graph Γ^n will be called the *truncated graph at level n* associated to the admissible graph Γ .

Examples. The graph Γ^0 has one vertex and $|J(\mathcal{C})|$ loops. Every loop corresponds to a hyperplane of the initial arrangement $\mathcal{C}$. The graph Γ^N coincides with Γ .

2.5. Orlik-Solomon algebra and flag complex. Here we recall the construction of the Orlik-Solomon algebra and the flag complex of an arrangement, see [SV]. For an arrangement $\mathcal{C}$ in $\mathbb{C}^N$ define the complex vector spaces $\mathcal{A}^p(\mathcal{C})$, $p = 0, \dots, N$. For $p = 0$ set $\mathcal{A}^0(\mathcal{C}) = \mathbb{C}$. For $p \geq 1$ the vector space $\mathcal{A}^p(\mathcal{C})$ is generated by symbols $(H_{j_1}, \dots, H_{j_p})$, where $H_{j_k} \in \mathcal{C}$, such that

(i)

$$(H_{j_1}, \dots, H_{j_p}) = 0$$

if $H_{j_1}, \dots, H_{j_p}$ are not in general position, that is if their intersection $H_{j_1} \cap \dots \cap H_{j_p}$ is empty or its codimension is less than p ;

(ii)

$$(H_{j_{\sigma(1)}}, \dots, H_{j_{\sigma(p)}}) = \text{sgn}(\sigma) \cdot (H_{j_1}, \dots, H_{j_p})$$

for any permutation $\sigma \in S_p$;

(iii)

$$(2.6) \quad \sum_{k=1}^{p+1} (-1)^k (H_{j_1}, \dots, \widehat{H_{j_k}}, \dots, H_{j_{p+1}}) = 0$$

for any $(p+1)$ -tuple $H_{j_1}, \dots, H_{j_{p+1}}$ of hyperplanes in $\mathcal{C}$ which are not in general position and such that $H_{j_1} \cap \dots \cap H_{j_{p+1}} \neq \emptyset$.

The direct sum $\mathcal{A}(\mathcal{C}) = \bigoplus_{p=1}^N \mathcal{A}^p(\mathcal{C})$ is a differential graded skew commutative algebra with respect to multiplication

$$(H_{j_1}, \dots, H_{j_p}) \cdot (H_{j_{p+1}}, \dots, H_{j_{p+q}}) = (H_{j_1}, \dots, H_{j_p}, H_{j_{p+1}}, \dots, H_{j_{p+q}})$$

and the differential $\delta_{\mathcal{A}}: \mathcal{A}^p(\mathcal{C}) \rightarrow \mathcal{A}^{p-1}(\mathcal{C})$,

$$(2.7) \quad \delta_{\mathcal{A}} : (H_{j_1}, \dots, H_{j_p}) \mapsto \sum_{k=0}^p (-1)^{k-1} (H_{j_1}, \dots, \widehat{H_{j_k}}, \dots, H_{j_p}).$$

The algebra is called *the Orlik-Solomon algebra* of the arrangement $\mathcal{C}$.

The Orlik-Solomon algebra is determined by the admissible graph Γ only. For an admissible graph Γ , its Orlik-Solomon algebra $\mathcal{A}(\Gamma)$ can be defined as the algebra generated by the skew-symmetric symbols $(H_{j_1}, \dots, H_{j_k})$, where $j_k \in I_1(\Gamma)$. We set $(H_{j_1}, \dots, H_{j_p}) = 0$ if $H_{j_1}, \dots, H_{j_k}$ are not in general position, and $H_{j_1}, \dots, H_{j_k}$ are not in general position if the vertex $j_1 \wedge \dots \wedge j_k$ does not exist or its codimension is less than k . Relation (2.6) has to take place for any $(p+1)$ -tuple of vertices $j_i \in I_1(\Gamma)$ for which the intersection $j_1 \wedge \dots \wedge j_{p+1}$ exists and its codimension is less than $p+1$.

For $\alpha \in I(\Gamma)$, a *flag starting at X_α* is a sequence

$$\overline{X}_{\alpha_0} \supset \dots \supset \overline{X}_{\alpha_p} = \overline{X}_\alpha$$

of closed strata such that $\text{codim}_{\mathbb{C}^N} X_{\alpha_k} = j$ for $j = 0, \dots, p$.

For $\alpha \in I(\Gamma)$, we define $\widetilde{\mathcal{F}}_\alpha$ to be the complex vector space with basis vectors $\widetilde{F}_{\alpha_0, \dots, \alpha_p = \alpha}$ labeled by the elements of the set of all flags starting at X_α .

Define $\mathcal{F}_\alpha$ as the quotient of $\tilde{\mathcal{F}}_\alpha$ over the subspace generated by the elements

$$(2.8) \quad \sum_{\beta, \alpha_{k-1} \succ \beta \succ \alpha_{k+1}} \tilde{F}_{\alpha_0, \dots, \alpha_{k-1}, \beta, \alpha_{k+1}, \dots, \alpha_p = \alpha} ,$$

where $k = 1, \dots, p-1$ and $\overline{X}_{\alpha_0} \supset \dots \supset \overline{X}_{\alpha_{k-1}} \supset \overline{X}_{\alpha_{k+1}} \supset \dots \supset \overline{X}_{\alpha_p} = \overline{X}_\alpha$ is any incomplete flag with $l(\alpha_j) = j$.

Denote by $F_{\alpha_0, \dots, \alpha_p}$ the image in $\mathcal{F}_\alpha$ of the basis vector $\tilde{F}_{\alpha_0, \dots, \alpha_p}$. Set

$$\mathcal{F}^p(\mathcal{C}) = \bigoplus_{\alpha \in \Gamma, l(\alpha)=p} \mathcal{F}_\alpha .$$

The direct sum

$$\mathcal{F}(\mathcal{C}) = \bigoplus_{p=0}^N \mathcal{F}^p(\mathcal{C})$$

is a complex with respect to the differential

$$(2.9) \quad d_{\mathcal{F}} : \mathcal{F}^p(\mathcal{C}) \rightarrow \mathcal{F}^{p+1}(\mathcal{C}), \quad F_{\alpha_0, \dots, \alpha_p} \mapsto (-1)^p \sum_{\alpha_{p+1}, \alpha_p \succ \alpha_{p+1}} F_{\alpha_0, \dots, \alpha_p, \alpha_{p+1}} .$$

The complex $(\mathcal{F}^\bullet(\mathcal{C}), d_{\mathcal{F}})$ is called *the flag complex of the arrangement $\mathcal{C}$* .

As well as the Orlik-Solomon algebra, it is determined by the graph $\Gamma = \Gamma(\mathcal{C})$ only and will be denoted in this context by $(\mathcal{F}^\bullet(\Gamma), d_{\mathcal{F}})$.

The vector spaces $\mathcal{A}^p(\mathcal{C})$ and $\mathcal{F}^p(\mathcal{C})$ are dual [SV]. The pairing $\mathcal{A}^p(\mathcal{C}) \otimes \mathcal{F}^p(\mathcal{C}) \rightarrow \mathbb{C}$ is defined as follows. For $H_{j_1}, \dots, H_{j_p}$ in general position, set

$$F(H_{j_1}, \dots, H_{j_p}) = F_{\alpha_0, \alpha_1, \dots, \alpha_p}$$

where

$$\overline{X}_{\alpha_0} = \mathbb{C}^N, \quad \overline{X}_{\alpha_1} = H_{j_1}, \quad \overline{X}_{\alpha_2} = H_{j_1} \cap H_{j_2}, \quad \dots, \quad \overline{X}_{\alpha_p} = H_{j_1} \cap H_{j_2} \cap \dots \cap H_{j_p}.$$

Then set

$$\langle (H_{j_1}, \dots, H_{j_p}), F \rangle = \text{sign}(\sigma) ,$$

if $F = F(H_{j_{\sigma(1)}}, \dots, H_{j_{\sigma(p)}})$ for some $\sigma \in S_p$, and

$$\langle (H_{j_1}, \dots, H_{j_p}), F \rangle = 0$$

otherwise.

The differentials $\delta_{\mathcal{A}}$ and $d_{\mathcal{F}}$ are adjoint with respect to this duality and the map dual to the multiplication map in $\mathcal{A}(\mathcal{C})$ turns $\mathcal{F}(\mathcal{C})$ into a skew cocommutative differential graded coalgebra.

Let $a : \mathcal{C} \rightarrow \mathbb{C}$ be a map which assigns to each hyperplane H a complex number $a(H)$ called *the exponent* of the hyperplane. Set

$$\omega(a) = \sum_{H \in \mathcal{C}} a(H) H \in \mathcal{A}^1(\mathcal{C}) .$$

The multiplication by $\omega(a)$ defines a differential

$$d_a : \mathcal{A}^p(\mathcal{C}) \rightarrow \mathcal{A}^{p+1}(\mathcal{C}), \quad x \mapsto \omega(a) \cdot x ,$$

in the vector space of the Orlik-Solomon algebra $\mathcal{A}(\mathcal{C})$. The complex $(\mathcal{A}^\bullet(\mathcal{C}), d_a)$ is called *the Aomoto complex*.

The collection of exponents determines *the Shapovalov map*

$$S_a : \mathcal{F}(\mathcal{C}) \rightarrow \mathcal{A}(\mathcal{C}) ,$$

$$F_{\alpha_0, \alpha_1, \dots, \alpha_p} \mapsto \sum a(H_{j_1}) a(H_{j_2}) \dots a(H_{j_p}) (H_{j_1}, \dots, H_{j_p}) ,$$

where the sum is taken over all p -tuples $(H_{j_1}, \dots, H_{j_p})$ such that

$$H_{j_1} \supset \overline{X}_{\alpha_1}, \dots, H_{j_p} \supset \overline{X}_{\alpha_p} .$$

The image $S_a(F_{\alpha_0, \alpha_1, \dots, \alpha_p}) \in \mathcal{A}^p$ of a flag $F_{\alpha_0, \alpha_1, \dots, \alpha_p}$ is called a *flag form*.

According to [SV], the map S_a is a morphism of the complex $(\mathcal{F}^\bullet(\mathcal{C}), d_{\mathcal{F}})$ to the complex $(\mathcal{A}^\bullet(\mathcal{C}), d_a)$. The image of the flag complex in the Aomoto complex is called *the complex of flag forms* of $\mathcal{C}$. We denote it by $(\overline{\mathcal{F}}_a^\bullet(\mathcal{C}), d_a)$.

Identifying $\mathcal{A}(\mathcal{C})$ with $\mathcal{F}^*(\mathcal{C})$, we can consider the map S_a as a bilinear form on the vector space $\mathcal{F}(\mathcal{C})$. The bilinear form is symmetric and called *the Shapovalov form*.

3. QUIVERS

3.1. Quivers of an admissible graph. Let $\Gamma = \{I(\Gamma), E(\Gamma), l : I(\Gamma) \rightarrow \mathbb{Z}_{\geq 0}\}$ be an admissible graph.

A *quiver* $\mathcal{V}_\Gamma = \{V_\alpha, A_{\alpha, \beta}\}$ of Γ is a collection of finite dimensional complex vector spaces V_α , $\alpha \in I(\Gamma)$, and a collection of linear maps $A_{\alpha, \beta} : V_\beta \rightarrow V_\alpha$, $\alpha, \beta \in I(\Gamma)$, such that

- (a) $A_{\alpha, \beta} = 0$ if α and β are not adjacent;
- (b) $\sum_\beta A_{\alpha, \beta} A_{\beta, \gamma} = 0$ if $|l(\alpha) - l(\gamma)| = 2$;
- (c) $\sum_\beta A_{\alpha, \beta} A_{\beta, \gamma} = 0$ if $l(\alpha) = l(\gamma)$, $\alpha \neq \gamma$, and there exists $\delta \in I(\Gamma)$, such that $\alpha \succ \delta, \gamma \succ \delta$.

The possibly nonzero operators $A_{\alpha, \beta}$ of a quiver $\mathcal{V}_\Gamma$ correspond to oriented edges of the graph Γ . There may be only one or two summands in the left hand side of any relation of type (c) due to properties (iii) and (iv) of Γ in Section 2.1

Let $\{V_\alpha, A_{\alpha, \beta}\}, \{V'_\alpha, A'_{\alpha, \beta}\}$ be two quivers of Γ . A *morphism of quivers* is a collection of maps $\{f_\alpha : V_\alpha \rightarrow V'_\alpha\}$ such that $f_\alpha A_{\alpha, \beta} = A'_{\alpha, \beta} f_\beta$ for all α, β .

Denote by Qui_Γ the category of all quivers of Γ .

A quiver can be considered as a module over the path algebra B_Γ .

The path algebra of Γ is the associative algebra B_Γ with unit, generated by two sets of generators i_α , $\alpha \in I(\Gamma)$, and $a_{\alpha, \beta}$, where (α, β) runs through the set of pairs of adjacent vertices of Γ . The generators are subject to the relations:

- (i) $i_\alpha^2 = i_\alpha$ for all α ;
- (ii) $i_\alpha i_\beta = 0$ for all $\alpha \neq \beta$;
- (iii) $\sum_\alpha i_\alpha = 1$;
- (iv) Let $\delta_{\beta, \gamma}$ be the Kronecker symbol, then $a_{\alpha, \beta} i_\gamma = \delta_{\beta, \gamma} a_{\alpha, \beta}$, and $i_\gamma a_{\alpha, \beta} = \delta_{\gamma, \alpha} a_{\alpha, \beta}$ for all α, β, γ .
- (v) $\sum_\beta a_{\alpha, \beta} a_{\beta, \gamma} = 0$ for all α, γ with $|l(\alpha) - l(\gamma)| = 2$;
- (vi) $\sum_\beta a_{\alpha, \beta} a_{\beta, \gamma} = 0$ for all α, γ , such that $\alpha \neq \gamma$, $l(\alpha) = l(\gamma)$, and there exists δ such that $\alpha \succ \delta, \gamma \succ \delta$.

A quiver in Qui_Γ has a natural structure of a B_Γ -module. The category Qui_Γ is equivalent to the category of finite-dimensional B_Γ -modules.

For $\alpha, \beta \in I(\Gamma)$, let $\varepsilon(\beta, \alpha)$ be defined as in Section 2.1.

Let $\mathcal{V}_\Gamma = \{V_\alpha, A_{\alpha, \beta}\}$ be a quiver of Γ . Then the collection $\mathcal{W}_\Gamma = \{W_\alpha, B_{\alpha, \beta}\}$, where $W_\alpha = V_\alpha^*$ and $B_{\alpha, \beta} = \varepsilon(\beta, \alpha) A_{\beta, \alpha}^* : W_\beta \rightarrow W_\alpha$ determines a quiver $\tau(\mathcal{V}_\Gamma)$, which we call *the dual quiver*. The assignment $\mathcal{V}_\Gamma \mapsto \tau(\mathcal{V}_\Gamma)$ determines an anti-automorphism of the category Qui_Γ of the fourth order. Moreover, for any quiver $\mathcal{V}_\Gamma$ the quiver $\tau^2(\mathcal{V}_\Gamma)$ is isomorphic to $\mathcal{V}_\Gamma$. The isomorphism $\phi : \mathcal{V}_\Gamma \rightarrow \tau^2(\mathcal{V}_\Gamma)$ is given by the relation $\phi(v_\alpha) = (-1)^{l(\alpha)} v_\alpha$.

The linear maps $A_{\alpha, \beta}$ of a quiver $\mathcal{V}_\Gamma$ can be collected into two differentials, acting in the total space of the quiver, $C(\mathcal{V}_\Gamma) = \bigoplus_{\alpha \in I(\Gamma)} V_\alpha$. Namely, let

$$(3.1) \quad C_k(\mathcal{V}_\Gamma) = \bigoplus_{\alpha, l(\alpha)=k} V_\alpha, \quad C(\mathcal{V}_\Gamma) = \bigoplus_k C_k(\mathcal{V}_\Gamma).$$

For $v \in V_\alpha$, set

$$(3.2) \quad d(v) = \sum_{\beta, \alpha \succ \beta} A_{\beta, \alpha}(v), \quad \text{and} \quad \partial(v) = \sum_{\beta, \beta \succ \alpha} A_{\beta, \alpha}(v).$$

We have

$$\begin{aligned} d(C_k(\mathcal{V}_\Gamma)) &\subset C_{k+1}(\mathcal{V}_\Gamma), & d^2 &= 0, \\ \partial(C_k(\mathcal{V}_\Gamma)) &\subset C_{k-1}(\mathcal{V}_\Gamma), & \partial^2 &= 0. \end{aligned}$$

Denote the complex $(C(\mathcal{V}_\Gamma), d)$ by $C_+(\mathcal{V}_\Gamma)$ and the complex $(C(\mathcal{V}_\Gamma), \partial)$ by $C_-(\mathcal{V}_\Gamma)$.

For a vertex β and an edge (α, β) , denote by A_β^α the operator

$$(3.3) \quad A_\beta^\alpha = A_{\beta, \alpha} A_{\alpha, \beta} : V_\beta \rightarrow V_\beta.$$

Let $\mathcal{O}_\beta(\mathcal{V}_\Gamma)$ be the $\mathbb{C}$ -algebra of endomorphisms of V_β , generated by operators A_β^α , where α runs through the set of all vertices adjacent to β . We call $\mathcal{O}_\beta(\mathcal{V}_\Gamma)$ *the local algebra of the quiver at the vertex β* .

To a vertex β , $\beta \neq \emptyset$, we attach three operators S_β , T_β and $\bar{T}_\beta$,

$$(3.4) \quad S_\beta = \sum_{\alpha, \alpha \succ \beta} A_\beta^\alpha : V_\beta \rightarrow V_\beta,$$

$$(3.5) \quad T_\beta = \sum_{\alpha, \alpha'} A_{\alpha, \beta} A_{\beta, \alpha'} : \bigoplus_{\alpha, \alpha \succ \beta} V_\alpha \rightarrow \bigoplus_{\alpha, \alpha \succ \beta} V_\alpha,$$

where the summation is over all α, α' such that $\alpha \succ \beta$, $\alpha' \succ \beta$ and

$$(3.6) \quad \bar{T}_\beta = \sum_{\alpha, \alpha', \delta} A_{\alpha, \delta} A_{\delta, \alpha'} : \bigoplus_{\alpha, \alpha \succ \beta} V_\alpha \rightarrow \bigoplus_{\alpha, \alpha \succ \beta} V_\alpha,$$

where the summation is over all α, α', δ such that $\alpha \succ \beta$, $\alpha' \succ \beta$, and $\delta \succ \alpha$, $\delta \succ \alpha'$.

For a vertex β , $l(\beta) \neq N$, we set

$$(3.7) \quad \tilde{S}_\beta = \sum_{\alpha, \beta \succ \alpha} A_\beta^\alpha : V_\beta \rightarrow V_\beta.$$

We also set $S_\emptyset = 0 : V_\emptyset \rightarrow V_\emptyset$ and $\tilde{S}_\beta = 0 : V_\beta \rightarrow V_\beta$, if $l(\beta) = N$. We define the operator $S : \oplus_{\alpha \in I(\Gamma)} V_\alpha \rightarrow \oplus_{\alpha \in I(\Gamma)} V_\alpha$ as

$$(3.8) \quad S = \sum_{\alpha \in I(\Gamma)} (S_\alpha + \tilde{S}_\alpha) = \sum_{\alpha, \beta \in I(\Gamma)} A_{\alpha, \beta} A_{\beta, \alpha} .$$

Proposition 3.1.

- (i) The operator S_β is a central element in the local algebra $\mathcal{O}_\beta(\mathcal{V})$.
- (ii) Any nonzero eigenvalue of T_β is equal to an eigenvalue of the operator $S_\alpha + A_\alpha^\beta$ for some α , $\alpha \succ \beta$.
- (iii) Any nonzero eigenvalue of $\bar{T}_\beta$ is equal to an eigenvalue of the operator $S_\alpha + A_\alpha^\beta$ for some α , $\alpha \succ \beta$.

Proposition 3.2. Let $\mathcal{C}$ be a central arrangement (that is, all hyperplanes of $\mathcal{C}$ contain $0 \in \mathbb{C}^N$). Then

- (i) The operator $\tilde{S}_\beta$ is a central element in the local algebra $\mathcal{O}_\beta(\mathcal{V})$.
- (ii) For any $\alpha, \beta \in I(\Gamma)$, we have

$$(3.9) \quad (S_\alpha + \tilde{S}_\alpha) A_{\alpha, \beta} = A_{\alpha, \beta} (S_\beta + \tilde{S}_\beta) .$$

In particular, the element S , defined in (3.8), is central in the path algebra B_Γ .

Propositions 3.1 and 3.2 are proved in Section 7.

A subset $G \subset \mathbb{C}$ is called *non-resonant* if it satisfies the following two conditions:

- (i) The intersection $G \cap \mathbb{Z}$ is either $\{0\}$ or empty.
- (ii) For any x, y in G , we have $x - y \in \mathbb{C} \setminus \mathbb{Z}$.

A subset $G \subset \mathbb{C}$ is called *weakly non-resonant* if it satisfies condition (ii).

Suppose that for every vertex $\beta \in I(\Gamma)$, $\beta \neq \emptyset$, a non-resonant subset $G_\beta \subset \mathbb{C}$ is chosen. Denote by $Qui_{\Gamma, \{G_\beta\}}$ the category of all quivers $\{V_\alpha, A_{\alpha, \beta}\} \in Qui_\Gamma$ such that

- (a) for any $\beta \in I(\Gamma)$, $\beta \neq \emptyset$, the eigenvalues of the operator T_β belong to G_β ,
- (b) for any $\beta \in I(\Gamma)$, $\beta \neq \emptyset$, none of the eigenvalues of the operator $\bar{T}_\beta$ is a positive integer.

Suppose we have

- a non-resonant subset $G_\beta \subset \mathbb{C}$ for every vertex $\beta \in I(\Gamma)$, $\beta \neq \emptyset$,
- a weakly non-resonant set $\tilde{G}_\beta$ for every vertex $\beta \in I(\Gamma)$,
- a weakly non-resonant subset $G_{\alpha, \beta}$ for any edge $(\alpha, \beta) \in E(\Gamma)$, $\alpha \succ \beta$.

Denote by $Qui_{\Gamma, \{G_\beta, \tilde{G}_\beta, G_{\alpha, \beta}\}}$ the subcategory of all quivers $\{V_\alpha, A_{\alpha, \beta}\} \in Qui_\Gamma$, such that

- (a) for any $\beta \in I(\Gamma)$, $\beta \neq \emptyset$, the eigenvalues of the operator T_β belong to G_β and
- (c) for any $\beta \in I(\Gamma)$, the eigenvalues of the operator $\tilde{S}_\beta$ belong to $\tilde{G}_\beta$ and
- (d) for any edge $(\alpha, \beta) \in E(\Gamma)$, $\alpha \succ \beta$, the eigenvalues of the operator A_α^β belong to $G_{\alpha, \beta}$.

The categories $Qui_{\Gamma, \{G_\beta, \tilde{G}_\beta, G_{\alpha, \beta}\}}$ and $Qui_{\Gamma, \{G_\beta\}}$ are full abelian subcategories of the category Qui_Γ . The categories $Qui_{\Gamma, \{G_\beta, \tilde{G}_\beta, G_{\alpha, \beta}\}}$ and $Qui_{\Gamma, \{G_\beta\}}$ are closed under extensions.

We call a quiver *non-resonant* if it belongs to the subcategory $Qui_{\Gamma, \{G_\beta\}}$ for some collection $\{G_\beta\}$. We call a quiver *strongly non-resonant* if it belongs to the subcategory $Qui_{\Gamma, \{G_\beta, \tilde{G}_\beta, G_{\alpha, \beta}\}}$ for some collections $\{G_\beta\}$, $\{\tilde{G}_\beta\}$, and $\{G_{\alpha, \beta}\}$.

3.2. Quivers of level n . Suppose that Γ is an admissible graph, $n \in \{0, \dots, N\}$, and Γ^n is the corresponding truncated graph.

A quiver $\mathcal{V}_{\Gamma^n}$ of Γ^n (or a level n quiver) consists of a collection of finite dimensional vector spaces V_α , $\alpha \in I(\Gamma^n)$, a collection of linear maps $A_{\alpha, \beta} : V_\beta \rightarrow V_\alpha$, $\alpha, \beta \in I(\Gamma^n)$, and a collection of linear maps $A_\alpha^\beta : V_\alpha \rightarrow V_\alpha$, labeled by loops $(\alpha, \alpha)^\beta \in E(\Gamma^n)$. The operators are subject to the following relations:

- (i) $A_{\alpha, \beta} = 0$ if α, β are not adjacent in Γ ;
- (ii) $\sum_{\beta \in I(\Gamma^n)} A_{\alpha, \beta} A_{\beta, \gamma} = 0$ if $|l(\alpha) - l(\gamma)| = 2$;
- (iii) $\sum_{\beta \in I(\Gamma^n)} A_{\alpha, \beta} A_{\beta, \gamma} = 0$ if $l(\alpha) = l(\gamma)$, $\alpha \neq \gamma$, and there exists $\delta \in I(\Gamma^n)$, such that $\alpha \succ \delta$, $\gamma \succ \delta$;
- (iv) If $l(\alpha) = n$ and $\delta \succ \alpha \succ \beta$ in Γ^{n+1} , then

$$A_\alpha^\beta A_{\alpha, \delta} = A_{\alpha, \delta} \left(\sum_{\gamma} A_{\delta, \gamma} A_{\gamma, \delta} \right),$$

$$A_{\delta, \alpha} A_\alpha^\beta = \left(\sum_{\gamma} A_{\delta, \gamma} A_{\gamma, \delta} \right) A_{\delta, \alpha}$$

where the summation is over $\gamma \in I(\Gamma^n)$ such that $\gamma \neq \alpha$, $\delta \succ \gamma \succ \beta$.

- (v) If $l(\alpha) = n$ and $\alpha \succ \beta \succ \gamma$ in Γ^{n+2} , then

$$\left[A_\alpha^\beta, \sum_{\delta} A_\alpha^\delta \right] = 0$$

where the summation is over $\delta \in I(\Gamma^n)$, $\alpha \succ \delta \succ \gamma$.

A morphism of two level n quivers is defined in the standard way.

The level n quivers of Γ^n form an abelian category denoted by Qui_{Γ^n} . It is equivalent to the category of finite-dimensional modules over the suitable path algebra.

Note that the structure of the defining relations (i-v) for a level n quiver is completely determined by the adjacency matrix of the level $n+2$ truncated graph Γ^{n+2} .

For $\alpha, \beta \in I(\Gamma)$, let $\varepsilon(\beta, \alpha)$ be defined as in Section 2.1.

Let $\mathcal{V}_{\Gamma^n} = \{V_\alpha, A_{\alpha, \beta}, A_\alpha^\beta\}$ be a level n quiver. Then the collection $\mathcal{W}_{\Gamma^n} = \{W_\alpha, B_{\alpha, \beta}, B_\alpha^\beta\}$, where $W_\alpha = V_\alpha^*$, $B_{\alpha, \beta} = \varepsilon(\beta, \alpha) A_{\beta, \alpha}^*$: $W_\beta \rightarrow W_\alpha$, $B_\alpha^\beta = -(A_\alpha^\beta)^*$: $W_\alpha \rightarrow W_\alpha$, determines a level n quiver $\tau_n(\mathcal{V}_{\Gamma^n})$ called the dual quiver. The assignment $\mathcal{V}_{\Gamma^n} \mapsto \tau_n(\mathcal{V}_{\Gamma^n})$ determines an anti-automorphism of the category Qui_{Γ^n} of the fourth order, such that for any level n quiver $\mathcal{V}_{\Gamma^n}$ the quiver $\tau_n^2(\mathcal{V}_{\Gamma^n})$ is isomorphic to $\mathcal{V}_{\Gamma^n}$. The isomorphism $\phi : \mathcal{V}_{\Gamma^n} \rightarrow \tau^2(\mathcal{V}_{\Gamma^n})$ is given by the relation $\phi(v_\alpha) = (-1)^{l(\alpha)} v_\alpha$.

For a vertex β and an edge (α, β) of Γ^n , denote by A_β^α the operator

$$A_\beta^\alpha = A_{\beta, \alpha} A_{\alpha, \beta} : V_\beta \rightarrow V_\beta.$$

For a quiver $\mathcal{V}_{\Gamma^n}$ of level n and $\beta \in I(\Gamma^n)$, $\beta \neq \emptyset$, define the *monodromy operators*

$$T_\beta : \bigoplus_{\alpha, \alpha \succ \beta} V_\alpha \rightarrow \bigoplus_{\alpha, \alpha \succ \beta} V_\alpha, \quad \bar{T}_\beta : \bigoplus_{\alpha, \alpha \succ \beta} V_\alpha \rightarrow \bigoplus_{\alpha, \alpha \succ \beta} V_\alpha,$$

by formulas (3.5) and (3.6) and for any $\beta \in I(\Gamma^n)$ define the operator

$$\tilde{S}_\beta : V_\beta \rightarrow V_\beta,$$

by formula (3.7).

Suppose that for every vertex $\beta \in I(\Gamma^n)$, $\beta \neq \emptyset$, a non-resonant subset $G_\beta \subset \mathbb{C}$ is chosen. Denote by $Qui_{\Gamma^n, \{G_\beta\}}$ the category of all quivers $\{V_\alpha, A_{\alpha, \beta}, A_\alpha^\beta\} \in Qui_{\Gamma^n}$ such that

- (a) for any $\beta \in I(\Gamma^n)$, $\beta \neq \emptyset$, the eigenvalues of the operator T_β belong to G_β ,
- (b) for any $\beta \in I(\Gamma^n)$, $\beta \neq \emptyset$, none of the eigenvalues of the operator $\bar{T}_\beta$ is a positive integer.

Suppose we have

- a non-resonant subset $G_\beta \subset \mathbb{C}$ for every vertex $\beta \in I(\Gamma^n)$, $\beta \neq \emptyset$,
- a weakly non-resonant set $\tilde{G}_\beta$ for every vertex $\beta \in I(\Gamma^n)$,
- a weakly non-resonant subset $G_{\alpha, \beta}$ for every edge $(\alpha, \beta) \in E(\Gamma^n)$, $\alpha \succ \beta$, and loop $(\alpha, \alpha)^\beta$ of Γ^n .

Denote by $Qui_{\Gamma^n, \{G_\beta, \tilde{G}_\beta, G_{\alpha, \beta}\}}$ the subcategory of all quivers $\{V_\alpha, A_{\alpha, \beta}, A_\alpha^\beta\} \in Qui_{\Gamma^n}$, such that

- (a) for any $\beta \in I(\Gamma^n)$, $\beta \neq \emptyset$, the eigenvalues of the operator T_β belong to G_β ,
- (c) for any $\beta \in I(\Gamma^n)$, the eigenvalues of the operator $\tilde{S}_\beta$ belong to $\tilde{G}_\beta$ and
- (d) for any edge $(\alpha, \beta) \in E(\Gamma^n)$, $\alpha \succ \beta$, and loop $(\alpha, \alpha)^\beta$ the eigenvalues of the operator A_α^β belong to $G_{\alpha, \beta}$.

The categories $Qui_{\Gamma^n, \{G_\beta, \tilde{G}_\beta, G_{\alpha, \beta}\}}$ and $Qui_{\Gamma^n, \{G_\beta\}}$ are full abelian subcategories of the category Qui_{Γ^n} . The categories $Qui_{\Gamma^n, \{G_\beta, \tilde{G}_\beta, G_{\alpha, \beta}\}}$ and $Qui_{\Gamma^n, \{G_\beta\}}$ are closed under extensions.

We call a level n quiver *non-resonant* if it belongs to the subcategory $Qui_{\Gamma^n, \{G_\beta\}}$ for some collection $\{G_\beta\}$. We call a level n quiver *strongly non-resonant* if it belongs to the subcategory $Qui_{\Gamma^n, \{G_\beta, \tilde{G}_\beta, G_{\alpha, \beta}\}}$ for some collections $\{G_\beta\}$, $\{\tilde{G}_\beta\}$, and $\{G_{\alpha, \beta}\}$.

3.3. Restrictions of quivers. For integers k, l , $0 \leq k < l \leq N$, we define a functor $\mathbb{J}_{l, k}^* : Qui_{\Gamma^l} \rightarrow Qui_{\Gamma^k}$. If $\mathcal{V}_{\Gamma^l} = \{V_\alpha, A_{\alpha, \beta}\}$ is a quiver from Qui_{Γ^l} , then the quiver $\mathbb{J}_{l, k}^*(\mathcal{V}_{\Gamma^l}) = \{W_\alpha, B_{\alpha, \beta}, B_\alpha^\beta\} \in Qui_{\Gamma^k}$ is defined as follows.

For every $\alpha \in I(\Gamma^k)$, the space W_α coincides with the space V_α . For every $\alpha, \beta \in I(\Gamma^k)$, $\alpha \neq \beta$, the map $B_{\alpha, \beta} : W_\beta \rightarrow W_\alpha$ coincides with the map $A_{\alpha, \beta} : V_\beta \rightarrow V_\alpha$. For every loop $(\alpha, \alpha)^\beta \in E(\Gamma^k)$, the map B_α^β is equal to the product $A_{\alpha, \beta} A_{\beta, \alpha}$.

One can check that relations (i-v) in Section 2.2 for the level l quiver $\mathcal{V}_{\Gamma^l}$ imply relations (i-v) in Section 2.2 for the level k quiver $\mathbb{J}_{l, k}^*(\mathcal{V}_{\Gamma^l})$. If $\mathcal{V}_{\Gamma^l} = \{V_\alpha, A_{\alpha, \beta}\}$ and $\mathcal{V}'_{\Gamma^l} = \{V'_\alpha, A'_{\alpha, \beta}\}$

are two level l quivers, and $\varphi = \{\varphi_\alpha : V_\alpha \rightarrow V'_\alpha \mid \alpha \in I(\Gamma^l)\}$ a morphism of these quivers, then the collection $\{\varphi_\alpha : V_\alpha \rightarrow V'_\alpha \mid \alpha \in I(\Gamma^k)\}$ defines a morphism $\mathbb{J}_{l,k}^*(\varphi)$ of level k quivers $\mathbb{J}_{l,k}^*(\mathcal{V}_{\Gamma^l})$ and $\mathbb{J}_{l,k}^*(\mathcal{V}'_{\Gamma^l})$.

One can check that the functor $\mathbb{J}_{l,k}^*$ is exact and it maps the subcategory of non-resonant Γ^l quivers to the subcategory of non-resonant Γ^k quivers. For $0 \leq k_1 < k_2 < k_3 \leq N$, we have

$$(3.10) \quad \mathbb{J}_{k_3,k_1}^* = \mathbb{J}_{k_2,k_1}^* \cdot \mathbb{J}_{k_3,k_2}^*,$$

and for any $0 \leq k < l \leq N$

$$(3.11) \quad \mathbb{J}_{l,k}^* \cdot \tau_l = \tau_k \cdot \mathbb{J}_{l,k}^*.$$

We will use the notation $\mathbb{J}_0^*$ for the functor $\mathbb{J}_{N,0}^* : Qui_\Gamma = Qui_{\Gamma^N} \rightarrow Qui_{\Gamma^0}$.

The quiver $\mathbb{J}_{l,k}^*(\mathcal{V}_{\Gamma^l})$ is called *the restriction at level k of the quiver $\mathcal{V}_{\Gamma^l}$* . The functors $\mathbb{J}_{l,k}^*$ are called *the restriction functors*.

3.4. Direct images of quivers. Now we recall the definition of the left and right adjoint functors.

Let $\mathcal{A}$ and $\mathcal{B}$ be additive categories; let $F : \mathcal{A} \rightarrow \mathcal{B}$ and $G : \mathcal{B} \rightarrow \mathcal{A}$ be additive functors. The functor F is called *a left adjoint* to G and the functor G is called *a right adjoint* to F , if there exists a collection of isomorphisms

$$(3.12) \quad \alpha_{X,Y} : \text{Hom}_{\mathcal{A}}(X, G(Y)) \xrightarrow{\sim} \text{Hom}_{\mathcal{B}}(F(X), Y), \quad X \in \mathcal{A}, \quad Y \in \mathcal{B},$$

such that

- for any $X, X' \in \mathcal{A}$, $Y \in \mathcal{B}$, $\phi \in \text{Hom}_{\mathcal{A}}(X, G(Y))$, $\gamma \in \text{Hom}_{\mathcal{A}}(X', X)$, we have $\alpha_{X',Y}(\phi\gamma) = \alpha_{X,Y}(\phi)F(\gamma)$,
- for any $X \in \mathcal{A}$, $Y, Y' \in \mathcal{B}$, $\phi \in \text{Hom}_{\mathcal{B}}(F(X), Y)$, $\delta \in \text{Hom}_{\mathcal{B}}(Y, Y')$, we have $\alpha_{X,Y'}(G(\delta)\phi) = \delta \cdot \alpha_{X,Y}(\phi)$.

For a given additive functor $F : \mathcal{A} \rightarrow \mathcal{B}$ its left adjoint functor $G : \mathcal{B} \rightarrow \mathcal{A}$ is unique up to an isomorphism, if exists. Analogously, for a given additive functor $G : \mathcal{B} \rightarrow \mathcal{A}$ its right adjoint functor $F : \mathcal{A} \rightarrow \mathcal{B}$ is unique up to an isomorphism, if exists.

The isomorphisms (3.12) determine collections of canonical morphisms

$$(3.13) \quad \sigma_{F,G} : X \rightarrow GF(X), \quad \tau_{F,G} : FG(Y) \rightarrow Y, \quad X \in \mathcal{A}, \quad Y \in \mathcal{B},$$

where $\sigma_{F,G} = \alpha_{X,GF(X)}^{-1}(\text{id}_{GF(X)})$ and $\tau_{F,G} = \alpha_{FG(Y),Y}(\text{id}_{FG(Y)})$. The collections determine natural transformations of the functors $\sigma_{F,G} : Id_{\mathcal{A}} \rightarrow GF$ and $\tau_{F,G} : FG \rightarrow Id_{\mathcal{B}}$, that is,

- for any $X, X' \in \mathcal{A}$, $\gamma \in \text{Hom}_{\mathcal{A}}(X', X)$ we have the equality $\sigma_{F,G}\gamma = GF(\gamma)\sigma_{F,G}$ in $\text{Hom}_{\mathcal{A}}(X', GF(X))$,
- for any $Y, Y' \in \mathcal{B}$, $\delta \in \text{Hom}_{\mathcal{B}}(Y, Y')$ we have the equality $\delta\tau_{F,G} = \tau_{F,G}FG(\delta)$ in $\text{Hom}_{\mathcal{B}}(FG(Y), Y')$.

Our next goal is to describe the functor $\mathbb{J}_{n,n-1,*} : Qui_{\Gamma^{n-1}} \rightarrow Qui_{\Gamma^n}$, which is the right adjoint functor to the functor $\mathbb{J}_{n,n-1}^*$.

Let $\mathcal{V}_{\Gamma^{n-1}} = \{V_\alpha, A_{\alpha,\beta}, A_\alpha^\beta\}$ be a level $n-1$ quiver. Define a new level n quiver $\mathcal{W}_{\Gamma^n} = \{W_\alpha, B_{\alpha,\beta}, B_\alpha^\beta\}$ as follows.

For $\alpha \in I(\Gamma^{n-1})$, set $W_\alpha = V_\alpha$.

For $\alpha \in I_n(\Gamma)$, denote by $\widetilde{W}_\alpha$ the direct sum $\widetilde{W}_\alpha = \bigoplus_{\gamma \in I_{n-1}(\Gamma), \gamma \succ \alpha} V_\gamma$. Set W_α to be the subspace of $\widetilde{W}_\alpha$, generated by the vectors $w = \sum_{\gamma \in I_{n-1}(\Gamma), \gamma \succ \alpha} v_\gamma$, $v_\gamma \in V_\gamma$, such that for any $\delta \in I_{n-2}(\Gamma)$ with $\delta > \alpha$, the sum $\sum_{\gamma \in I_{n-1}(\Gamma), \gamma \succ \alpha} A_{\delta, \gamma}(v_\gamma)$ equals zero.

Set $B_{\alpha, \beta} = A_{\alpha, \beta}$, if α and β both belong to $I(\Gamma^{n-1})$.

Let $\alpha \in I_{n-1}(\Gamma)$, $\beta \in I_n(\Gamma)$, and $\alpha \succ \beta$. Set $B_{\alpha, \beta} : W_\beta \rightarrow W_\alpha$ to be the restriction to the subspace W_β of the projection of the space $\widetilde{W}_\beta$ to the space $W_\alpha = V_\alpha$ along the other direct summands of $\widetilde{W}_\beta$. Define the map $B_{\beta, \alpha} : W_\alpha \rightarrow W_\beta$ by the relation

$$B_{\beta, \alpha}(v_\alpha) = A_\alpha^\beta(v_\alpha) - \sum A_{\alpha', \delta} A_{\delta, \alpha}(v_\alpha),$$

where $v_\alpha \in V_\alpha = W_\alpha$ and the sum is over all $\alpha' \in I_{n-1}(\Gamma)$ and $\delta \in I_{n-2}(\Gamma)$, such that $\delta \succ \alpha$ and $\delta \succ \alpha'$.

Let $\alpha \in I_n(\Gamma)$, $\beta \in I_{n+1}(\Gamma)$, and $\alpha \succ \beta$. Define the map $B_\alpha^\beta : W_\alpha \rightarrow W_\alpha$ by the relation

$$B_\alpha^\beta\left(\sum_{\gamma \in I_{n-1}(\Gamma), \gamma \succ \alpha} v_\gamma\right) = \sum_{\gamma, \gamma \succ \alpha} \sum_{\delta, \gamma \succ \delta \succ \beta, \delta \neq \alpha} A_\gamma^\delta(v_\gamma),$$

where $v_\gamma \in V_\gamma$.

Proposition 3.3. *The collection $\mathcal{W}_{\Gamma^n} = \{W_\alpha, B_{\alpha, \beta}, B_\alpha^\beta\}$ is a level n quiver.* $\square$

The assignment $\mathcal{V}_{\Gamma^{n-1}} \mapsto \mathcal{W}_{\Gamma^n}$ determines a functor $\mathbb{J}_{n, n-1, *}: \text{Qui}_{\Gamma^{n-1}} \rightarrow \text{Qui}_{\Gamma^n}$.

Let $\mathcal{V}'_{\Gamma^n} = \{V'_\alpha, A'_{\alpha, \beta}, A'^\beta_\alpha\}$ be a level n quiver and ϕ a morphism of Γ_{n-1} quivers $\phi : \mathbb{J}_{n, n-1}^*(\mathcal{V}'_{\Gamma^n}) \rightarrow \mathcal{V}_{\Gamma^{n-1}}$, given by a collection of linear maps $\phi_\alpha : V'_\alpha \rightarrow V_\alpha$, $\alpha \in I(\Gamma^{n-1})$.

Define a new morphism of level n quivers $\tilde{\phi} : \mathcal{V}'_{\Gamma^n} \rightarrow \mathcal{W}_{\Gamma^n}$ by the following rules.

Set $\tilde{\phi}_\alpha = \phi_\alpha : V'_\alpha \rightarrow W_\alpha = V_\alpha$ if $l(\alpha) < n$.

Let $l(\beta) = n$ and $v'_\beta \in V'_\beta$. Set $\tilde{\phi}_\beta(v'_\beta) = \sum_{\alpha, \alpha \succ \beta} \phi_\alpha A'_{\alpha, \beta}(v'_\beta)$. We have $\tilde{\phi}_\beta(v'_\beta) \subset W_\beta$.

Proposition 3.4.

(i) *The assignment $\beta_{\mathcal{V}'_{\Gamma^n}, \mathcal{V}_{\Gamma^{n-1}}} : \phi \mapsto \tilde{\phi}$ determines an isomorphism*

$$\beta_{\mathcal{V}'_{\Gamma^n}, \mathcal{V}_{\Gamma^{n-1}}} : \text{Hom}_{\text{Qui}_{\Gamma^{n-1}}}(\mathbb{J}_{n, n-1}^*(\mathcal{V}'_{\Gamma^n}), \mathcal{V}_{\Gamma^{n-1}}) \xrightarrow{\sim} \text{Hom}_{\text{Qui}_{\Gamma^n}}(\mathcal{V}'_{\Gamma^n}, \mathbb{J}_{n, n-1, *}(\mathcal{V}_{\Gamma^{n-1}})).$$

(ii) *The functors $\mathbb{J}_{n, n-1}^*$ and $\mathbb{J}_{n, n-1, *}$ form a pair of left and right adjoint functors with respect to the collection of isomorphisms*

$$\alpha_{\mathcal{V}'_{\Gamma^n}, \mathcal{V}_{\Gamma^{n-1}}} : \text{Hom}_{\text{Qui}_{\Gamma^n}}(\mathcal{V}'_{\Gamma^n}, \mathbb{J}_{n, n-1, *}(\mathcal{V}_{\Gamma^{n-1}})) \xrightarrow{\sim} \text{Hom}_{\text{Qui}_{\Gamma^{n-1}}}(\mathbb{J}_{n, n-1}^*(\mathcal{V}'_{\Gamma^n}), \mathcal{V}_{\Gamma^{n-1}}),$$

where $\alpha_{\mathcal{V}'_{\Gamma^n}, \mathcal{V}_{\Gamma^{n-1}}} = \beta_{\mathcal{V}'_{\Gamma^n}, \mathcal{V}_{\Gamma^{n-1}}}^{-1}$. $\square$

For $0 \leq k < l \leq N$, define the functors

$$\mathbb{J}_{l, k, *}: \text{Qui}_{\Gamma^k} \rightarrow \text{Qui}_{\Gamma^l}, \quad \mathbb{J}_{l, k, !}: \text{Qui}_{\Gamma^k} \rightarrow \text{Qui}_{\Gamma^l}$$

by the formulas

$$(3.14) \quad \mathbb{J}_{l, k, *} = \mathbb{J}_{l, l-1, *} \cdots \mathbb{J}_{k+1, k, *}(\mathcal{V}_{\Gamma^k}), \quad \mathbb{J}_{l, k, !} = \tau_l^{-1} \cdot \mathbb{J}_{l, k, *} \cdot \tau_k.$$

The functors $\mathbb{J}_{l, k, *}$ and $\mathbb{J}_{l, k, !}$ are called *the quiver direct image functors*.

For $0 \leq k_1 < k_2 < k_3 \leq N$, we have

$$\mathbb{J}_{k_3, k_1, *} = \mathbb{J}_{k_3, k_2, *} \cdot \mathbb{J}_{k_2, k_1, *}, \quad \mathbb{J}_{k_3, k_1, !} = \mathbb{J}_{k_3, k_2, !} \cdot \mathbb{J}_{k_2, k_1, !}.$$

We will use the notation $\mathbb{J}_{0,*}$ and $\mathbb{J}_{0,!}$, respectively, for the functors $\mathbb{J}_{N,0,*} : Qui_{\Gamma^0} \rightarrow Qui_{\Gamma^N} = Qui_{\Gamma}$ and $\mathbb{J}_{N,0,!} : Qui_{\Gamma^0} \rightarrow Qui_{\Gamma^N} = Qui_{\Gamma}$.

The functor $\mathbb{J}_{l,k,*}$ is the right adjoint to the functor $\mathbb{J}_{l,k}^*$. For any $\mathcal{V}'_{\Gamma^l}$ and $\mathcal{V}_{\Gamma^k}$ the isomorphism $\alpha_{\mathcal{V}'_{\Gamma^l}, \mathcal{V}_{\Gamma^k}} : \text{Hom}_{Qui_{\Gamma^l}}(\mathcal{V}'_{\Gamma^l}, \mathbb{J}_{l,k,*}(\mathcal{V}_{\Gamma^k})) \xrightarrow{\sim} \text{Hom}_{Qui_{\Gamma^k}}(\mathbb{J}_{l,k}^*(\mathcal{V}'_{\Gamma^l}), \mathcal{V}_{\Gamma^k})$ is the composition

$$\alpha_{\mathcal{V}'_{\Gamma^l}, \mathcal{V}_{\Gamma^k}} = \alpha_{\mathbb{J}_{l,k+1}^*(\mathcal{V}'_{\Gamma^l}), \mathcal{V}_{\Gamma^k}} \cdots \alpha_{\mathbb{J}_{l,l-1}^*(\mathcal{V}'_{\Gamma^l}), \mathbb{J}_{l-2,k,*}(\mathcal{V}_{\Gamma^k})} \alpha_{\mathcal{V}'_{\Gamma^l}, \mathbb{J}_{l-1,k,*}(\mathcal{V}_{\Gamma^k})}.$$

The functor $\mathbb{J}_{l,k,!}$ is left adjoint to the functor $\mathbb{J}_{l,k}^*$. For any $\mathcal{V}'_{\Gamma^l}$ and $\mathcal{V}_{\Gamma^k}$ the corresponding isomorphism $\alpha'_{\mathcal{V}_{\Gamma^k}, \mathcal{V}'_{\Gamma^l}} : \text{Hom}_{Qui_{\Gamma^k}}(\mathcal{V}_{\Gamma^k}, \mathbb{J}_{l,k}^*(\mathcal{V}'_{\Gamma^l})) \xrightarrow{\sim} \text{Hom}_{Qui_{\Gamma^l}}(\mathbb{J}_{l,k,!}(\mathcal{V}_{\Gamma^k}), \mathcal{V}'_{\Gamma^l})$ is defined by the relation

$$\alpha'_{\mathcal{V}_{\Gamma^k}, \mathcal{V}'_{\Gamma^l}}(\phi) = \tau_l^{-1} \cdot \alpha_{\tau_l(\mathcal{V}'_{\Gamma^l}), \tau_k(\mathcal{V}_{\Gamma^k})} \cdot \tau_k(\phi).$$

Thus $\mathbb{J}_{l,k}^* : Qui_{\Gamma^l} \rightarrow Qui_{\Gamma^k}$ and $\mathbb{J}_{l,k,*} : Qui_{\Gamma^k} \rightarrow Qui_{\Gamma^l}$ form a left and right adjoint pair, as well as $\mathbb{J}_{l,k,!} : Qui_{\Gamma^k} \rightarrow Qui_{\Gamma^l}$ and $\mathbb{J}_{l,k}^* : Qui_{\Gamma^l} \rightarrow Qui_{\Gamma^k}$ form a left and right adjoint pair.

Remark. Below we give a direct description of the functor $\mathbb{J}_{n,n-1,!}$.

Let $\mathcal{V}_{\Gamma^{n-1}} = \{V_{\alpha}, A_{\alpha,\beta}, A_{\alpha}^{\beta}\}$ be a level $n-1$ quiver. Define a new level n quiver $\mathcal{U}_{\Gamma^n} = \{U_{\alpha}, C_{\alpha,\beta}, C_{\alpha}^{\beta}\}$ as follows.

For $\alpha \in I(\Gamma^{n-1})$, we set $U_{\alpha} = V_{\alpha}$.

For $\alpha \in I_n(\Gamma)$, we set $\widetilde{U}_{\alpha} = \bigoplus_{\gamma \in I_{n-1}(\Gamma), \gamma \succ \alpha} V_{\gamma}$. We set U_{α} to be the quotient of $\widetilde{U}_{\alpha}$ over the subspace, generated by the vectors $\sum_{\alpha, \delta \succ \alpha \succ \beta} A_{\alpha,\delta}(v_{\delta})$, where $\delta \in I_{n-2}(\Gamma)$, $\delta \succ \beta$, and $v_{\delta} \in V_{\delta}$. Let $\pi : \widetilde{U}_{\alpha} \rightarrow U_{\alpha}$ be the natural projection.

We set $C_{\alpha,\beta} = A_{\alpha,\beta}$, if α and β both belong to $I(\Gamma^{n-1})$.

Let $\alpha \in I_{n-1}(\Gamma)$, $\beta \in I_n(\Gamma)$, and $\alpha \succ \beta$. We set $C_{\beta,\alpha} : U_{\alpha} \rightarrow U_{\beta}$ to be the composition of the inclusion $U_{\alpha} = V_{\alpha} \hookrightarrow \widetilde{U}_{\beta} = \bigoplus_{\gamma \in I_{n-1}(\Gamma), \gamma \succ \alpha} V_{\gamma}$ and the projection $\pi : \widetilde{U}_{\beta} \rightarrow U_{\beta}$. Define the map $C_{\alpha,\beta} : U_{\beta} \rightarrow U_{\alpha}$, $u \mapsto C_{\alpha,\beta}(u)$ by the following rules:

- Assume that there exists $v_{\alpha} \in V_{\alpha}$ such that $\pi(v_{\alpha}) = u$, then we set $C_{\alpha,\beta}(u) = \pi(A_{\alpha}^{\beta}(v_{\alpha}))$.
- Assume that there exists $\gamma \in I_{n-1}(\Gamma)$, $\gamma \neq \alpha$, and $v_{\gamma} \in V_{\gamma}$ such that $\pi(v_{\gamma}) = u$. Assume that there exists $\delta \in I_{n-2}(\Gamma)$ such that $\delta \succ \alpha, \delta \succ \gamma$. Then we set $C_{\alpha,\beta}(u) = -\pi(A_{\alpha,\delta}A_{\delta,\gamma}(v_{\gamma}))$.
- Assume that there exists $\gamma \in I_{n-1}(\Gamma)$, $\gamma \neq \alpha$, and $v_{\gamma} \in V_{\gamma}$ such that $\pi(v_{\gamma}) = u$. Assume that there is no $\delta \in I_{n-2}(\Gamma)$ such that $\delta \succ \alpha, \delta \succ \gamma$. Then we set $C_{\alpha,\beta}(u) = 0$.

Let $\alpha \in I_n(\Gamma)$, $\beta \in I_{n+1}(\Gamma)$, and $\alpha \succ \beta$. Define the map $C_{\alpha}^{\beta} : U_{\alpha} \rightarrow U_{\alpha}$ by the relation

$$C_{\alpha}^{\beta} \pi \left(\sum_{\gamma \in I_{n-1}(\Gamma), \gamma \succ \alpha} v_{\gamma} \right) = \sum_{\gamma, \gamma \succ \alpha} \sum_{\delta, \gamma \succ \delta \succ \beta, \delta \neq \alpha} \pi(A_{\gamma}^{\delta}(v_{\gamma})),$$

where $v_{\gamma} \in V_{\gamma}$.

Proposition 3.5. *The collection $\mathcal{U}_{\Gamma^n} = \{U_{\alpha}, C_{\alpha,\beta}, C_{\alpha}^{\beta}\}$ is a level n quiver. This level n quiver is isomorphic to $\mathbb{J}_{n,n-1,!}(\mathcal{V}_{\Gamma^{n-1}})$.* $\square$

Proposition 3.6. *For any $k < l$ and any level k quiver $\mathcal{V}_{\Gamma^k} = \{V_\alpha, A_{\alpha,\beta}, A_\alpha^\beta\}$, the canonical morphisms*

$$(3.15) \quad \begin{aligned} \sigma_{\mathbb{J}_{l,k,!}, \mathbb{J}_{l,k}^*} &: \mathcal{V}_{\Gamma^k} \rightarrow \mathbb{J}_{l,k}^* \mathbb{J}_{l,k,!}(\mathcal{V}_{\Gamma^k}), \\ \tau_{\mathbb{J}_{l,k}^*, \mathbb{J}_{l,k,*}} &: \mathbb{J}_{l,k}^* \mathbb{J}_{l,k,*}(\mathcal{V}_{\Gamma^k}) \rightarrow \mathcal{V}_{\Gamma^k} \end{aligned}$$

are isomorphisms. □

The proofs of Propositions 3.3–3.6 are direct linear algebra calculations.

Define the morphism

$$s_{l,k} : \mathbb{J}_{l,k,!}(\mathcal{V}_{\Gamma^k}) \rightarrow \mathbb{J}_{l,k,*}(\mathcal{V}_{\Gamma^k})$$

by the formula

$$(3.16) \quad s_{l,k} = \alpha_{\mathcal{V}_{\Gamma^k}, \mathbb{J}_{l,k,*}(\mathcal{V}_{\Gamma^k})} \left(\tau_{\mathbb{J}_{l,k}^*, \mathbb{J}_{l,k,*}}^{-1} \right),$$

where $\alpha_{X,Y}$ are isomorphisms (3.12), establishing the adjunction property of the functors $\mathbb{J}_{l,k,!}$ and $\mathbb{J}_{l,k}^*$.

For $0 \leq k < l \leq N$ and a level k quiver $\mathcal{V}_{\Gamma^k} = \{V_\alpha, A_{\alpha,\beta}, A_\alpha^\beta\}$, denote by $\mathbb{J}_{l,k,!}(\mathcal{V}_{\Gamma^k}) \in Qui_{\Gamma^l}$ the image of $\mathbb{J}_{l,k,!}(\mathcal{V}_{\Gamma^k})$ in $\mathbb{J}_{l,k,*}(\mathcal{V}_{\Gamma^k})$ with respect to $s_{l,k}$.

We use the notation s_0 for the morphism $s_{N,0}$ and $\mathbb{J}_{0,!}(\mathcal{V}_{\Gamma^0})$ for the quiver $\mathbb{J}_{0,N,!}(\mathcal{V}_{\Gamma^0}) \in Qui_{\Gamma}$.

3.5. Example. Let X be $\mathbb{C}^2$ with coordinates z_1, z_2 . Let $\mathcal{C} = \{H_1, H_2, H_3\}$ be the arrangement of three lines, $H_1 = \{z_1 = 0\}$, $H_2 = \{z_2 = 0\}$, $H_3 = \{z_1 = z_2\}$, intersecting at one point. The stratification of X consists of 5 strata:

- The complement to the union of lines, $X_\emptyset = \{(z_1, z_2) \mid z_1 \neq 0, z_2 \neq 0, z_1 \neq z_2\}$;
- The lines without the point, $X_{\alpha_i} = H_i \setminus \{(0, 0)\}$, $i = 1, 2, 3$;
- The point, $X_\beta = \{(0, 0)\}$.

The principal open subset X^0 is $X_\emptyset$. The principal open subset X^1 is the complement to the point, $X^1 = X \setminus \{(0, 0)\}$. The principal open subset X^2 is X .

The graph Γ of the arrangement has five vertices, $\emptyset, \alpha_1, \alpha_2, \alpha_3, \beta$, such that $l(\emptyset) = 0$, $l(\alpha_i) = 1$, $l(\beta) = 2$. The graph has six edges $(\emptyset, \alpha_i)$, (α_i, β) . We have $\emptyset \succ \alpha_i \succ \beta$.

The level two graph Γ^2 coincides with Γ . The level one graph Γ^1 has four vertices $\emptyset, \alpha_1, \alpha_2, \alpha_3$, three edges $(\emptyset, \alpha_i)$ and three loops $(\alpha_i, \alpha_i)^\beta$. The level zero graph Γ^0 consists of one vertex $\emptyset$ and three loops $(\emptyset, \emptyset)^{\alpha_i}$.

- (i) A quiver $\mathcal{V}_\Gamma$ (the same as a level two quiver) is a collection of five vector spaces, $V_\emptyset, V_\beta, V_{\alpha_i}$, $i = 1, 2, 3$, and twelve linear maps, $A_{\alpha_i, \emptyset} : V_\emptyset \rightarrow V_{\alpha_i}$, $A_{\emptyset, \alpha_i} : V_{\alpha_i} \rightarrow V_\emptyset$, $A_{\alpha_i, \beta} : V_\beta \rightarrow V_{\alpha_i}$, $A_{\beta, \alpha_i} : V_{\alpha_i} \rightarrow V_\beta$, such that

$$\sum_{i=1}^3 A_{\beta, \alpha_i} A_{\alpha_i, \emptyset} = 0, \quad \sum_{i=1}^3 A_{\emptyset, \alpha_i} A_{\alpha_i, \beta} = 0,$$

and

$$A_{\alpha_j, \emptyset} A_{\emptyset, \alpha_i} + A_{\alpha_j, \beta} A_{\beta, \alpha_i} = 0$$

for all pairs (i, j) , $i \neq j$.

- (ii) A level one quiver $\mathcal{V}_{\Gamma^1}$ is a collection of four vector spaces, $V_\emptyset, V_{\alpha_i}, i = 1, 2, 3$, and nine linear maps $A_{\alpha_i, \emptyset} : V_\emptyset \rightarrow V_{\alpha_i}, A_{\emptyset, \alpha_i} : V_{\alpha_i} \rightarrow V_\emptyset, A_{\alpha_i}^\beta : V_{\alpha_i} \rightarrow V_{\alpha_i}$, such that

$$A_{\alpha_i}^\beta A_{\alpha_i, \emptyset} = A_{\alpha_i, \emptyset} \sum_{j \neq i} A_{\emptyset, \alpha_j} A_{\alpha_j, \emptyset},$$

$$A_{\emptyset, \alpha_i} A_{\alpha_i}^\beta = \sum_{j \neq i} A_{\emptyset, \alpha_j} A_{\alpha_j, \emptyset} A_{\emptyset, \alpha_i}.$$

- (iii) A level zero quiver $\mathcal{V}_{\Gamma^0}$ consists of the vector space $V = V_\emptyset$ and three linear maps $A^i : V \rightarrow V, i = 1, 2, 3$. subject to the relations

$$[A^i, A^1 + A^2 + A^3] = 0, \quad i = 1, 2, 3.$$

Let $\mathcal{V}_\Gamma$ be a quiver as in (i). Then its restriction $\mathbb{J}_{2,1}^*(\mathcal{V}_\Gamma)$ to level one is the level one quiver $\{W_\emptyset, W_{\alpha_i}, B_{\emptyset, \alpha_i}, B_{\alpha_i, \emptyset}, B_{\alpha_i}^\beta\}$, where $W_\emptyset = V_\emptyset, W_{\alpha_i} = V_{\alpha_i}, B_{\emptyset, \alpha_i} = A_{\emptyset, \alpha_i}, B_{\alpha_i, \emptyset} = A_{\alpha_i, \emptyset}$, and $B_{\alpha_i}^\beta = A_{\alpha_i, \beta} A_{\beta, \alpha_i}$.

Let $\mathcal{V}_{\Gamma^1}$ be a level one quiver as in (ii). Its restriction $\mathbb{J}_{1,0}^*(\mathcal{V}_{\Gamma^1})$ to level zero is the level zero quiver $\{W_\emptyset, B^i, \}$, where $W_\emptyset = V_\emptyset, B^i = A_{\emptyset, \alpha_i} A_{\alpha_i, \emptyset}$.

Let $\mathcal{V}_{\Gamma^0}$ be a level zero quiver as in (iii). Its direct image $\mathbb{J}_{1,0,*}(\mathcal{V}_{\Gamma^0})$ is the level one quiver $\{W_\emptyset, W_{\alpha_i}, B_{\emptyset, \alpha_i}, B_{\alpha_i, \emptyset}, B_{\alpha_i}^\beta\}$, where $W_\emptyset = V, W_{\alpha_i} = V, B_{\emptyset, \alpha_i} = \text{Id}, B_{\alpha_i, \emptyset} = A^i$, and $B_{\alpha_i}^\beta = \sum_{j \in \{1,2,3\}, j \neq i} A^j$.

The direct image $\mathbb{J}_{1,0,!}(\mathcal{V}_{\Gamma^0})$ is the level one quiver $\{U_\emptyset, U_{\alpha_i}, C_{\emptyset, \alpha_i}, C_{\alpha_i, \emptyset}, C_{\alpha_i}^\beta\}$, where $U_\emptyset = V, U_{\alpha_i} = V, C_{\alpha_i, \emptyset} = \text{Id}, C_{\emptyset, \alpha_i} = A^i$, and $C_{\alpha_i}^\beta = \sum_{j \in \{1,2,3\}, j \neq i} A^j$.

Let $\mathcal{V}_{\Gamma^1}$ be a level one quiver as in (ii). Its direct image $\mathbb{J}_{2,1,*}(\mathcal{V}_{\Gamma^1})$ is the level two quiver $\{W_\emptyset, W_{\alpha_i}, W_\beta, B_{\emptyset, \alpha_i}, B_{\alpha_i, \emptyset}, B_{\beta, \alpha_i}, B_{\alpha_i, \beta}\}$, where $W_\emptyset = V_\emptyset, W_{\alpha_i} = V_{\alpha_i}, W_\beta$ is the subspace of the direct sum $\widetilde{W}_\beta = \oplus_i V_{\alpha_i}$, generated by the vectors $\sum_{i=1}^3 v_{\alpha_i}$, such that $\sum_i A_{\emptyset, \alpha_i}(v_{\alpha_i}) = 0$. The maps $B_{\emptyset, \alpha_i}$ are equal to $A_{\emptyset, \alpha_i}$. The maps $B_{\alpha_i, \emptyset}$ are equal to $A_{\alpha_i, \emptyset}$. The maps $B_{\alpha_i, \beta}$ and B_{β, α_i} are defined by the formulas $B_{\alpha_i, \beta}(\sum_j v_{\alpha_j}) = v_{\alpha_i}$ and $B_{\beta, \alpha_i}(v_{\alpha_i}) = A_{\alpha_i}^\beta(v_{\alpha_i}) - \sum_{j \in \{1,2,3\}, j \neq i} A_{\alpha_j, \emptyset} A_{\emptyset, \alpha_i}(v_{\alpha_j})$.

The direct image $\mathbb{J}_{2,1,!}(\mathcal{V}_{\Gamma^1})$ is the level two quiver $\{U_\emptyset, U_{\alpha_i}, U_\beta, C_{\emptyset, \alpha_i}, C_{\alpha_i, \emptyset}, C_{\beta, \alpha_i}, C_{\alpha_i, \beta}, \}$, where $U_\emptyset = V_\emptyset$ and $U_{\alpha_i} = V_{\alpha_i}$. The space U_β is the quotient space of $\widetilde{U}_\beta = \oplus_i V_{\alpha_i}$ over the subspace, generated by the vectors $\sum_{i=1}^3 A_{\alpha_i, \emptyset}(v_\emptyset), v_\emptyset \in V_\emptyset$. Let $\pi : \widetilde{U}_\beta \rightarrow U_\beta$ be the canonical projection. The maps $C_{\emptyset, \alpha_i}$ are equal to $A_{\emptyset, \alpha_i}$. The maps $C_{\alpha_i, \emptyset}$ are equal to $A_{\alpha_i, \emptyset}$. The maps C_{β, α_i} are defined by the formulas $C_{\beta, \alpha_i}(v_{\alpha_i}) = \pi(v_{\alpha_i})$. The maps $C_{\alpha_i, \beta}$ are defined by the formulas $C_{\alpha_i, \beta}(\pi(v_{\alpha_i})) = \pi(A_{\alpha_i}^\beta(v_{\alpha_i}))$ and $C_{\alpha_i, \beta}(\pi(v_{\alpha_j})) = -\pi(A_{\alpha_i, \emptyset} A_{\emptyset, \alpha_j}(v_{\alpha_j}))$ if $j \neq i$.

3.6. Direct images of level zero quivers. The direct images of level zero quivers can be obtained by inductive application of the construction of the previous section. Here we give a direct description of the functors $\mathbb{J}_{0,*}$ and $\mathbb{J}_{0,!}$.

A level zero quiver $\mathcal{W}_{\Gamma^0} \in \text{Qui}_{\Gamma^0}$ consists of a vector space $W = W_{\emptyset}$ and linear operators $B^i = B_{\emptyset}^i : W \rightarrow W$, $i \in I_1(\Gamma)$, such that for each $\alpha \in I_2(\Gamma)$, $i \succ \alpha$, we have

$$(3.16) \quad [B^i, \sum_{j, j \succ \alpha} B^j] = 0.$$

Let $\mathcal{W}_{\Gamma^0} = \{W, B^i\}$ be a level zero quiver. Define a quiver $\mathcal{V}_{\Gamma} = \{V_{\alpha}, A_{\alpha, \beta}\} \in \text{Qui}_{\Gamma}$ as follows.

For $\alpha \in I(\Gamma)$, set $V_{\alpha} = \mathcal{F}_{\alpha} \otimes W$, where $\mathcal{F}_{\alpha}$ is the subspace of the flag space $\mathcal{F}(\mathcal{C})$, corresponding to $\alpha \in I(\Gamma)$.

For $\beta \succ \alpha$, $l(\beta) = m$ define the map $A_{\alpha, \beta} : V_{\beta} \rightarrow V_{\alpha}$ by the rule

$$(3.17) \quad A_{\alpha, \beta} : F_{\beta_0, \dots, \beta_{m-1}, \beta} \otimes w \mapsto (-1)^m F_{\beta_0, \dots, \beta_{m-1}, \beta, \alpha} \otimes w.$$

To define the maps $A_{\alpha, \beta} : V_{\beta} \rightarrow V_{\alpha}$ for $\alpha \succ \beta$ we need the following construction.

Let $F_{\beta_0, \dots, \beta_{m-1}, \beta_m} \in \mathcal{F}(\mathcal{C})$ be a flag and $\alpha \in I_{m-1}(\Gamma)$, $\alpha \succ \beta_m$, a vertex. Assume that there exists a flag $F_{\alpha_0, \dots, \alpha_{m-1}} \in \mathcal{F}(\mathcal{C})$ such that $\alpha_{m-1} = \alpha$ and $\alpha_k \succ \beta_{k+1}$ for $k = 0, \dots, m-1$. It is easy to see such a flag is unique. We call $F_{\alpha_0, \dots, \alpha_{m-1}}$ the *cutoff* of $F_{\beta_0, \dots, \beta_m}$ in the direction of α and denote by $F_{\beta_0, \dots, \beta_m}^{\alpha}$. If such a flag does not exist, then we say that the cutoff is zero and set $F_{\beta_0, \dots, \beta_m}^{\alpha} = 0$.

We define the number $(\alpha; \beta_0, \dots, \beta_{m-1}, \beta_m)$ to be the maximal codimension in which the flags $F_{\beta_0, \dots, \beta_{m-1}}$ and $F_{\alpha_0, \dots, \alpha_{m-1}}$ coincide. More precisely, we define $(\alpha; \beta_0, \dots, \beta_m) = k$ if $\beta_0 = \alpha_0, \dots, \beta_k = \alpha_k$ for $k < m-1$ and $\beta_{k+1} \neq \alpha_{k+1}$. We define $(\alpha; \beta_0, \dots, \beta_m) = m-1$ if $\beta_k = \alpha_k$ for $k = 0, \dots, m-1$. We define $(\alpha; \beta_0, \dots, \beta_m) = 0$ if $F_{\beta_0, \dots, \beta_m}^{\alpha} = 0$.

For $\beta \succ \alpha$, define the map $A_{\alpha, \beta} : V_{\beta} \rightarrow V_{\alpha}$ by the rule

$$(3.18) \quad A_{\alpha, \beta} : F_{\beta_0, \dots, \beta_{m-1}, \beta_m = \beta} \otimes w \mapsto (-1)^k F_{\beta_0, \dots, \beta_{m-1}, \beta_m}^{\alpha} \otimes \sum B^i(w),$$

where $k = (\alpha; \beta_0, \dots, \beta_m)$ and the sum is over all $i \in I_1(\Gamma)$ such that $i \geq \alpha_{k+1}$ and $i \not\geq \alpha_k$. Another form of formula (3.18) is

$$A_{\alpha, \beta} : F_{\beta_0, \dots, \beta_{m-1}, \beta_m = \beta} \otimes w \mapsto (-1)^k F_{\beta_0, \dots, \beta_k, \alpha_{k+1}, \dots, \alpha_{m-1} = \alpha} \otimes \sum B^i(w),$$

where the sum is over $i \in I_1(\Gamma)$ such that $i \wedge \beta_k = \beta_{k+1}$, $i \wedge \alpha_{k+1} = \beta_{k+2}$, $\dots$, $i \wedge \alpha_{m-1} = \beta_m$. Here $k = (\alpha; \beta_0, \dots, \beta_m)$ and $F_{\beta_0, \dots, \beta_k, \alpha_{k+1}, \dots, \alpha_{m-1}} = F_{\beta_0, \dots, \beta_{m-1}, \beta}^{\alpha}$.

Proposition 3.7.

- (i) The spaces V_{α} and operators $A_{\alpha, \beta}$ form a quiver $\mathcal{V}_{\Gamma} \in \text{Qui}_{\Gamma}$.
- (ii) The quiver $\mathcal{V}_{\Gamma}$ is isomorphic to $\mathbb{J}_{0,!}(\mathcal{W}_{\Gamma^0})$.
- (iii) For any quiver $\mathcal{U} = \{U_{\alpha}, C_{\alpha, \beta}\} \in \text{Qui}_{\Gamma}$ and a morphism $\varphi \in \text{Hom}_{\text{Qui}_{\Gamma^0}}(\mathcal{W}_{\Gamma^0}, \mathbb{J}_0^*(\mathcal{U}))$, the adjoint morphism $\alpha_{\mathcal{W}_{\Gamma^0}, \mathcal{U}}(\varphi) \in \text{Hom}_{\text{Qui}}(\mathbb{J}_{0,!}(\mathcal{W}_{\Gamma^0}), \mathcal{U})$ is given by the relations

$$(3.19) \quad \alpha_{\mathcal{W}_{\Gamma^0}, \mathcal{U}}(\varphi) : \begin{array}{ll} F_{\emptyset} \otimes w & \mapsto \varphi(w) \\ F_{\alpha_0, \dots, \alpha_m} \otimes w & \mapsto (-1)^{m-1} C_{\alpha_m \alpha_{m-1}} \cdots C_{\alpha_1 \alpha_0} \varphi(w), \quad m \geq 1. \end{array}$$

- (iii) The functor $\mathbb{J}_{0,!} : \text{Qui}_{\Gamma^0} \rightarrow \text{Qui}_{\Gamma}$ is exact. □

For a level zero quiver $\mathcal{W}_{\Gamma^0} = \{W, B^i\}$, define a quiver $\tilde{\mathcal{V}}_{\Gamma} = \{\tilde{V}_{\alpha}, \tilde{A}_{\alpha, \beta}\} \in \text{Qui}_{\Gamma}$ as follows.

Let $\mathcal{A}(\Gamma)$ be the Orlik-Solomon algebra of the graph Γ . For $\alpha \in I(\Gamma)$, define the projection operator $P_\alpha : \mathcal{A}(\Gamma) \rightarrow \mathcal{A}(\Gamma)$ by the formula

$$P_\alpha(H_{j_1}, \dots, H_{j_m}) = \begin{cases} (H_{j_1}, \dots, H_{j_m}), & \text{if } j_1 \wedge \dots \wedge j_m = \alpha, \\ 0, & \text{otherwise} \end{cases}$$

Set $\tilde{V}_\alpha = P_\alpha(\mathcal{A}) \otimes W$. Define the linear maps $\tilde{A}_{\alpha,\beta} : \tilde{V}_\beta \rightarrow \tilde{V}_\alpha$ by the formula

$$(3.20) \quad \tilde{A}_{\alpha,\beta} : (H_{j_1}, \dots, H_{j_m}) \otimes w \mapsto \sum_{k=1}^m (-1)^{k-1} P_\alpha((H_{j_1}, \dots, \hat{H}_{j_k}, \dots, H_{j_m})) \otimes w$$

if $\alpha \succ \beta$, and by the formula

$$(3.21) \quad \tilde{A}_{\alpha,\beta} : (H_{j_1}, \dots, H_{j_m}) \otimes w \mapsto \sum_{k \in I_1(\Gamma)} P_\alpha((H_k, H_{j_1}, \dots, H_{j_m})) \otimes B^k(w).$$

if $\beta \succ \alpha$.

Proposition 3.8.

- (i) The spaces $\tilde{V}_\alpha$ and linear maps $\tilde{A}_{\alpha,\beta}$ form a quiver $\tilde{\mathcal{V}}_\Gamma \in \text{Qui}_\Gamma$.
- (ii) The quiver $\tilde{\mathcal{V}}_\Gamma$ is isomorphic to $\mathbb{J}_{0,*}(\mathcal{W}_{\Gamma^0})$.
- (iii) The morphism $s_0 : \mathbb{J}_{0,!}(\mathcal{W}_{\Gamma^0}) \rightarrow \mathbb{J}_{0,*}(\mathcal{W}_{\Gamma^0})$ is given by the formula

$$(3.22) \quad s_0 : F_{\alpha_0, \dots, \alpha_m} \otimes w \mapsto \sum (H_{j_1}, \dots, H_{j_m}) \otimes B^{j_m} \dots B^{j_1}(w),$$

where the sum is taken over all tuples $(H_{j_1}, \dots, H_{j_m})$, such that $H_{j_1} \supset \overline{X}_{\alpha_1}, \dots, H_{j_m} \supset \overline{X}_{\alpha_m}$.

- (iv) The functor $\mathbb{J}_{0,*} : \text{Qui}_{\Gamma^0} \rightarrow \text{Qui}_\Gamma$ is exact. $\square$

The morphism $s_0 : \mathbb{J}_{0,!}(\mathcal{W}_{\Gamma^0}) \rightarrow \mathbb{J}_{0,*}(\mathcal{W}_{\Gamma^0})$ is called *the quiver Shapovalov map*. It is a matrix generalization of the Shapovalov map S_a from the flag complex to the Aomoto complex, see Section 2.5.

Identifying $\mathcal{A}(\mathcal{C})$ with $\mathcal{F}^*(\mathcal{C})$, as in Section 2.5, we can consider the map s_0 as a bilinear form on the vector space $\mathcal{F}(\mathcal{C})$ with values in $\text{End } W$. This *quiver Shapovalov form* is given by the formula:

$$s_0(F_{\alpha_0, \alpha_1, \dots, \alpha_m}, F_{\alpha'_0, \alpha'_1, \dots, \alpha'_m}) = \sum_{\sigma \in S_m} (-1)^{l(\sigma)} \sum B^{j_m} \dots B^{j_1},$$

where the inner sum is taken over all $j_1, \dots, j_m \in J(\mathcal{C})$, such that $H_{j_1} \supset \overline{X}_{\alpha_1}, \dots, H_{j_m} \supset \overline{X}_{\alpha_m}$, and $H_{j_{\sigma(1)}} \supset \overline{X}_{\alpha'_1}, \dots, H_{j_{\sigma(m)}} \supset \overline{X}_{\alpha'_m}$.

If all operators $B^j : W \rightarrow W$, $j \in J(\mathcal{C})$, commute, then the quiver Shapovalov form is symmetric.

3.7. Monodromy operators for quivers $\mathbb{J}_{0,!}(\mathcal{W}_{\Gamma^0})$ and $\mathbb{J}_{0,*}(\mathcal{W}_{\Gamma^0})$. Suppose that each operator B^i of the level zero quiver $\mathcal{W}_{\Gamma^0} = \{W, B^i\}$ has a single eigenvalue λ_i . We call the collection $\{\lambda_i\}$, $i \in I_1(\Gamma)$, *the spectrum* of $\mathcal{W}_{\Gamma^0}$. For $\alpha \in I(\Gamma)$, set

$$(3.23) \quad \lambda_\alpha = \sum_{i \in I_1(\Gamma), i \geq \alpha} \lambda_i.$$

We call the spectrum $\{\lambda_i\}$ *non-resonant* if

$$(3.24) \quad \lambda_\alpha \notin \mathbb{Z} \setminus \{0\} \quad \text{for all } \alpha \in I(\Gamma).$$

Let $\mathbb{J}_{0,!}(\mathcal{W}_{\Gamma^0}) = \{V_\alpha, A_{\alpha,\beta}\}$ and $\mathbb{J}_{0,*}(\mathcal{W}_{\Gamma^0}) = \{\tilde{V}_\alpha, \tilde{A}_{\alpha,\beta}\}$.

Proposition 3.9.

- (i) For $\alpha \in I(\Gamma)$ and the quiver $\mathbb{J}_{0,!}(\mathcal{W}_{\Gamma^0})$, consider the central element S_α of the local algebra $\mathcal{O}_\alpha(\mathbb{J}_{0,!}(\mathcal{W}_{\Gamma^0}))$, see (3.4). Then S_α is the scalar operator with eigenvalue λ_α .
- (ii) For $\alpha \in I(\Gamma)$, $\alpha \succ \beta$, and the quiver $\mathbb{J}_{0,!}(\mathcal{W}_{\Gamma^0})$, consider the operator $A_\alpha^\beta = A_{\alpha,\beta}A_{\beta,\alpha} : V_\alpha \rightarrow V_\alpha$. Then A_α^β is scalar with eigenvalue $\lambda_\beta - \lambda_\alpha$.
- (iii) For $\alpha \in I(\Gamma)$ and the quiver $\mathbb{J}_{0,!}(\mathcal{W}_{\Gamma^0})$, consider the operators T_α and $\bar{T}_\alpha$, defined in (3.5) and (3.6), respectively. Then every eigenvalue of T_α is either 0 or λ_α and every eigenvalue of $\bar{T}_\alpha$ is either 0 or λ_α .

Proposition 3.10.

- (i) For $\alpha \in I(\Gamma)$ and the quiver $\mathbb{J}_{0,*}(\mathcal{W}_{\Gamma^0})$, consider the central element S_α of the local algebra $\mathcal{O}_\alpha(\mathbb{J}_{0,*}(\mathcal{W}_{\Gamma^0}))$, see (3.4). Then S_α is the scalar operator with eigenvalue λ_α .
- (ii) For $\alpha \in I(\Gamma)$, $\alpha \succ \beta$, and the quiver $\mathbb{J}_{0,*}(\mathcal{W}_{\Gamma^0})$, consider the operator $\tilde{A}_\alpha^\beta = \tilde{A}_{\alpha,\beta}\tilde{A}_{\beta,\alpha} : \tilde{V}_\alpha \rightarrow \tilde{V}_\alpha$. Then $\tilde{A}_\alpha^\beta$ is scalar with eigenvalue $\lambda_\beta - \lambda_\alpha$.
- (iii) For $\alpha \in I(\Gamma)$ and the quiver $\mathbb{J}_{0,*}(\mathcal{W}_{\Gamma^0})$, consider the operators T_α and $\bar{T}_\alpha$, defined in (3.5) and (3.6), respectively. Then every eigenvalue of T_α is either 0 or λ_α and every eigenvalue of $\bar{T}_\alpha$ is either 0 or λ_α .

Proposition 3.11. Let $\mathcal{C}$ be a central arrangement (that is, all hyperplanes of $\mathcal{C}$ contain $0 \in \mathbb{C}^N$). Then for each of the quivers $\mathbb{J}_{0,!}(\mathcal{W}_{\Gamma^0})$ and $\mathbb{J}_{0,*}(\mathcal{W}_{\Gamma^0})$, the operator S , defined in (3.8), has a single eigenvalue which is equal to $\lambda_\infty = \sum_{j \in J(\mathcal{C})} \lambda_j$.

Corollary 3.12.

- (i) Let $\mathcal{W}_{\Gamma^0} = \{W, B^i\}$ be a level zero quiver with non-resonant spectrum. Then the quivers $\mathbb{J}_{0,!}(\mathcal{W}_{\Gamma^0})$, $\mathbb{J}_{0,*}(\mathcal{W}_{\Gamma^0})$, and $\mathbb{J}_{0,!*}(\mathcal{W}_{\Gamma^0})$ are non-resonant. The same conclusion is valid if all operators B^i are close to zero.
- (ii) Let $\mathcal{C}$ be a central arrangement and $\mathcal{W}_{\Gamma^0} = \{W, B^i\}$ a level zero quiver with non-resonant spectrum. Then the quivers $\mathbb{J}_{0,!}(\mathcal{W}_{\Gamma^0})$, $\mathbb{J}_{0,*}(\mathcal{W}_{\Gamma^0})$ and $\mathbb{J}_{0,!*}(\mathcal{W}_{\Gamma^0})$ are strongly non-resonant. The same conclusion is valid if all operators B^i are close to zero.

Statements of the corollary for level zero quivers with non-resonant spectrum follow directly from Propositions 3.9-3.11. If $\mathcal{W}_{\Gamma^0} = \{W, B^i\}$ is a level zero quiver with operators B^i close to zero, then all monodromy operators of quivers $\mathbb{J}_{0,!}(\mathcal{W}_{\Gamma^0})$, $\mathbb{J}_{0,*}(\mathcal{W}_{\Gamma^0})$, and $\mathbb{J}_{0,!*}(\mathcal{W}_{\Gamma^0})$ are close to zero and their eigenvalues belong to some non-resonant sets. $\square$

Propositions 3.9– 3.11 are proved in Section 7.

3.8. Example. Let X be $\mathbb{C}^2$ and $\mathcal{C} = \{H_1, H_2, H_3\}$ the arrangement of three lines, considered in Section 3.5.

In notations of Sections 2.5 and 3.5, the flag complex $\mathcal{F}(\mathcal{C})$ of the arrangement has seven flags $F_\emptyset, F_{\emptyset,\alpha_1}, F_{\emptyset,\alpha_2}, F_{\emptyset,\alpha_3}, F_{\emptyset,\alpha_1,\beta}, F_{\emptyset,\alpha_2,\beta}, F_{\emptyset,\alpha_3,\beta}$, subject to the relation

$$F_{\emptyset,\alpha_1,\beta} + F_{\emptyset,\alpha_2,\beta} + F_{\emptyset,\alpha_3,\beta} = 0.$$

We have $\mathcal{F}_\emptyset = \mathbb{C} \cdot F_\emptyset$, $\mathcal{F}_{\alpha_i} = \mathbb{C} \cdot F_{\emptyset, \alpha_i}$, $i = 1, 2, 3$, and $\mathcal{F}_\beta$ is generated by the flags $F_{\emptyset, \alpha_i, \beta}$, $i = 1, 2, 3$.

The Orlik–Solomon algebra $\mathcal{A}(\mathcal{C})$ is generated by symbols 1 , (H_i) with $i = 1, 2, 3$, and by symbols (H_i, H_j) , where $i, j = 1, 2, 3$. We have $(H_i, H_j) = -(H_j, H_i)$ and

$$(H_1, H_2) - (H_1, H_3) + (H_2, H_3) = 0.$$

Let $\mathcal{V}_{\Gamma^0}$ be a level zero quiver as in Section 3.5, (iii).

The direct image $\mathbb{J}_{0,!}(\mathcal{V}_{\Gamma^2})$ is the quiver $\{W_\emptyset, W_{\alpha_i}, W_\beta, B_{\emptyset, \alpha_i}, B_{\alpha_i, \emptyset}, B_{\beta, \alpha_i}, B_{\alpha_i, \beta}\}$ of Γ , where $W_\gamma = \mathcal{F}_\gamma \otimes V$ for all $\gamma \in I(\Gamma)$, and

$$B_{\alpha_i, \emptyset}(F_\emptyset \otimes v) = F_{\emptyset, \alpha_i} \otimes v, \quad B_{\emptyset, \alpha_i}(F_{\emptyset, \alpha_i} \otimes v) = F_\emptyset \otimes A^i(v),$$

$$\begin{aligned} B_{\beta, \alpha_i}(F_{\emptyset, \alpha_i} \otimes v) &= -F_{\emptyset, \alpha_i, \beta} \otimes v, \\ B_{\alpha_i, \beta}(F_{\emptyset, \alpha_i, \beta} \otimes v) &= F_{\emptyset, \alpha_i} \otimes \sum_{j, j \neq i} A^j(v), \\ B_{\alpha_j, \beta}(F_{\emptyset, \alpha_i, \beta} \otimes v) &= -F_{\emptyset, \alpha_j} \otimes A^i(v), \quad j \neq i. \end{aligned}$$

The direct image $\mathbb{J}_{0,*}(\mathcal{V}_{\Gamma^0})$ is the quiver $\{U_\emptyset, U_{\alpha_i}, U_\beta, C_{\emptyset, \alpha_i}, C_{\alpha_i, \emptyset}, C_{\beta, \alpha_i}, C_{\alpha_i, \beta}\}$ of Γ , where $U_\emptyset = \mathbb{C} \cdot 1 \otimes V$, $U_{\alpha_i} = \mathbb{C} \cdot (H_i) \otimes V$, and U_β is generated by the vectors $(H_i, H_j) \otimes v$, $i, j = 1, 2, 3$, $v \in V$. We have

$$C_{\alpha_i, \emptyset}(1 \otimes v) = (H_i) \otimes A^i(v), \quad C_{\emptyset, \alpha_i}((H_i) \otimes v) = 1 \otimes v,$$

$$C_{\beta, \alpha_i}((H_i) \otimes v) = \sum_{j, j \neq i} (H_j, H_i) \otimes A^j(v),$$

$$C_{\alpha_i, \beta}((H_j, H_i) \otimes v) = (H_i) \otimes v, \quad j \neq i,$$

$$C_{\alpha_i, \beta}((H_j, H_k) \otimes v) = 0, \quad i \notin \{j, k\}.$$

The Shapovalov map $s_0 : \mathbb{J}_{0,!}(\mathcal{V}_{\Gamma^0}) \rightarrow \mathbb{J}_{0,*}(\mathcal{V}_{\Gamma^0})$ is given by the formulas:

$$s_0(F_\emptyset \otimes v) = v, \quad s_0(F_{\emptyset, \alpha_i} \otimes v) = (H_i) \otimes A^i(v),$$

$$s_0(F_{\emptyset, \alpha_i, \beta} \otimes v) = \sum_{j, j \neq i} (H_i, H_j) \otimes A^j A^i(v).$$

The quiver $\mathbb{J}_{0,!}(\mathcal{V}_{\Gamma^0})$ is the subquiver of $\mathbb{J}_{0,*}(\mathcal{V}_{\Gamma^0})$, generated by the vectors $1 \otimes v$, $(H_i) \otimes A^i(v)$, $\sum_{j, j \neq i} (H_i, H_j) \otimes A^j A^i(v)$, where $i = 1, 2, 3$ and $v \in V$.

Suppose that each of the operators $A^i : V \rightarrow V$ of the level 0 quiver $\mathcal{V}_{\Gamma^0}$ has a single eigenvalue λ_i . Then the quivers $\mathbb{J}_{0,!}(\mathcal{V}_{\Gamma^0})$, $\mathbb{J}_{0,*}(\mathcal{V}_{\Gamma^0})$ and $\mathbb{J}_{0,!}(\mathcal{V}_{\Gamma^0})$ are strongly non-resonant if $\lambda_1 + \lambda_2 + \lambda_3 \notin \mathbb{Z} \setminus \{0\}$ and $\lambda_i \notin \mathbb{Z} \setminus \{0\}$ for all $i = 1, 2, 3$.

4. QUIVER $\mathcal{D}$ -MODULES

4.1. Modules over rings of differential operators. Here we briefly recall the basic terminology of modules over rings of differential operators [K1, K4, B].

Let Y be a smooth algebraic variety over $\mathbb{C}$. We denote by $\mathcal{D}_Y$ the sheaf of rings of linear algebraic differential operators over Y . Let M be a sheaf of modules over the sheaf $\mathcal{D}_Y$. We call M a $\mathcal{D}_Y$ -module. A $\mathcal{D}_Y$ -module M is called coherent if the sheaf M can be locally presented as the cokernel of a morphism of free $\mathcal{D}_Y$ -modules of finite rank. We denote by $\mathcal{M}_Y$ the category of coherent $\mathcal{D}_Y$ -modules. If Y is affine, then the space $\overline{M} = \Gamma(Y, M)$ of global sections of a coherent $\mathcal{D}_Y$ -module M is a module over the ring $D_M = \Gamma(Y, \mathcal{D}_Y)$ of global algebraic differential operators over Y , and the sheaf M is determined by localizations of $\overline{M}$, that is, for any open affine $U \subset Y$, we have

$$\Gamma(U, M) = D_U \otimes_{D_Y} \overline{M}.$$

Let U be an affine open subset in an algebraic variety Y and T^*U the cotangent bundle of U . A coherent D_U -module M_U has singular support $ss.M \subset T^*U$, which is defined as follows. Denote by $D_U^k \subset \mathcal{D}_U$ the subspace of differential operators of order not greater than k . The spaces D_U^k define an increasing filtration of the ring D_U such that the corresponding graded ring $Gr D_U$ is a commutative ring isomorphic to the polynomial ring $\mathbb{C}[T^*U]$. An increasing filtration $\{M_U^k\}$ of the module M_U is called *good* if it agrees with the filtration $\{D_U^k\}$ of the ring D_U :

- $D_U^k M_U^l \subset M_U^{k+l}$ for all k, l ;
- $D_U^k M_U^l = M_U^{k+l}$ for all $l \geq l_0, k \geq 0$;
- All M_U^k are finitely generated $\mathcal{O}_U$ -modules.

For a good filtration, the graded space $gr M_U$ is a coherent module over $Gr D_U$ and thus defines an algebraic variety $ss.M_U \subset T^*U$. More precisely, the variety is defined by the radical of the annihilator ideal $Ann(gr M_U)$. The variety is called *the singular support or characteristic variety* of the D_U -module M_U .

Bernstein's theorem [Bn] says that a coherent $\mathcal{D}_Y$ -module admits a good filtration and the singular support of M_U does not depend on the choice of the good filtration. This implies that the characteristic variety for a coherent $\mathcal{D}_Y$ -module M_Y on a nonsingular variety Y is well defined. The characteristic variety is an involutive subvariety of T^*Y with respect to the canonical symplectic structure on T^*Y , thus its dimension is not less than the dimension of Y . A $\mathcal{D}_Y$ -module M_Y is called *holonomic* if its characteristic variety is lagrangian. Denote by $\mathcal{M}_Y^{hol} \subset \mathcal{M}_Y$ the full subcategory of sheaves of holonomic $\mathcal{D}_Y$ -modules.

Let $\mathcal{O}_Y$ denote the structure sheaf of regular functions on Y , $\mathcal{O}_Y^{an}$ the sheaf of holomorphic functions on Y ; Ω_Y^k the sheaf of k -forms on Y ; $\Omega_Y^{k,an}$ the sheaf of holomorphic k -forms on Y .

The simplest examples of holonomic $\mathcal{D}_Y$ -modules are provided by flat algebraic connections in algebraic vector bundles over Y . Recall that a connection ∇ in a locally free $\mathcal{O}_Y$ -module $\mathcal{S}$ of local sections of a vector bundle S is a $\mathbb{C}$ -linear map

$$\nabla : \mathcal{S} \rightarrow \Omega_Y^1 \otimes_{\mathcal{O}_Y} \mathcal{S},$$

satisfying the condition $\nabla(fs) = df \otimes s + f\nabla(s)$ for $f \in \mathcal{O}_Y$ and $s \in \mathcal{S}$. The curvature $\nabla^{(2)}$ of the connection ∇ ,

$$\nabla^{(2)} : \Omega_Y^1 \otimes_{\mathcal{O}_Y} \mathcal{S} \rightarrow \Omega_Y^2 \otimes_{\mathcal{O}_Y} \mathcal{S},$$

is determined by conditions $\nabla^{(2)}(\omega \otimes s) = d\omega \otimes s - \omega \wedge \nabla(s)$ for any $\omega \in \Omega_Y^1$ and $s \in \mathcal{S}$. The connection is flat if its curvature is zero.

A flat connection $(\nabla, \mathcal{S})$ can be canonically extended to a holomorphic flat connection, $(\nabla^{an}, \mathcal{O}_Y^{an} \otimes_{\mathcal{O}_Y} \mathcal{S})$.

A flat connection $(\nabla, \mathcal{S})$ provides the $\mathcal{O}_Y$ -module $\mathcal{S}$ with a structure of a $\mathcal{D}_Y$ -module $\mathcal{S}_\nabla$, where the action of a vector field ξ is given by the relation $\xi(s) = i_\xi(\nabla(s))$.

Let Y be a stratified algebraic variety with smooth algebraic strata Y_j , $j \in J(Y)$. It means that Y is presented as a finite disjoint union of strata Y_j , such that the closure of each stratum is a union of strata. Denote by $\widetilde{\mathcal{M}}_Y^{hol} \subset \mathcal{M}_Y$ the full subcategory of holonomic $\mathcal{D}_Y$ -modules M whose singular support is contained in the union of the conormal bundles of strata of Y , $ss.M \subset \bigcup_{j \in J(Y)} T_{Y_j}^* Y$. This category is abelian and is closed under extensions. If $M \in \widetilde{\mathcal{M}}_Y^{hol}$, then we say that M is *smooth along the stratification of Y* .

4.2. Quiver $\mathcal{D}_X$ -modules. Let $\mathcal{C}$ be an arrangement in $X = \mathbb{C}^N$. Let $\{f_{\beta,\alpha}, \xi_{\alpha,\beta} \mid \alpha, \beta \in I(\Gamma), \alpha \succ \beta\}$ be an edge framing of $\mathcal{C}$.

Let $\mathcal{V}$ be a quiver of the graph Γ of $\mathcal{C}$. We define *the associated quiver $\mathcal{D}_X$ -module $E\mathcal{V}$* as the quotient of the free sheaf $\mathcal{R}^0 E\mathcal{V} = \mathcal{D}_X \otimes_{\mathbb{C}} (\bigoplus_{\alpha \in I(\Gamma)} \overline{\Omega}_\alpha \otimes V_\alpha)$ of $\mathcal{D}_X$ -modules over its subsheaf $\mathcal{R}'E\mathcal{V}$ of coherent $\mathcal{D}_X$ -modules, whose global sections are the following global sections of $\mathcal{R}^0 E\mathcal{V}$:

(i) the sections

$$(4.25) \quad u \xi \otimes \overline{\omega}_\alpha \otimes v_\alpha - \sum_{\beta, \alpha \succ \beta} \langle \xi, f_{\beta,\alpha} \rangle u \otimes \pi_{\beta,\alpha}(\overline{\omega}_\alpha) \otimes A_{\beta,\alpha}(v_\alpha),$$

where $\alpha \in I(\Gamma)$, $u \otimes \overline{\omega}_\alpha \otimes v_\alpha \in D_X \otimes_{\mathbb{C}} \overline{\Omega}_\alpha \otimes V_\alpha$ and $\xi \in T_\alpha$,

(ii) the sections

$$(4.26) \quad u f \otimes \overline{\omega}_\alpha \otimes v_\alpha - \sum_{\beta, \beta \succ \alpha} \langle \xi_{\beta,\alpha}, f \rangle u \otimes \pi_{\beta,\alpha}(\overline{\omega}_\alpha) \otimes A_{\beta,\alpha}(v_\alpha),$$

where $\alpha \in I(\Gamma)$, $u \otimes \overline{\omega}_\alpha \otimes v_\alpha \in D_X \otimes_{\mathbb{C}} \overline{\Omega}_\alpha \otimes V_\alpha$ and $f \in F_\alpha$.

It is easy to see that the space generated by sections in (4.25)–(4.26) does not depend on the choice of the edge framing. Thus the sheaf $E\mathcal{V}$ is determined by the quiver $\mathcal{V}$ only.

For any $\alpha \in I(\Gamma)$, $\overline{\omega}_\alpha \in \overline{\Omega}_\alpha$, $v_\alpha \in V_\alpha$, we denote by $\overline{\omega}_\alpha \otimes v_\alpha$ the image in $\Gamma(X, E\mathcal{V})$ of the global section $1 \otimes \overline{\omega}_\alpha \otimes v_\alpha \in \Gamma(X, \mathcal{R}^0 E\mathcal{V})$ with respect to the canonical projection $\mathcal{R}^0 E\mathcal{V} \rightarrow E\mathcal{V}$. The vectors $\overline{\omega}_\alpha \otimes v_\alpha$, where $\alpha \in I(\Gamma)$, $\overline{\omega}_\alpha \in \overline{\Omega}_\alpha$, $v_\alpha \in V_\alpha$, generate $\Gamma(X, E\mathcal{V})$ as $\mathcal{D}_X$ -module.

Remark. In [Kh, KhS2] relations (4.25)–(4.26) were written in a different but equivalent form. Namely, let $\widehat{\Omega}_\alpha$ be the one-dimensional complex vector space of $l(\alpha)$ -exterior forms

$df_1 \wedge df_2 \wedge \dots \wedge df_{l(\alpha)}$ with $f_1, \dots, f_{l(\alpha)} \in F_\alpha$. For any $\alpha \in I(\Gamma)$, choose an element $\widehat{\omega}_\alpha \in \widehat{\Omega}_\alpha$. Then the relations of (4.25)–(4.26) are equivalent to the next relations,

$$(4.27) \quad \begin{aligned} \xi(\overline{\omega}_\alpha \otimes v_\alpha) &= \sum_{\beta, \alpha \succ \beta} \frac{i_\xi(\widehat{\omega}_\beta) \wedge \overline{\omega}_\alpha}{\widehat{\omega}_\beta \wedge \overline{\omega}_\beta} \overline{\omega}_\beta \otimes A_{\beta, \alpha}(v_\alpha), \\ f(\overline{\omega}_\alpha \otimes v_\alpha) &= \sum_{\beta, \beta \succ \alpha} \frac{df \wedge \widehat{\omega}_\beta \wedge \overline{\omega}_\alpha}{\widehat{\omega}_\beta \wedge \overline{\omega}_\beta} \overline{\omega}_\beta \otimes A_{\beta, \alpha}(v_\alpha). \end{aligned}$$

The right hand side of formulas (4.27) does not depend on the choice of the forms $\widehat{\omega}_\alpha \in \widehat{\Omega}_\alpha$.

It is easy to see that a morphism of quivers $\phi : \mathcal{V} \rightarrow \mathcal{V}'$ induces a natural morphism $E\phi : E\mathcal{V} \rightarrow E\mathcal{V}'$ of $\mathcal{D}_X$ -modules. Thus the correspondence $E : \mathcal{V} \mapsto E\mathcal{V}$, $E : \phi \mapsto E\phi$ defines a functor from the category Qui_Γ to the category of $\mathcal{D}_X$ -modules.

Choose a vector space structure and a non-degenerate bilinear form $\langle, \rangle$ on $\mathbb{C}^N$. For $\alpha \in I(\Gamma)$, introduce the spaces T'_α and F'_α as in Section 2.2. Let $S(T'_\alpha \oplus F'_\alpha)$ be the symmetric tensor algebra over the space $T'_\alpha \oplus F'_\alpha$. Denote by $M_\alpha = M_\alpha(\mathcal{V})$ the vector space

$$(4.28) \quad M_\alpha = S(T'_\alpha \oplus F'_\alpha) \otimes \overline{\Omega}_\alpha \otimes V_\alpha.$$

Clearly, there is a natural linear map of the vector space M_α to the vector space of global section of $E\mathcal{V}$.

Proposition 4.1. *The vector space of global sections of the $\mathcal{D}_X$ -module $E\mathcal{V}$ is isomorphic to the direct sum of the spaces M_α ,*

$$(4.29) \quad \Gamma(E\mathcal{V}) \simeq \bigoplus_{\alpha \in I(\Gamma)} M_\alpha.$$

The proposition is proved in Section 7.

For $k = 0, \dots, N$, denote by $\overline{V}_k$ the following finite dimensional subspace of global sections of the $\mathcal{D}_X$ -module $E\mathcal{V}$:

$$(4.30) \quad \overline{V}_k = \bigoplus_{\alpha \in I_k(\Gamma)} \overline{\Omega}_\alpha \otimes V_\alpha.$$

For any $m \geq 0$ and l , $0 \leq l \leq N$, set

$$F_p^m E\mathcal{V} = \sum_{k=0}^N \mathcal{D}_X^{m-k} \overline{V}_k, \quad F_p^{m,l} E\mathcal{V} = \sum_{k=l}^N \mathcal{D}_X^{m-k} \overline{V}_k$$

where $\mathcal{D}_X^l$ is the sheaf of differential operators over X of order less than or equal to l . We have

$$\mathcal{D}_X^l F_p^m E\mathcal{V} \subset F_p^{l+m} E\mathcal{V} \quad \text{and} \quad F_p^{m_1} E\mathcal{V} \subset F_p^{m_2} E\mathcal{V}$$

for $m_1 < m_2$, so that the graded factor $\overline{E\mathcal{V}} = \bigoplus_{m \geq 0} F_p^m E\mathcal{V} / F_p^{m-1} E\mathcal{V}$ is a module over $\text{gr}\mathcal{D}_X \simeq \mathbb{C}[T^*X]$. We call the filtration

$$(4.31) \quad 0 = F_p^{-1} E\mathcal{V} \subset F_p^0 E\mathcal{V} \subset F_p^1 E\mathcal{V} \subset \dots \subset F_p^m E\mathcal{V} \subset \dots \subset E\mathcal{V}$$

the principal filtration of the quiver $\mathcal{D}_X$ -module $E\mathcal{V}$. The sheaves $F_p^{m,l}E\mathcal{V}$ form a subfiltration

$$(4.32) \quad \dots \subset F_p^{m-1}E\mathcal{V} = F_p^{m-1,0}E\mathcal{V} \subset F_p^{m,N}E\mathcal{V} \subset \dots \subset F_p^{m,0}E\mathcal{V} = F_p^mE\mathcal{V} \subset \dots$$

of the principal filtration of the quiver $\mathcal{D}_X$ -module $E\mathcal{V}$. We call it *the subtle filtration* of the quiver $\mathcal{D}_X$ -module $E\mathcal{V}$.

Proposition 4.2.

- (i) *The principal filtration (4.31) of the quiver $\mathcal{D}_X$ -module $E\mathcal{V}$ is good.*
- (ii) *The associated graded factor $\overline{E\mathcal{V}} = \text{gr } E\mathcal{V}$ of the sheaf $E\mathcal{V}$ with respect to the principal filtration $F_pE\mathcal{V}$ is isomorphic to*

$$\bigoplus_{\alpha \in I(\Gamma)} \mathcal{O}_{\overline{X}_\alpha} \otimes (S(T'_\alpha) \otimes \overline{\Omega}_\alpha \otimes V_\alpha)$$

as the sheaf of $\mathcal{O}_X$ -modules.

- (iii) *The $\mathcal{D}_X$ -module $E\mathcal{V}$ is holonomic and smooth along the stratification $\{X_\alpha\}$.*

The proposition is proved in Section 7.

4.3. Example.

- (i) Let X be $\mathbb{C}$ with coordinate z . Let $\mathcal{C} = \{H_1, \dots, H_n\}$, where $H_j = \{z - t_j = 0\}$. Denote the vertices of the graph of the arrangement by the symbols $\emptyset$, and j , $j = 1, \dots, n$. Let $\mathcal{V} = \{V_\emptyset, V_j, A_{\emptyset,j}, A_{j,\emptyset}\}$ be a quiver of Γ .

The vector space $\Gamma(X, E\mathcal{V})$ of global sections of the sheaf $E\mathcal{V}$ is generated over D_X by the vectors $dz \otimes v_\emptyset \in \overline{\Omega}_\emptyset \otimes V_\emptyset$, and $1 \otimes v_j \in \overline{\Omega}_j \otimes V_j$, subject to the relations:

$$\partial_z(dz \otimes v_\emptyset) = \sum_{j=1}^n 1 \otimes A_{j,\emptyset}(v_\emptyset), \quad (z - t_j)(1 \otimes v_j) = dz \otimes A_{\emptyset,j}(v_j),$$

where $\partial_z = \frac{d}{dz}$.

The vectors

$$\begin{aligned} z^k(dz \otimes v_\emptyset), \quad k \geq 0, \quad v_\emptyset \in V_\emptyset, \\ \partial_z^l(1 \otimes v_j), \quad l \geq 0, \quad v_j \in V_j, \end{aligned}$$

form a $\mathbb{C}$ -basis of the vector space $\Gamma(X, E\mathcal{V})$.

Let $\{F_p^l E\mathcal{V}\}$ be the principal filtration of the $\mathcal{D}_X$ -module $E\mathcal{V}$.

The vectors $z^k(dz \otimes v_\emptyset)$, $k \geq 0$, form a $\mathbb{C}$ -basis of the space of global sections of the sheaf $F_p^0 E\mathcal{V}$.

For any $l > 0$, the vectors $\partial_z^m(1 \otimes v_j)$, $0 \leq m \leq l$, and $z^k(dz \otimes v_\emptyset)$, $k \geq 0$, form a $\mathbb{C}$ -basis of the space of global sections of the sheaf $F_p^l E\mathcal{V}$.

- (ii) Let X be $\mathbb{C}^2$ with coordinates z_1, z_2 . Let $\mathcal{C} = \{H_1, \dots, H_n\}$ be the arrangement of n lines, $H_j = \{z_2 + k_j z_1 = 0\}$, $j = 1, \dots, n$, intersecting at one point. Denote the vertices of the graph of the arrangement by the symbols $\emptyset$, α_j , where $j = 1, \dots, n$, and β , such that X_β is the point $(0, 0)$.

Let $\mathcal{V} = \{V_\emptyset, V_{\alpha_j}, V_\beta, A_{\emptyset,\alpha_j}, A_{\alpha_j,\emptyset}, A_{\beta,\alpha_j}, A_{\alpha_j,\beta}\}$ be a quiver.

The vector space $\Gamma(X, E\mathcal{V})$ of global sections of the sheaf $E\mathcal{V}$ is generated over D_X by the vectors $dz_1 \wedge dz_2 \otimes v_\emptyset$, where $v_\emptyset \in V_\emptyset$, the vectors $dz_1 \otimes v_{\alpha_j}$, where $v_{\alpha_j} \in V_{\alpha_j}$, and the vectors $1 \otimes v_\beta$, where $v_\beta \in V_\beta$, subject to the following relations:

$$\begin{aligned} \partial_{z_1}(dz_1 \wedge dz_2 \otimes v_\emptyset) &= \sum_{j=1}^n k_j dz_1 \otimes A_{\alpha_j, \emptyset}(v_\emptyset), & \partial_{z_2}(dz_1 \wedge dz_2 \otimes v_\emptyset) &= \sum_{j=1}^n dz_1 \otimes A_{\alpha_j, \emptyset}(v_\emptyset), \\ (z_2 + k_j z_1)(dz_1 \otimes v_{\alpha_j}) &= 1 \otimes A_{\emptyset, \alpha_j}(v_{\alpha_j}), & (\partial_{z_1} - k_j \partial_{z_2})(dz_1 \otimes v_{\alpha_j}) &= 1 \otimes A_{\beta, \alpha_j}(v_{\alpha_j}), \\ z_1(1 \otimes v_\beta) &= \sum_{j=1}^n dz_1 \otimes A_{\beta, \alpha_j}(v_\beta), & z_2(1 \otimes v_\beta) &= - \sum_{j=1}^n k_j dz_1 \otimes A_{\beta, \alpha_j}(v_\beta) \end{aligned}$$

The vectors:

$$\begin{aligned} z_1^{k_1} z_2^{k_2} (dz_1 \wedge dz_2 \otimes v_\emptyset) , & \quad k_1, k_2 \geq 0 , \\ z_1^{k_1} \partial_{z_2}^{k_2} (dz_1 \otimes v_{\alpha_j}) , & \quad k_1, k_2 \geq 0 , \quad j = 1, \dots, n , \\ \partial_{z_1}^{k_1} \partial_{z_2}^{k_2} (1 \otimes v_\beta) , & \quad k_1, k_2 \geq 0 , \end{aligned}$$

form a $\mathbb{C}$ -basis of the vector space $\Gamma(X, E\mathcal{V})$.

The vectors $z_1^{k_1} z_2^{k_2} (dz_1 \wedge dz_2 \otimes v_\emptyset)$, $k_1, k_2 \geq 0$, form a $\mathbb{C}$ -basis of the space of global sections of the sheaf $F_p^0 E\mathcal{V}$.

The vectors $z_1^{k_1} z_2^{k_2} (dz_1 \wedge dz_2 \otimes v_\emptyset)$, $k_1, k_2 \geq 0$, and the vectors $dz_1 \otimes v_{\alpha_j} \in V_{\alpha_j}$, $j = 1, \dots, n$, form a $\mathbb{C}$ -basis of the space of global sections of the sheaf $F_p^1 E\mathcal{V}$.

For $l \geq 2$, the vectors $z_1^{k_1} z_2^{k_2} (dz_1 \wedge dz_2 \otimes v_\emptyset)$, $k_1, k_2 \geq 0$, the vectors $z_1^{k_1} \partial_{z_2}^{k_2} (dz_1 \otimes v_{\alpha_j})$, $k_1 \geq 0$, $0 \leq k_2 < l$, $j = 1, \dots, n$, and the vectors $\partial_{z_1}^{k_1} \partial_{z_2}^{k_2} (1 \otimes v_\beta)$, $k_1 + k_2 < l - 1$, form a $\mathbb{C}$ -basis of the space of global sections of the sheaf $F_p^l E\mathcal{V}$.

4.4. Quiver $\mathcal{D}_{X^n}$ -modules. Let X^n be a principal open subset of X , see Section 2.3. Let $\{f_{\beta, \alpha}, \xi_{\alpha, \beta} \mid \alpha, \beta \in I(\Gamma), \alpha \succ \beta\}$ be an edge framing of $\mathcal{C}$, and $\mathcal{V}_{\Gamma^n} = \{V_\alpha, A_{\alpha, \beta}, A_\alpha^\beta\} \in \text{Qui}_{\Gamma^n}$ a quiver of Γ^n .

We define *the associated quiver $\mathcal{D}_{X^n}$ -module $E^n \mathcal{V}_{\Gamma^n}$* as the quotient of the free sheaf $\mathcal{R}^0 E^n \mathcal{V}_{\Gamma^n} = \mathcal{D}_{X^n} \otimes_{\mathbb{C}} (\oplus_{\alpha \in I(\Gamma^n)} \overline{\Omega}_\alpha \otimes V_\alpha)$ of $\mathcal{D}_{X^n}$ -modules over its subsheaf $\mathcal{R}' E^n \mathcal{V}_{\Gamma^n}$ of coherent $\mathcal{D}_{X^n}$ -modules, which is described by its restrictions to open affine subsets $U_{\mathcal{F}_n}$, where $\mathcal{F}_n = \{\overline{f}_\alpha \mid \alpha \in I_{n+1}(\Gamma)\}$ is a level n vertex framing.

The sections $\Gamma(U_{\mathcal{F}_n}, \mathcal{R}' E^n \mathcal{V}_{\Gamma^n})$ of the sheaf $\mathcal{R}' E^n \mathcal{V}_{\Gamma^n}$ over $U_{\mathcal{F}_n}$ are the following sections in $\Gamma(U_{\mathcal{F}_n}, \mathcal{R}^0 E^n \mathcal{V}_{\Gamma^n})$:

(i) the sections

$$(4.33) \quad u \xi \otimes \overline{\omega}_\alpha \otimes v_\alpha - \sum_{\beta, \alpha \succ \beta} \langle \xi, f_{\beta, \alpha} \rangle u \otimes \pi_{\beta, \alpha}(\overline{\omega}_\alpha) \otimes A_{\beta, \alpha}(v_\alpha) ,$$

where $\alpha \in I(\Gamma^n)$ with $l(\alpha) < n$, $u \otimes \overline{\omega}_\alpha \otimes v_\alpha \in D_{U_{\mathcal{F}_n}} \otimes_{\mathbb{C}} \overline{\Omega}_\alpha \otimes V_\alpha$, and $\xi \in T_\alpha$,

(ii) the sections

$$(4.34) \quad \begin{aligned} u \xi \otimes \bar{\omega}_\alpha \otimes v_\alpha - \sum_{\beta, \alpha \succ \beta} u \bar{f}_\beta^{-1} \otimes \left(\langle \xi, \bar{f}_\beta \rangle \bar{\omega}_\alpha \otimes A_\alpha^\beta(v_\alpha) + \right. \\ \left. + \sum_{\gamma, \gamma \succ \beta, \gamma \neq \alpha} \langle \xi, f_{\beta, \alpha} \rangle \langle \xi_{\gamma, \beta}, \bar{f}_\beta \rangle \pi_{\gamma, \beta} \pi_{\beta, \alpha}(\bar{\omega}_\alpha) \otimes A_{\gamma, \delta(\alpha, \gamma)} A_{\delta(\alpha, \gamma), \alpha}(v_\alpha) \right) \end{aligned}$$

where $\alpha \in I(\Gamma^n)$ with $l(\alpha) = n$, $u \otimes \bar{\omega}_\alpha \otimes v_\alpha \in D_{U_{\mathcal{F}_n}} \otimes_{\mathbb{C}} \bar{\Omega}_\alpha \otimes V_\alpha$, and $\xi \in T_\alpha$.

Here $\delta(\alpha, \gamma)$ is the unique (if exists) element of $I(\Gamma^n)$, such that $\delta(\alpha, \gamma) \succ \alpha$ and $\delta(\alpha, \gamma) \succ \gamma$. If it does not exist, the corresponding summand in (4.34) is set to be zero.

(iii) the sections

$$(4.35) \quad u f \otimes \bar{\omega}_\alpha \otimes v_\alpha - \sum_{\beta, \beta \succ \alpha} \langle \xi_{\beta, \alpha}, f \rangle u \otimes \pi_{\beta, \alpha}(\bar{\omega}_\alpha) \otimes A_{\beta, \alpha}(v_\alpha),$$

where $\alpha \in I(\Gamma^n)$, $u \otimes \bar{\omega}_\alpha \otimes v_\alpha \in D_{U_{\mathcal{F}_n}} \otimes_{\mathbb{C}} \bar{\Omega}_\alpha \otimes V_\alpha$, and $f \in F_\alpha$.

One can see that the space generated by sections in (4.33)–(4.35) does not depend on the choice of the edge framing of $\mathcal{C}$. Thus the sheaf $E^n \mathcal{V}_{\Gamma^n}$ is determined by the quiver $\mathcal{V}_{\Gamma^n}$ only.

For any $\alpha \in I(\Gamma^n)$, $\bar{\omega}_\alpha \in \bar{\Omega}_\alpha$, $v_\alpha \in V_\alpha$, we denote by $\bar{\omega}_\alpha \otimes v_\alpha$ the image in $\Gamma(U_{\mathcal{F}_n}, E^n \mathcal{V}_{\Gamma^n})$ of the sections $1 \otimes \bar{\omega}_\alpha \otimes v_\alpha \in \Gamma(U_{\mathcal{F}_n}, \mathcal{R}^0 E^n \mathcal{V}_{\Gamma^n})$ with respect to the canonical projection $\mathcal{R}^0 E^n \mathcal{V}_{\Gamma^n} \rightarrow E^n \mathcal{V}_{\Gamma^n}$.

The vectors $\bar{\omega}_\alpha \otimes v_\alpha$, where $\alpha \in I(\Gamma^n)$, $\bar{\omega}_\alpha \in \bar{\Omega}_\alpha$, $v_\alpha \in V_\alpha$, generate $\Gamma(U_{\mathcal{F}_n}, E^n \mathcal{V}_{\Gamma^n})$ as a $D_{U_{\mathcal{F}_n}}$ -module.

All statements of the previous section have straightforward analogs for the quiver $\mathcal{D}_{X^n}$ -modules on X^n . In particular, the principal filtration $F_p E^n \mathcal{V}_{\Gamma^n}$ of a quiver $\mathcal{D}_{X^n}$ -module $E^n \mathcal{V}_{\Gamma^n}$ is defined by the rule

$$\Gamma(U_{\mathcal{F}_n}, F_p^m E^n \mathcal{V}_{\Gamma^n}) = \sum_{k=0}^n D_{U_{\mathcal{F}_n}}^{m-k} \bar{V}_k,$$

where $\bar{V}_k$ is defined in (4.30). The graded factor $\bar{E}^n \mathcal{V}_{\Gamma^n} = \text{gr } E^n \mathcal{V}_{\Gamma^n}$ of the sheaf $E^n \mathcal{V}_{\Gamma^n}$ with respect to the principal filtration $F_p E^n \mathcal{V}_{\Gamma^n}$ is isomorphic to

$$\bigoplus_{\alpha \in I(\Gamma^n)} \mathcal{O}_{\bar{X}_\alpha \cap X^n} \otimes (S(T'_\alpha) \otimes \bar{\Omega}_\alpha \otimes V_\alpha)$$

as a sheaf of $\mathcal{O}_{X^n}$ -modules. This implies that the space of global sections $\Gamma(X^n, E^n \mathcal{V}_{\Gamma^n})$ of the $\mathcal{D}_{X^n}$ -module $E^n \mathcal{V}_{\Gamma^n}$ is isomorphic to

$$\bigoplus_{\alpha \in I(\Gamma^n)} \mathbb{C}[\bar{X}_\alpha \cap X^n] \otimes (S(T'_\alpha) \otimes \bar{\Omega}_\alpha \otimes V_\alpha).$$

For any $n = 0, 1, \dots, N$, consider the category $Qui_{\Gamma^n, \{G_\beta, \tilde{G}_\beta, G_{\alpha, \beta}\}}$ associated with given collections $\{G_\alpha\}$, $\{\tilde{G}_\beta\}$, and $\{G_{\alpha, \beta}\}$.

Proposition 4.3.

(i) The assignment $\mathcal{V}_{\Gamma^n} \mapsto E^n \mathcal{V}_{\Gamma^n}$ defines a faithful functor $E^n : Qui_{\Gamma^n} \rightarrow \mathcal{M}_{X^n}^{hol}$.

(ii) Let $\mathcal{C}$ be a central arrangement (that is, all hyperplanes of $\mathcal{C}$ contain $0 \in \mathbb{C}^N$). Then the restriction of E^n to the subcategory $Qui_{\Gamma^n, \{G_\beta, \tilde{G}_\beta, G_{\alpha, \beta}\}}$ is full and faithful.

In particular, the restriction of the functor E to the subcategory $Qui_{\Gamma, \{G_\beta, \tilde{G}_\beta, G_{\alpha, \beta}\}}$ defines an equivalence of the category $Qui_{\Gamma, \{G_\beta, \tilde{G}_\beta, G_{\alpha, \beta}\}}$ with a full subcategory of the category $\mathcal{M}_X^{hol}$, closed under extensions.

Proposition 4.3 is proved in Section 7 by induction on n . The following proposition plays the key role in the inductive proof.

Proposition 4.4. *Let $\mathcal{C}$ be a central arrangement. Let $\{G_j \mid j \in J(\mathcal{C})\} \cup \{\tilde{G}_\emptyset\}$ be a collection of weakly non-resonant sets. Let $\mathcal{V} = \{V, A^j\}$ and $\mathcal{W} = \{W, B^j\}$ be level zero quivers, such that*

- (a) *for any $j \in J(\mathcal{C})$ the eigenvalues of A^j and eigenvalues of B^j are in G_j ,*
- (b) *the eigenvalues of $\sum_{j \in J(\mathcal{C})} A^j$ and the eigenvalues of $\sum_{j \in J(\mathcal{C})} B^j$ are in $\hat{G}_\emptyset$.*

Then the vector spaces $\text{Hom}_{Qui_{\Gamma^0}}(\mathcal{V}, \mathcal{W})$ and $\text{Hom}_{\mathcal{D}_{X^0}}(E^0 \mathcal{V}, E^0 \mathcal{W})$ are isomorphic.

Remark. Proposition 4.3 is a generalization of the corresponding statement in [Kh]. Note that the condition of the centrality of the arrangement is missing in [Kh]. This is the mistake, which is in the ignorance of Proposition 4.4 in the inductive proof of the main theorem of [Kh]. The statement of the main theorem of [Kh] is valid for central arrangements.

Proposition 4.4 is proved in Section 7.

Example. Let X^0 be the complement to the union of hyperplanes of the arrangement $\mathcal{C} = \{H_j\}, j \in J(\mathcal{C})$, in $X = \mathbb{C}^N$. Let $z_1, \dots, z_N$ be affine coordinates in X . For $j \in J(\mathcal{C})$, let f_j be an affine function on $\mathbb{C}^N$ defining the hyperplane H_j . Let $\mathcal{V}_{\Gamma^0} = \{V, A^j\} \in Qui_{\Gamma^0}$ be a quiver of level 0. Recall that $A^j : V \rightarrow V$ are linear maps such that for each $\alpha \in I_2(\Gamma)$, $j \succ \alpha$, we have

$$(4.36) \quad [A^j, \sum_{i, i \succ \alpha} A^i] = 0.$$

The associated $\mathcal{D}_{X^0}$ -module $E^0 \mathcal{V}_{\Gamma^0}$ is generated by the vectors $\omega_\emptyset \otimes v$, where $\omega_\emptyset = dz_1 \wedge \dots \wedge dz_N$ and $v \in V$, subject to the relations, labeled by vectors $v \in V$ and constant vector fields ξ on $\mathbb{C}^N$,

$$(4.37) \quad \xi(\omega_\emptyset \otimes v) = \sum_j \frac{\langle \xi, f_j \rangle}{f_j} (\omega_\emptyset \otimes A^j(v)).$$

That means that $E^0 \mathcal{V}_{\Gamma^0}$ is isomorphic to the $\mathcal{D}_{X^0}$ -module associated to the following flat connection.

Consider the trivial bundle on X^0 with fiber V , the $\text{End } V$ -valued differential one-form

$$(4.38) \quad \sum_{j \in J(\mathcal{C})} A^j \otimes d \log f_j,$$

and the associated connection on the trivial bundle. The connection is flat due to (4.36). The connection is regular singular. The $\mathcal{D}_{X^0}$ -module associated with the connection is

isomorphic to $E^0 \mathcal{V}_{\Gamma^0}$. As a sheaf of $\mathcal{O}_{X^0}$ -modules, it is isomorphic to the free sheaf $\mathcal{O}_{X^0} \otimes V$.

4.5. Direct and inverse images of quiver $\mathcal{D}$ -modules. For $0 \leq k \leq l \leq N$, let X^k, X^l be the principal open subsets of X . Since the canonical embedding $j_{l,k} : X^k \rightarrow X^l$ is open, the functors $j_{l,k}^* : \mathcal{M}_{X^l} \rightarrow \mathcal{M}_{X^k}$ and $j_{l,k,*} : \mathcal{M}_{X^k} \rightarrow \mathcal{M}_{X^l}$ of the inverse and direct images of $\mathcal{D}$ -modules are well defined and can be described in the sheaf-theoretical language. Namely, for $M \in \mathcal{M}_{X^l}$, the sheaf $j_{l,k}^*(M)$ is the sheaf-theoretical inverse image $j_{l,k}^*(M)$ of the sheaf M equipped with the natural structure of the sheaf of $\mathcal{D}_{X^k}$ -modules. Also for $N \in \mathcal{M}_{X^k}$, the sheaf $j_{l,k,*}(N)$ is the sheaf-theoretical direct image $(j_{l,k})_*(N)$ of the sheaf N equipped with the natural structure of the sheaf of $\mathcal{D}_{X^l}$ -modules.

More precisely, let M be a coherent $\mathcal{D}_{X^l}$ -module and $U \subset X^k$ an open subset. Then U is an open subset of X^l as well and

$$(4.39) \quad \Gamma(U, j_{l,k}^*(M)) = \Gamma(U, M) .$$

Since the space $\Gamma(U, M)$ is a D_U -module, the relation (4.39) describes the structure of a D_U -module on the space $\Gamma(U, j_{l,k}^*(M))$.

Let now N be a coherent $\mathcal{D}_{X^k}$ -module and $U \subset X^l$ an open subset. Then $j_{l,k}^{-1}(U) = U \cap X^k \subset U$ is an open subset of X^k and

$$(4.40) \quad \Gamma(U, j_{l,k,*}(N)) = \Gamma(j_{l,k}^{-1}(U), N) .$$

The structure of a $\mathcal{D}_X$ -module on the sheaf $j_{l,k,*}(N)$ is described as follows.

Let $n \in \Gamma(U, j_{l,k,*}(N))$ be a section corresponding to the section $\tilde{n} \in \Gamma(j_{l,k}^{-1}(U), N)$ via identification (4.40). Let $d \in D_U$ be a differential operator on U . Then the product $dn \in \Gamma(U, j_{l,k,*}(N))$ corresponds via (4.40) to the section $\tilde{d} \tilde{n} \in \Gamma(j_{l,k}^{-1}(U), N)$, where $\tilde{d} = d|_{U \cap X^k}$.

The functor $j_{l,k}^*$ is exact and maps the subcategory $\mathcal{M}_{X^l}^{hol} \subset \mathcal{M}_{X^l}$ to the subcategory $\mathcal{M}_{X^k}^{hol} \subset \mathcal{M}_{X^k}$. For any $0 \leq k_1 < k_2 < k_3 \leq N$, we have

$$j_{k_3,k_1}^* = j_{k_2,k_1}^* \cdot j_{k_3,k_2}^* .$$

We use the notation j_0^* for the functor $j_{N,0}^* : \mathcal{M}_X = \mathcal{M}_{X^N} \rightarrow \mathcal{M}_{X^0}$.

The functor $j_{l,k,*} : \mathcal{M}_{X^k} \rightarrow \mathcal{M}_{X^l}$ is right adjoint to the functor of inverse image $j_{l,k}^* : \mathcal{M}_{X^l} \rightarrow \mathcal{M}_{X^k}$ and for any $0 \leq k_1 < k_2 < k_3 \leq N$:

$$j_{k_3,k_1,*} = j_{k_3,k_2,*} \cdot j_{k_2,k_1,*} .$$

We use the notation $j_{0,*}$ for the functor $j_{N,0,*} : \mathcal{M}_{X^0} \rightarrow \mathcal{M}_{X^N} = \mathcal{M}_X$. It is exact since the map $j_0 : X^0 \rightarrow X$ is affine.

The following theorem is an immediate consequence of the constructions in the previous section.

Theorem 4.5. *The diagram below is commutative :*

$$\begin{array}{ccc} \text{Qui}_{\Gamma^l} & \xrightarrow{\mathbb{J}_{l,k}^*} & \text{Qui}_{\Gamma^k} \\ \downarrow E^l & & \downarrow E^k \\ \mathcal{M}_{X^l}^{hol} & \xrightarrow{j_{l,k}^*} & \mathcal{M}_{X^k}^{hol} . \end{array}$$

□

Let $0 \leq k < l \leq N$. For any Γ^k -quiver $\mathcal{V}_{\Gamma^k}$ and Γ^l -quiver $\mathcal{U}_{\Gamma^l}$ denote by $\alpha_{\mathcal{U}_{\Gamma^l}, \mathcal{V}_{\Gamma^k}} : \text{Hom}_{\text{Qui}_{\Gamma^l}}(\mathcal{U}_{\Gamma^l}, \mathbb{J}_{l,k,*}(\mathcal{V}_{\Gamma^k})) \rightarrow \text{Hom}_{\text{Qui}_{\Gamma^k}}(\mathbb{J}_{l,k}^*(\mathcal{U}_{\Gamma^l}), \mathcal{V}_{\Gamma^k})$ the isomorphism of adjunction of the functors $\mathbb{J}_{l,k}^*$ and $\mathbb{J}_{l,k,*}$. For any $\mathcal{D}_{X^k}$ -module M_k and any $\mathcal{D}_{X^l}$ -module M_l denote by $\tilde{\alpha}_{M_l, M_k} : \text{Hom}_{\mathcal{M}_{X^l}}(M_l, j_{l,k,*}(M_k)) \rightarrow \text{Hom}_{\mathcal{M}_{X^k}}(j_{l,k}^*(M_l), M_k)$ the isomorphism of adjunction of the functors $j_{l,k}^*$ and $j_{l,k,*}$.

Theorem 4.6. *Let $\mathcal{V}_{\Gamma^k}$ be a Γ^k quiver such that its quiver direct image $\mathbb{J}_{l,k,*}(\mathcal{V}_{\Gamma^k})$ is strongly non-resonant.*

- (i) *Then the direct image of the $\mathcal{D}_{X^k}$ -module $E^k(\mathcal{V}_{\Gamma^k})$ with respect to $j_{l,k}$ is isomorphic to the quiver $\mathcal{D}_{X^l}$ -module, associated with $\mathbb{J}_{l,k,*}(\mathcal{V}_{\Gamma^k})$:*

$$(4.41) \quad j_{l,k,*}(E^k(\mathcal{V}_{\Gamma^k})) \simeq E^l(\mathbb{J}_{l,k,*}(\mathcal{V}_{\Gamma^k})) .$$

- (ii) *Identify $\mathcal{D}_{X^l}$ -modules $j_{l,k,*}(E^k(\mathcal{V}_{\Gamma^k}))$ and $E^l(\mathbb{J}_{l,k,*}(\mathcal{V}_{\Gamma^k}))$ by isomorphism (4.41). Then for any Γ^l -quiver $\mathcal{U}_{\Gamma^l}$ and any morphism $\phi : \mathcal{U}_{\Gamma^l} \rightarrow \mathbb{J}_{l,k,*}(\mathcal{V}_{\Gamma^k})$ of Γ^l -quivers we have $E^k \alpha_{\mathcal{U}_{\Gamma^l}, \mathcal{V}_{\Gamma^k}} = \tilde{\alpha}_{E^l(\mathcal{U}_{\Gamma^l}), E^k(\mathcal{V}_{\Gamma^k})}(E^l \phi)$.*

The theorem is proved in Section 7.

Corollary 4.7. *Let $\mathcal{V}_{\Gamma^0} = \{V, A^i\}$ be a level zero quiver. Suppose that each operator A^i has a single eigenvalue. Suppose that $\mathcal{V}_{\Gamma^0}$ has a non-resonant spectrum, see Section 3.7. Then*

$$j_{0,*}(E^0(\mathcal{V}_{\Gamma^0})) \simeq E(\mathbb{J}_{0,*}(\mathcal{V}_{\Gamma^0})) .$$

4.6. Specialization of a quiver $\mathcal{D}_X$ -module to a stratum. There is a *specialization* construction in the theory of $\mathcal{D}$ -modules due to Kashiwara [K3]. This construction gives a collection of functors between quiver $\mathcal{D}$ -modules.

Let Y be a smooth complex algebraic variety and $Z \subset Y$ a smooth complex algebraic subvariety. Let $T_Z Y \rightarrow Z$ be the normal bundle of the subvariety Z . Let $I \subset \mathcal{O}_Y$ be the sheaf of ideals of functions vanishing on Z .

Define a decreasing filtration $F(\mathcal{D}_Y)$ of the sheaf $\mathcal{D}_Y$ by the formula

$$(4.42) \quad F^k(\mathcal{D}_Y) = \{P \in \mathcal{D}_Y \mid P(I_\beta^j) \subset I_\beta^{j+k} \text{ for any } j\} .$$

The associated graded quotient is isomorphic to the sheaf of differential operators $\mathcal{D}_{T_Z Y}$ on the total space $T_Z Y$ of the normal bundle to Z .

Let M be a $\mathcal{D}_Y$ -module. A decreasing filtration $F(M)$ of M is called *good* with respect to $F(\mathcal{D}_Y)$, if

- (i) $F^k(\mathcal{D}_Y)F^j(M) \subset F^{k+j}(M)$ for any k and j ,
- (ii) $F^k(\mathcal{D}_Y)F^j(M) = F^{k+j}(M)$ if $j \gg 0$ and $k \geq 0$ or $j \ll 0$ and $k \leq 0$,
- (iii) $F^j(M)$ is a coherent $F^0(\mathcal{D}_Y)$ -module,
- (iv) $M = \cup_j F^j(M)$.

Let θ be a vector field on Y tangent to Z and acting on I/I^2 as the identity. Let $\overline{G} \subset \mathbb{C}$ be a *non-resonant section* of $\mathbb{C}$, that is $\overline{G}$ contains zero and for any $a \in \mathbb{C}$ the intersection $\overline{G} \cap (a + \mathbb{Z})$ consists of a single point.

Kashiwara's theorem [K3] states that for any holonomic $\mathcal{D}_Y$ -module M with regular singularities there exists a unique filtration of M which is good with respect to $F(\mathcal{D}_Y)$ and such that

(v) there exists a polynomial $b(x)$ with zeros in $\overline{G}$ such that

$$b(\theta - k) F^k(M) \subset F^{k+1}(M)$$

for any k .

The associated graded quotient $\text{gr } M$ has a structure of a $\mathcal{D}_{T_Z Y}$ -module and is called *the specialization* of M to Z . We denote the specialization by $Sp_Z(M)$.

Thus, having $Y, Z, \theta, M, \overline{G}$, Kashiwara's theorem gives the specialization $\mathcal{D}_{T_Z Y}$ -module $Sp_Z(M)$.

In this section, we consider the case in which

- Y is a vector space $X = \mathbb{C}^N$ with a central arrangement $\mathcal{C}$,
- Z is the closure $\overline{X}_\alpha$ of a stratum of that arrangement, so Z is a vector subspace,
- $\theta = z_1 \frac{\partial}{\partial z_1} + \dots + z_k \frac{\partial}{\partial z_k}$, where $z_1, \dots, z_N$ are linear coordinates on X , such that Z is defined by equations $z_1 = \dots = z_k = 0$,
- M is a quiver $\mathcal{D}_Y$ -module associated with that arrangement.

Then it turns out that the specialization is also a quiver $\mathcal{D}_{T_Z Y}$ -module for a suitable arrangement of hyperplanes in $T_Z Y$ and a suitable quiver associated with that arrangement in $T_Z Y$.

More precisely, let $\mathcal{C}$ be a central arrangement of hyperplanes in a vector space $X = \mathbb{C}^N$. Let Γ be the graph of the arrangement $\mathcal{C}$. Fix $\alpha \in I(\Gamma)$. The arrangement $\mathcal{C}$ induces an arrangement $\mathcal{C}_\alpha$ in the total space $T_{\overline{X}_\alpha} X$ of the normal bundle $T_{\overline{X}_\alpha} X \rightarrow \overline{X}_\alpha$.

It is defined as follows. Identify $T_{\overline{X}_\alpha} X$ with the direct sum $\overline{X}_\alpha \oplus X/\overline{X}_\alpha$ and let $\pi_\alpha : X \rightarrow X/\overline{X}_\alpha$ be the natural projection. Assign to a hyperplane H of $\mathcal{C}$ the hyperplane

$$\tilde{H} = H \cap \overline{X}_\alpha \oplus \pi_\alpha(H) \subset \overline{X}_\alpha \oplus X/\overline{X}_\alpha.$$

Then the arrangement $\mathcal{C}_\alpha$ consists of all distinct hyperplanes among the hyperplanes $\tilde{H}_j, j \in J(\mathcal{C})$.

The graph Γ_α of the arrangement $\mathcal{C}_\alpha$ can be described as follows. Define an equivalence relation $\equiv_\alpha$ on points of the set $I(\Gamma)$: we say that $\beta \equiv_\alpha \gamma$ if

- (i) $\overline{X}_\alpha \cap \overline{X}_\beta = \overline{X}_\alpha \cap \overline{X}_\gamma$,
- (ii) the set of all $\delta \in I(\Gamma)$, such that $\delta \geq \alpha$ and $\delta \geq \beta$ coincides with the set of all $\delta \in I(\Gamma)$, such that $\delta \geq \alpha$ and $\delta \geq \gamma$.

Then the vertices of Γ_α are the equivalence classes $\bar{\beta}$ relative to the relation $\equiv_\alpha$.

Two vertices $\bar{\beta}, \bar{\gamma} \in I(\Gamma_\alpha)$ are connected by an arrow in Γ_α if there exist representatives $\beta \in \bar{\beta}$ and $\gamma \in \bar{\gamma}$, connected by an arrow in Γ .

The length function $l : I(\Gamma_\alpha) \rightarrow \mathbb{Z}_{\geq 0}$ is given by the rule: the length $l(\bar{\beta})$ of any equivalence class $\bar{\beta} \in I(\Gamma_\alpha)$ is the length of any representative $\beta' \in \bar{\beta} \subset I(\Gamma)$.

Let $\mathcal{V} = \{V_\beta, A_{\beta,\gamma}\}$ be a quiver of Γ . We define the new quiver $Sp_\alpha(\mathcal{V}) = \{W_{\bar{\beta}}, C_{\bar{\beta},\bar{\gamma}}\}$ of the graph Γ_α which will be called *the specialization* of the quiver $\mathcal{V}$ at $\alpha \in I(\Gamma)$. Namely, we set

$$W_{\bar{\beta}} = \oplus_{\beta'} V_{\beta'} ,$$

where the sum is over all β' from the equivalence class $\bar{\beta}$. We set

$$C_{\bar{\beta},\bar{\gamma}} = \sum_{\beta',\gamma'} A_{\beta',\gamma'} ,$$

where the sum is over all β' from the equivalence class $\bar{\beta}$ and γ' from the equivalence class $\bar{\gamma}$. It is easy to check that $Sp_\alpha(\mathcal{V})$ is a quiver of Γ_α .

Let $F(\mathcal{D}_X)$ be the decreasing filtration (4.42) of the sheaf $\mathcal{D}_X$, related to the ideal of functions vanishing on $\overline{X}_\alpha$. Define the decreasing filtration $F(E\mathcal{V})$ of the sheaf $E\mathcal{V}$ as follows. For any $j = 0, 1, \dots, l = l(\alpha)$, denote by $\overline{V}_j$ the vector space

$$\overline{V}_j = \oplus_\gamma \overline{\Omega}_\gamma \otimes V_\gamma ,$$

where the sum is taken over all γ with $\text{codim}_{\overline{X}_\gamma} \overline{X}_\gamma \cap \overline{X}_\alpha = j$. We set

$$(4.43) \quad F^k(E\mathcal{V}) = \begin{cases} \sum_{j=0}^l F^{k+l-j}(\mathcal{D}_X) \overline{V}_j , & k \leq -l , \\ \sum_{j=k+l}^l F^{k+l-j}(\mathcal{D}_X) \overline{V}_j , & -l \leq k \leq 0 , \\ F^k(\mathcal{D}_X) \overline{V}_l , & 0 \leq k . \end{cases}$$

Fix a nondegenerate bilinear form $(\cdot, \cdot)$ on $X = \mathbb{C}^N$. For any vertex $\gamma \in I(\Gamma)$, the bilinear form determines an isomorphism $\nu_\gamma : \overline{X}_\gamma \rightarrow \overline{X}_{\bar{\gamma}}$ of vector spaces where $\overline{X}_\gamma \subset X$ and $\overline{X}_{\bar{\gamma}} \subset T_{\overline{X}_\alpha} X \equiv \overline{X}_\alpha \oplus X/\overline{X}_\alpha$. Namely, let $\overline{X}'_\gamma$ be the orthogonal complement to $\overline{X}_\gamma \cap \overline{X}_\alpha$ in $\overline{X}_\gamma$. We set $\nu_\gamma(x + x') = x + \pi_\alpha(x')$ for any $x \in \overline{X}_\gamma \cap \overline{X}_\alpha$ and $x' \in \overline{X}'_\gamma$, where $\pi_\alpha : X \rightarrow X/\overline{X}_\alpha$ is the natural projection. Let $\overline{\nu}_\gamma = (\nu_\gamma^*)^{-1} : \overline{\Omega}_\gamma \rightarrow \overline{\Omega}_{\bar{\gamma}}$ be the associated isomorphism of the spaces of top degree holomorphic differential forms invariant with respect to translations.

For any vertex $\beta \in I(\Gamma)$ define the linear operator $S_\beta^{(\alpha)} : V_\beta \rightarrow V_\beta$ by the formula

$$S_\beta^{(\alpha)} = \sum_\gamma A_{\beta,\gamma} A_{\gamma,\beta} ,$$

where the sum is over all $\gamma \in I(\Gamma)$ such that $\overline{X}_\alpha \cap \overline{X}_\gamma = \overline{X}_\beta \cap \overline{X}_\gamma$. We define the operator

$$S^{(\alpha)} : \oplus_{\beta \in I(\Gamma)} V_\beta \rightarrow \oplus_{\beta \in I(\Gamma)} V_\beta$$

by the formula

$$S^{(\alpha)} = \sum_{\beta \in I(\Gamma)} S_\beta^{(\alpha)} .$$

Lemma 4.8.

- (i) *The filtration $F(E\mathcal{V})$ is good with respect to $F(\mathcal{D}_X)$.*
- (ii) *The assignment $\text{gr}(\overline{\omega}_\gamma \otimes v_\gamma) \mapsto \overline{\nu}_\gamma(\overline{\omega}_\gamma) \otimes v_\gamma$, with $\gamma \in I(\Gamma)$, $v_\gamma \in V_\gamma$, $\overline{\omega}_\gamma \in \overline{\Omega}_\gamma$, establishes an isomorphism of the $\mathcal{D}_{T_{\overline{X}_\alpha} X}$ -modules $\text{gr } E\mathcal{V}$ and $ESp_\alpha(\mathcal{V})$.*

(iii) For $\beta \in I(\Gamma)$ let $j = \text{codim } \overline{X}_\beta \cap \overline{X}_\alpha$. Then for any $\overline{\omega}_\beta \otimes v_\beta \in \overline{\Omega}_\beta \otimes V_\beta$ we have

$$(\theta - j + l)(\overline{\omega}_\beta \otimes v_\beta) - 1 \otimes S_\beta^{(\alpha)}(\overline{\omega}_\beta \otimes v_\beta) \in F^{j-l+1}(E\mathcal{V}).$$

□

The proof is by direct calculation like in the proof of Proposition 4.3.

Lemma 4.9. *Let $G^{(\alpha)}$ be a non-resonant set and $\overline{G}^{(\alpha)}$ a non-resonant section such that $\overline{G}^{(\alpha)} \supset G^{(\alpha)}$. Assume that all eigenvalues of $S^{(\alpha)}$ belong to $G^{(\alpha)}$. Then the $\mathcal{D}_{T_{\overline{X}_\alpha} X}$ -module $\text{gr } E\mathcal{V}$ from Lemma 4.8 is the specialization module $Sp_{\overline{X}_\alpha}(E\mathcal{V})$ relative to the non-resonant section $\overline{G}^{(\alpha)}$ and vector field θ defined above.*

Proof. Due to statements (i) and (ii) of Lemma 4.8, it is sufficient to verify condition (v) for the filtration $F(E\mathcal{V})$. But condition (v) follows from the definition of the filtration in (4.43), statement (iii) of Lemma 4.8, and Proposition 4.2. □

Corollary 4.10. *Under the above assumptions the $\mathcal{D}_{T_{\overline{X}_\alpha} X}$ -modules $ESp_\alpha(\mathcal{V})$ and $Sp_{\overline{X}_\alpha}(E\mathcal{V})$ are isomorphic with respect to the isomorphism from Lemma 4.8.*

4.7. Free resolution of quiver $\mathcal{D}_{X^0}$ -modules. Let $\mathcal{V}_{\Gamma^0} = \{V, A^j \mid j \in J(\mathcal{C})\}$ be a quiver in Qui_{Γ^0} and $E^0\mathcal{V}_{\Gamma^0}$ the corresponding $\mathcal{D}_{X^0}$ -module.

Set

$$\mathcal{R}^r E^0\mathcal{V}_{\Gamma^0} = \mathcal{D}_{X^0} \otimes \bigwedge^{-r} T_\emptyset \otimes \overline{\Omega}_\emptyset \otimes V, \quad \mathcal{R}E^0\mathcal{V}_{\Gamma^0} = \bigoplus_{r \in \mathbb{Z}} \mathcal{R}^r E^0\mathcal{V}_{\Gamma^0}.$$

These are free left $\mathcal{D}_{X^0}$ -modules.

Set $\deg \mathcal{R}^r E^0\mathcal{V}_{\Gamma^0} = r$. Define $\mathcal{D}_{X^0}$ -linear maps $d_i : \mathcal{R}E^0\mathcal{V}_{\Gamma^0} \rightarrow \mathcal{R}E^0\mathcal{V}_{\Gamma^0}$, $i = 0, 1$, of degree 1 and a $\mathcal{D}_{X^0}$ -linear map $\nu : \mathcal{R}^0 E^0\mathcal{V}_{\Gamma^0} \rightarrow E^0\mathcal{V}_{\Gamma^0}$ by the formulas:

$$\begin{aligned} d_1(1 \otimes \xi_1 \wedge \dots \wedge \xi_r \otimes \overline{\omega}_\emptyset \otimes v) &= \sum_{i=1}^r (-1)^{i+1} \xi_i \otimes \xi_1 \wedge \dots \widehat{\xi_i} \dots \wedge \xi_r \otimes \overline{\omega}_\emptyset \otimes v, \\ d_0(1 \otimes \xi_1 \wedge \dots \wedge \xi_r \otimes \overline{\omega}_\emptyset \otimes v) &= \sum_{j \in J(\mathcal{C})} f_j^{-1} \otimes i_{f_j}(\xi_1 \wedge \dots \wedge \xi_r) \otimes \overline{\omega}_\emptyset \otimes A^j(v), \\ \nu(1 \otimes \overline{\omega}_\emptyset \otimes v) &= \overline{\omega}_\emptyset \otimes v. \end{aligned}$$

Here $f_j = 0$ is an equation of the hyperplane H_j . Set also $d = d_1 - d_0$. We have

$$d_i^2 = 0, \quad \text{for } i = 0, 1, \quad d_0 d_1 + d_1 d_0 = 0.$$

Theorem 4.11. *The complex*

$$0 \rightarrow \mathcal{R}^{-N} E^0\mathcal{V}_{\Gamma^0} \xrightarrow{d} \mathcal{R}^{-N+1} E^0\mathcal{V}_{\Gamma^0} \xrightarrow{d} \dots \xrightarrow{d} \mathcal{R}^0 E^0\mathcal{V}_{\Gamma^0} \xrightarrow{\nu} E^0\mathcal{V}_{\Gamma^0} \rightarrow 0$$

is acyclic and presents a free $\mathcal{D}_{X^0}$ -module resolution of the $\mathcal{D}_{X^0}$ -module $E^0\mathcal{V}_{\Gamma^0}$.

The theorem is proved in Section 7.

4.8. Free resolution of quiver $\mathcal{D}_X$ -modules. In this section we generalize the result of the previous section and attach to a quiver $\mathcal{V} \in \text{Qui}_\Gamma$ an acyclic complex

$$\overline{\mathcal{R}}E\mathcal{V} : \quad 0 \rightarrow \mathcal{R}^{-N}E\mathcal{V} \rightarrow \mathcal{R}^{-N+1}E\mathcal{V} \rightarrow \dots \rightarrow \mathcal{R}^0E\mathcal{V} \rightarrow E\mathcal{V} \rightarrow 0$$

of left $\mathcal{D}_X$ -modules, where all $\mathcal{R}^iE\mathcal{V}$ are finitely generated free left $\mathcal{D}_X$ -modules. This complex gives a free resolution of the quiver $\mathcal{D}_X$ -module $E\mathcal{V}$.

In the construction of the resolution we will use the following objects introduced in Section 2.2.

For $\alpha \in I(\Gamma)$, we will use the associated spaces $\mathcal{T}_\alpha = T_\alpha \oplus F_\alpha$, and $\overline{\mathcal{Q}}_\alpha$. For strata X_α and X_β , $\overline{X}_\alpha \supset X_\beta$, we will use the inclusions

$$\mu_{\beta,\alpha} : \bigwedge^p F_\alpha \hookrightarrow \bigwedge^p F_\beta, \quad \overline{\mu}_{\alpha,\beta} : \bigwedge^p T_\beta \hookrightarrow \bigwedge^p T_\alpha.$$

For $\alpha \succ \beta$, $f \in F_\beta$, $\xi \in T_\alpha$, we will use the linear maps

$$i_f^{\beta,\alpha} : \bigwedge^p T_\alpha \rightarrow \bigwedge^{p-1} T_\beta, \quad i_\xi^{\alpha,\beta} : \bigwedge^p F_\beta \rightarrow \bigwedge^{p-1} F_\alpha.$$

For $r = 0, \dots, -N$, set

$$(4.44) \quad \mathcal{R}^r E\mathcal{V} = \bigoplus_{\alpha \in I(\Gamma)} \mathcal{D}_X \otimes \bigwedge^{-r} \mathcal{T}_\alpha \otimes \overline{\mathcal{Q}}_\alpha \otimes V_\alpha.$$

This is a free left $\mathcal{D}_X$ -module. Set $\mathcal{R}^r E\mathcal{V} = 0$ for $r > 0$ and $r < -N$. Set

$$\mathcal{R}E\mathcal{V} = \bigoplus_{r \in \mathbb{Z}} \mathcal{R}^r E\mathcal{V}.$$

For an edge $(\alpha, \beta) \in E(\Gamma)$, and r , $1 \leq r \leq N$, we define a linear map

$$A_{\beta,\alpha}^r : \bigwedge^r \mathcal{T}_\alpha \otimes \overline{\mathcal{Q}}_\alpha \otimes V_\alpha \rightarrow \bigwedge^{r-1} \mathcal{T}_\beta \otimes \overline{\mathcal{Q}}_\beta \otimes V_\beta.$$

The construction of the map uses an edge framing $\{f_{\alpha,\beta}, \xi_{\alpha,\beta}\}$, but the map does not depend on the choice of the framing.

The map

$$A_{\beta,\alpha}^1 : \bigwedge^1 \mathcal{T}_\alpha \otimes \overline{\mathcal{Q}}_\alpha \otimes V_\alpha \rightarrow \bigwedge^0 \mathcal{T}_\beta \otimes \overline{\mathcal{Q}}_\beta \otimes V_\beta$$

is defined as follows. For $\xi \in T_\alpha$, $f \in F_\alpha$, $\overline{w}_\alpha \otimes v_\alpha \in \overline{\mathcal{Q}}_\alpha \otimes V_\alpha$ we set

$$A_{\beta,\alpha}^1(\xi \otimes \overline{w}_\alpha \otimes v_\alpha) = \langle \xi, f_{\beta,\alpha} \rangle \pi_{\beta,\alpha}(\overline{w}_\alpha) \otimes A_{\beta,\alpha}(v_\alpha), \quad A_{\beta,\alpha}^1(f \otimes \overline{w}_\alpha \otimes v_\alpha) = 0$$

if $\alpha \succ \beta$ and

$$A_{\beta,\alpha}^1(f \otimes \overline{w}_\alpha \otimes v_\alpha) = \langle \xi_{\beta,\alpha}, f \rangle \pi_{\beta,\alpha}(\overline{w}_\alpha) \otimes A_{\beta,\alpha}(v_\alpha), \quad A_{\beta,\alpha}^1(\xi \otimes \overline{w}_\alpha \otimes v_\alpha) = 0$$

if $\beta \succ \alpha$.

Define the map $A_{\beta,\alpha}^r$ for an arbitrary r . Let $\vec{\xi} \wedge \vec{f} \in \bigwedge^p T_\alpha \wedge \bigwedge^{r-p} F_\alpha \subset \bigwedge^r \mathcal{T}_\alpha$ and $\overline{w}_\alpha \otimes v_\alpha \in \overline{\mathcal{Q}}_\alpha \otimes V_\alpha$. If $\alpha \succ \beta$, set

$$(4.45) \quad A_{\beta,\alpha}^r(\vec{\xi} \wedge \vec{f} \otimes \overline{w}_\alpha \otimes v_\alpha) = i_{f_{\beta,\alpha}}^{\beta,\alpha}(\vec{\xi}) \wedge \vec{f} \otimes \pi_{\beta,\alpha}(\overline{w}_\alpha) \otimes A_{\beta,\alpha}(v_\alpha),$$

where the polyvector $\vec{f}$ in the right hand side of (4.45) is considered as an element of $\bigwedge^{r-p} F_\beta$ by means of the tautological embedding $\mu_{\alpha,\beta}$. If $\beta \succ \alpha$, set

$$(4.46) \quad A_{\beta,\alpha}^r(\vec{\xi} \wedge \vec{f} \otimes \bar{\omega}_\alpha \otimes v_\alpha) = (-1)^p \vec{\xi} \wedge i_{\xi_{\beta,\alpha}}^{\beta,\alpha}(\vec{f}) \otimes \pi_{\beta,\alpha}(\bar{\omega}_\alpha) \otimes A_{\beta,\alpha}(v_\alpha),$$

where the polivector $\vec{\xi}$ in the right hand side of (4.46) is considered as an element of $\bigwedge^p T_\beta$ by means of the tautological embedding $\bar{\mu}_{\alpha,\beta}$.

For $i = 0, 1$, define $\mathcal{D}_X$ -linear maps $d_i : \mathcal{R}E\mathcal{V} \rightarrow \mathcal{R}E\mathcal{V}$ by the formulas:

$$(4.47) \quad \begin{aligned} d_1(1 \otimes t_1 \wedge \dots \wedge t_r \otimes \bar{\omega}_\alpha \otimes v_\alpha) &= \sum_{i=1}^r (-1)^{i+1} t_i \otimes t_1 \wedge \dots \widehat{t_i} \dots \wedge t_r \otimes \bar{\omega}_\alpha \otimes v_\alpha, \\ d_0(1 \otimes t_1 \wedge \dots \wedge t_r \otimes \bar{\omega}_\alpha \otimes v_\alpha) &= \sum_{\beta, (\alpha,\beta) \in E(\Gamma)} 1 \otimes A_{\beta,\alpha}^r(t_1 \wedge \dots \wedge t_r \otimes \bar{\omega}_\alpha \otimes v_\alpha), \end{aligned}$$

where $t_i \in \mathcal{T}_\alpha$, $\bar{\omega}_\alpha \otimes v_\alpha \in \bar{\Omega}_\alpha \otimes V_\alpha$. Set

$$d_i(1 \otimes \bar{\omega}_\alpha \otimes v_\alpha) = 0.$$

It is clear that $d_i(\mathcal{R}^r E\mathcal{V}) \subset \mathcal{R}^{r+1} E\mathcal{V}$ for $i = 0, 1$.

Define a morphism $\nu : \mathcal{R}^0 E\mathcal{V} \rightarrow E\mathcal{V}$ of $\mathcal{D}_X$ -modules by the formula

$$\nu(1 \otimes \bar{\omega}_\alpha \otimes v_\alpha) = \bar{\omega}_\alpha \otimes v_\alpha.$$

Proposition 4.12.

(i) *The maps d_i are commuting differentials,*

$$d_i^2 = 0, \quad \text{for } i = 0, 1, \quad d_0 d_1 + d_1 d_0 = 0;$$

(ii) *Set $d = d_1 - d_0$. Then*

$$\nu d = 0.$$

□

Denote by $E\mathcal{V}^{(\cdot)}$ the $\mathcal{D}_X$ -module $E\mathcal{V}$ considered as a complex with the zero differential, concentrated in degree zero. Then, by Proposition 4.12, we can introduce two complexes,

$$\begin{aligned} \mathcal{R}E\mathcal{V} : \quad 0 \rightarrow \mathcal{R}^{-N} E\mathcal{V} \xrightarrow{d} \mathcal{R}^{-N+1} E\mathcal{V} \xrightarrow{d} \dots \xrightarrow{d} \mathcal{R}^0 E\mathcal{V} \rightarrow 0, \\ \overline{\mathcal{R}}E\mathcal{V} : \quad 0 \rightarrow \mathcal{R}^{-N} E\mathcal{V} \xrightarrow{d} \mathcal{R}^{-N+1} E\mathcal{V} \xrightarrow{d} \dots \xrightarrow{d} \mathcal{R}^0 E\mathcal{V} \xrightarrow{\nu} E\mathcal{V} \rightarrow 0 \end{aligned}$$

with $\deg \mathcal{R}^k E\mathcal{V} = k$. The second complex can be considered as a morphism of complexes $\nu : \mathcal{R}E\mathcal{V} \rightarrow E\mathcal{V}^{(\cdot)}$.

The following statement is one of the main results of the paper.

Theorem 4.13. *For any quiver $\mathcal{V} \in \text{Qui}_\Gamma$, the complex $\overline{\mathcal{R}}E\mathcal{V}$ is acyclic.*

The theorem is proved in Section 7.

By Theorem 4.13, the complex $\overline{\mathcal{R}}E\mathcal{V}$ gives a free $\mathcal{D}_X$ -module resolution of the $\mathcal{D}_X$ -module $E\mathcal{V}$. Equivalently, the complex $\mathcal{R}E\mathcal{V}$ of free $\mathcal{D}_X$ modules is quasi-isomorphic to $E\mathcal{V}^{(\cdot)}$, that is, the morphism $\nu : \mathcal{R}E\mathcal{V} \rightarrow E\mathcal{V}^{(\cdot)}$ induces an isomorphism of cohomology groups of complexes.

Example. Let $X = \mathbb{C}^1$, $\mathcal{C} = \{z = 0\}$. The graph Γ of the arrangement has two vertices, $\emptyset$ and α , $l(\alpha) = 1$, connected by an edge. Let $\mathcal{V} = \{V_\emptyset, V_\alpha, A_{\emptyset, \alpha}, A_{\alpha, \emptyset}\}$ be a quiver of Γ . The free $\mathcal{D}_X$ -module resolution $\mathcal{R}E\mathcal{V}$ of the quiver $\mathcal{D}_X$ -module $E\mathcal{V}$ is the complex

$$\mathcal{R}E\mathcal{V} : 0 \rightarrow \mathcal{R}^{-1}E\mathcal{V} \xrightarrow{d} \mathcal{R}^0E\mathcal{V} \xrightarrow{\nu} E\mathcal{V} \rightarrow 0.$$

Here $\mathcal{R}^0E\mathcal{V}$ is the free $\mathcal{D}_X$ -module $\mathcal{D}_X \otimes \overline{\Omega}_\emptyset \otimes V_\emptyset \oplus \mathcal{D}_X \otimes \overline{\Omega}_\alpha \otimes V_\alpha$. The term $\mathcal{R}^{-1}E\mathcal{V}$ is the free $\mathcal{D}_X$ -module $\mathcal{D}_X \otimes T_\emptyset \otimes \overline{\Omega}_\emptyset \otimes V_\emptyset \oplus \mathcal{D}_X \otimes F_\alpha \otimes \overline{\Omega}_\alpha \otimes V_\alpha$, where $T_\emptyset$ is the one-dimensional space $\mathbb{C} \cdot \frac{d}{dz}$ of vector fields with constant coefficients, parallel to $X_\emptyset$, F_α is the one-dimensional space $\mathbb{C} \cdot z$ of affine functions, vanishing at X_α . The differentials d, ν are given by the relations:

$$\begin{aligned} d(D \otimes, \frac{d}{dz} \otimes dz \otimes v_\emptyset) &= D \cdot \frac{d}{dz} \otimes dz \otimes v_\emptyset - D \otimes 1 \otimes A_{\alpha, \emptyset}(v_\emptyset), \\ d(D \otimes z \otimes 1 \otimes v_\alpha) &= D \cdot z \otimes 1 \otimes v_\alpha - D \otimes dz \otimes A_{\emptyset, \alpha}(v_\alpha), \\ \nu(D \otimes 1 \otimes v_\alpha) &= D \cdot (1 \otimes v_\alpha), \quad \nu(D \otimes dz \otimes v_\emptyset) = D \cdot (dz \otimes v_\emptyset), \end{aligned}$$

for any $D \in \mathcal{D}_X$, $v_\emptyset \in V_\emptyset$, $v_\alpha \in V_\alpha$. Here $D \cdot (1 \otimes v_\alpha)$ and $D \cdot (dz \otimes v_\emptyset)$ denote the application of D to elements of $E\mathcal{V}$.

4.9. Derived categories and derived functors. In the following we use the language of derived categories. Let us briefly recall necessary definitions and constructions, see e.g. [GM].

Let $\mathcal{A}$ be an abelian category. Denote by $\mathcal{K}om(\mathcal{A})$ (resp. $\mathcal{K}om^b(\mathcal{A})$) the category of complexes (resp. bounded complexes) over $\mathcal{A}$ with differential of degree one. Morphisms in both categories are morphisms of complexes, commuting with differentials: $f d_A = d_B f$.

For any complex $A^\bullet \in \mathcal{K}om(\mathcal{A})$ and $n \in \mathbb{Z}$ denote by $A[n]^\bullet$ the same complex with the shifted gradation, $A[n]^k = A^{n+k}$. For any object $A \in \mathcal{A}$ denote by $A^{(\cdot)}$ the complex with zero differential, defined by the condition $(A^{(\cdot)})^0 = A$ and $(A^{(\cdot)})^k = 0$ for $k \neq 0$.

A morphism of complexes $f : A^\bullet \rightarrow B^\bullet$ is called a quasi-isomorphism, if it induces an isomorphism of cohomology groups $H(A^\bullet)$ and $H(B^\bullet)$. It is denoted by $f : A^\bullet \xrightarrow{quis} B^\bullet$. Morphisms $f, f' : A^\bullet \rightarrow B^\bullet$ are called homotopic, if there exists a map of complexes $h : A^\bullet \rightarrow B^\bullet$ of degree -1 , (that is, a morphism $h : A^\bullet \rightarrow B[1]^\bullet$), such that $f - f' = h d_A + d_B h$.

Denote by $\mathcal{K}(\mathcal{A})$ (resp. $\mathcal{K}^b(\mathcal{A})$) the homotopy (respectively, bounded homotopy) category of complexes over $\mathcal{A}$. Objects of $\mathcal{K}(\mathcal{A})$ (resp. $\mathcal{K}^b(\mathcal{A})$) are objects of $\mathcal{K}om(\mathcal{A})$, (resp. $\mathcal{K}om^b(\mathcal{A})$). Morphisms are classes of morphisms of complexes modulo homotopy equivalence.

The derived category $D(\mathcal{A})$ is defined as the localization of the category $\mathcal{K}(\mathcal{A})$ over the set of all classes of quasi-isomorphisms. The bounded derived category $D^b(\mathcal{A})$ is defined as the localization of the category $\mathcal{K}^b(\mathcal{A})$ over the set of all classes of quasi-isomorphisms.

Objects of $D(\mathcal{A})$ are complexes over $\mathcal{A}$, and every morphism $f : A^\bullet \rightarrow B^\bullet$ can be presented as a triple $\{C^\bullet \in \mathcal{K}(\mathcal{A}), f' \in \text{Hom}_{\mathcal{K}(\mathcal{A})}(A^\bullet, C^\bullet), f'' \in \text{Hom}_{\mathcal{K}(\mathcal{A})}(B^\bullet, C^\bullet)\}$, where f' is a class of quasi-isomorphisms. An analogous description holds for the bounded derived category $D^b(\mathcal{A})$.

Let $\mathcal{A}$ and $\mathcal{B}$ be abelian categories and $F : \mathcal{A} \rightarrow \mathcal{B}$ a left (resp. right) exact additive functor. It admits a natural extension to a functor $F^\bullet : K(\mathcal{A}) \rightarrow K(\mathcal{B})$ of homotopy categories. Usually it is denoted by the same symbol F .

A class $\mathcal{A}_F$ of objects in $Ob(\mathcal{A})$ is called adjusted to F if

- (i) It is closed under direct sums;
- (ii) The functor $F^\bullet$ transforms acyclic bounded from below (resp. from above) complexes with graded components in $\mathcal{A}_F$ to acyclic complexes;
- (iii) For any object $A \in \mathcal{A}$, there exists an object $A' \in \mathcal{A}_F$ and an inclusion $\alpha : A \rightarrow A'$ (resp. an epimorphism $A' \rightarrow A$).

If a class $\mathcal{A}_F$ of objects adjusted to F is chosen, then the right derived functor $RF : D(\mathcal{A}) \rightarrow D(\mathcal{B})$ (resp. the left derived functor $LF : D(\mathcal{A}) \rightarrow D(\mathcal{B})$) is defined as follows. By (iii), for any complex $A^\bullet \in D(\mathcal{A})$, there exists a complex $A'^\bullet$ over $\mathcal{A}_F$ and a quasi-isomorphism $\phi : A^\bullet \rightarrow A'^\bullet$ (resp. a complex $A''^\bullet$ and a quasi-isomorphism $\psi : A''^\bullet \rightarrow A^\bullet$). By definition, $RF(A^\bullet) = F^\bullet(A'^\bullet)$ (resp. $LF(A^\bullet) = F^\bullet(A''^\bullet)$).

It can be proved that the derived functor does not depend on the choice of the class of adjusted objects.

We keep the natural notation $RF(A) = RF(A^{(\cdot)})$ and $LF(A) = LF(A^{(\cdot)})$ for any object $A \in \mathcal{A}$.

In most cases left and right derived functors admit well defined restrictions to bounded derived categories. It takes place, in particular, if any object A in $\mathcal{A}$ admits a finite resolution $\phi : A^{(\cdot)} \xrightarrow{quis} A'^\bullet$ over $\mathcal{A}_F$ for the case of a left exact functor F (resp. admits a finite resolution $\phi : A'^\bullet \xrightarrow{quis} A^{(\cdot)}$ over $\mathcal{A}_F$ for the case of a right exact functor F). It is always so in our considerations.

For a smooth algebraic variety Y we denote by $D^b(\mathcal{M}_Y)$ the bounded derived category of the category of coherent $\mathcal{D}_Y$ -modules. For a topological space Y we denote by $D^b(Y, \mathbb{C})$ the bounded derived category of the category $Sh(Y, \mathbb{C})$ of sheaves of complex vector spaces on Y .

4.10. Duality of quiver $\mathcal{D}_X$ -modules. Let Y be a smooth algebraic variety over $\mathbb{C}$ and Ω_Y the sheaf of germs of top exterior differential forms on Y . The sheaf Ω_Y is an invertible locally free sheaf of $\mathcal{O}_Y$ -modules of rank 1. Let Ω_Y^{-1} be the corresponding inverse sheaf. Denote by $\tilde{\mathcal{D}}_Y$ the sheaf $\mathcal{D}_Y \otimes_{\mathcal{O}_Y} \Omega_Y^{-1}$. It has two structures of a (locally free) left $\mathcal{D}_Y$ -module.

The first structure is given locally by the relations:

$$(4.48) \quad f \cdot (d \otimes \tilde{\omega}) = df \otimes \tilde{\omega}, \quad \xi \cdot (d \otimes \tilde{\omega}) = -d\xi \otimes \tilde{\omega} + d \otimes Lie_\xi(\tilde{\omega}),$$

where $f, d, \tilde{\omega}$ are local sections of the sheaves $\mathcal{O}_Y, \mathcal{D}_Y$, and Ω_Y^{-1} , respectively, ξ is a vector field over Y and Lie_ξ is the Lie derivative in the direction of ξ .

The second structure of a left $\mathcal{D}_Y$ module on $\tilde{\mathcal{D}}_Y$ is given by the left multiplication of $\mathcal{D}_Y$ on itself:

$$d_1 \star (d_2 \otimes \tilde{\omega}) = d_1 d_2 \otimes \tilde{\omega}.$$

Let $M^\bullet$ be a complex of $\mathcal{D}_Y$ -modules. We set

$$\Delta_Y M^\bullet = RHom_{\mathcal{D}_Y}(M^\bullet, \tilde{\mathcal{D}}_Y)[\dim Y].$$

This formula has the following meaning. The complex $\Delta_Y M^\bullet$ of left $\mathcal{D}_Y$ -modules is quasi-isomorphic to the complex $Q^\bullet$, where Q^i is the sheaf

$$Q^i = Hom_{\mathcal{D}_Y}(P^{-i-\dim Y}, \tilde{\mathcal{D}}_Y).$$

The complex $P^\bullet$ is a complex of locally free left $\mathcal{D}_Y$ -modules, quasi-isomorphic to $M^\bullet$. The local Hom functor is taken between the left $\mathcal{D}_Y$ -module $P^{-i-\dim Y}$ and the left $\mathcal{D}_Y$ -module $\widetilde{\mathcal{D}}_Y$, considered with respect to the first structure. The structure of a left $\mathcal{D}_Y$ -module on Q^i is induced from the second module structure on $\widetilde{\mathcal{D}}_Y$ as follows. Let ϕ be a local homomorphism $\phi : P^{-i-\dim Y} \rightarrow \widetilde{\mathcal{D}}_Y$, satisfying the condition $\phi(d' \cdot p) = d' \cdot \phi(p)$ for any $p \in P^{-i-\dim Y}$ and $d' \in \mathcal{D}_Y$. Then for any $d \in \mathcal{D}_Y$, the local $\mathcal{D}_Y$ -linear map $d \cdot \phi$ is defined by the rule $(d \cdot \phi)(p) = d \star \phi(p)$.

It is known that if M is a holonomic $\mathcal{D}_Y$ -module, then $H^i(\Delta_Y M^{(\cdot)}) = 0$ for $i \neq 0$. The map $M \mapsto H^0(\Delta_Y M^{(\cdot)})$ defines an exact contravariant functor $\mathcal{M}_Y^{\text{hol}} \rightarrow \mathcal{M}_Y^{\text{hol}}$ which we denote by the same symbol Δ_Y . For any $M \in \mathcal{M}_Y^{\text{hol}}$, the $\mathcal{D}_Y$ -module $\Delta_Y(\Delta_Y(M))$ is isomorphic to M .

Let $j : Z \rightarrow Y$ be an open embedding of smooth algebraic varieties. Then, besides the functors $j^* : \mathcal{M}_Y \rightarrow \mathcal{M}_Z$ and $j_* : \mathcal{M}_Z \rightarrow \mathcal{M}_Y$ of the inverse and direct images of $\mathcal{D}$ -modules, the second direct image functor $j_!^{(D)} : \mathcal{M}_Z^{\text{hol}} \rightarrow \mathcal{M}_Y^{\text{hol}}$ can be defined by the rule:

$$(4.49) \quad j_!^{(D)}(M) = \Delta_Y j_* \Delta_Z(M), \quad M \in \mathcal{M}_Z^{\text{hol}}.$$

It is the left adjoint functor to the functor $j^* : \mathcal{M}_Y^{\text{hol}} \rightarrow \mathcal{M}_Z^{\text{hol}}$. For any $M \in \mathcal{M}_Z^{\text{hol}}$, we denote by $j_{!*}^{(D)}(M)$ the image of $j_!^{(D)}(M)$ in $j_*(M)$ with respect to the natural map $s_j^{(D)} : j_!^{(D)}(M) \rightarrow j_*(M)$, induced by canonical transformations between the pairs of adjoint functors (j^*, j_*) and $(j_!^{(D)}, j^*)$, see Section 3.3.

Denote by $j_{l,k,!} : \mathcal{M}_{X^k}^{\text{hol}} \rightarrow \mathcal{M}_{X^l}^{\text{hol}}$ the direct image functor $(j_{l,k})_!^{(D)}$. It is left adjoint to the functor of inverse image $j_{l,k}^* : \mathcal{M}_{X^l}^{\text{hol}} \rightarrow \mathcal{M}_{X^k}^{\text{hol}}$. For $0 \leq k_1 < k_2 < k_3 \leq N$, we have

$$j_{k_3,k_1,!} = j_{k_3,k_2,!} \cdot j_{k_2,k_1,!}.$$

We use the notation $j_{0,!}$ for the functor $j_{N,0,!} : \mathcal{M}_{X^0}^{\text{hol}} \rightarrow \mathcal{M}_{X^N}^{\text{hol}} = \mathcal{M}_X^{\text{hol}}$. It is exact since the map X^0 is an affine variety [B]. For any $M \in \mathcal{M}_{X^0}^{\text{hol}}$, we denote by $j_{0,!}(M)$ the $\mathcal{D}_X$ -module $(j_{N,0})_!^{(D)}(M)$.

Theorem 4.14.

(i) Let $\mathcal{V}_{\Gamma^0}$ be a level zero quiver. Then the following $\mathcal{D}_{X^0}$ -modules are isomorphic:

$$\Delta_{X^0}(E^0(\mathcal{V}_{\Gamma^0})) \approx E^0(\tau(\mathcal{V}_{\Gamma^0})).$$

(ii) Let $\mathcal{V}$ be a quiver of Γ . Then the following $\mathcal{D}_X$ -modules are isomorphic:

$$\Delta_X(E(\mathcal{V})) \approx E(\tau(\mathcal{V})).$$

The theorem is proved in Section 7.

Corollary 4.15. Let $\mathcal{V}_{\Gamma^0} = \{V, A^i\}$ be a level zero quiver. Suppose that each operator A^i has a single eigenvalue. Suppose that $\mathcal{V}_{\Gamma^0}$ has a non-resonant spectrum, see Section 3.7. Then the following $\mathcal{D}_X$ -modules are isomorphic:

$$\begin{aligned} j_{0,!}(E^0(\mathcal{V}_{\Gamma^0})) &\approx E(\mathbb{J}_{0,!}(\mathcal{V}_{\Gamma^0})), \\ j_{0,!}(E^0(\mathcal{V}_{\Gamma^0})) &\approx E(\mathbb{J}_{0,!}(\mathcal{V}_{\Gamma^0})). \end{aligned}$$

Proof. The first statement follows from the definitions of the functors $j_{0,!} = (j_{N,0})_!$ and $\mathbb{J}_{0,!} = \mathbb{J}_{N,0,!}$, see (3.14), (4.49), and Theorem 4.14.

The canonical morphism $s_0 : \mathbb{J}_{0,!}(\mathcal{V}_{\Gamma^0}) \rightarrow \mathbb{J}_{0,*}(\mathcal{V}_{\Gamma^0})$, see (3.16), is determined by the collection of isomorphisms of adjunction

$$\alpha_{\mathcal{U}_{\Gamma}, \mathcal{V}_{\Gamma^0}} : \text{Hom}_{\text{Qui}_{\Gamma}}(\mathcal{U}_{\Gamma}, \mathbb{J}_{0,*}(\mathcal{V}_{\Gamma^0})) \rightarrow \text{Hom}_{\text{Qui}_{\Gamma^0}}(\mathbb{J}_0^*(\mathcal{U}_{\Gamma}), \mathcal{V}_{\Gamma^0}) ,$$

where $\mathcal{U}_{\Gamma} \in \text{Qui}_{\Gamma}$, and by the automorphisms τ_0 and τ_N . Analogously, for any $M_0 \in \mathcal{M}_{X^0}$, the canonical morphism $s_0^{(D)} : j_{0,!}(M_0) \rightarrow \mathbb{J}_{0,*}(M_0)$, is determined by the collection of isomorphisms of adjunction

$$\tilde{\alpha}_{M, M_0} : \text{Hom}_{\mathcal{M}_X}(M, j_{0,*}(M_0)) \rightarrow \text{Hom}_{\mathcal{M}_{X^0}}(j_0^*(M), M_0) ,$$

where $M \in \mathcal{M}_X$, and by the automorphisms Δ_{X^0} and Δ_X . By Theorem 4.14 and Theorem 4.6, statement (ii), we have an equality $Es_0(\mathbb{J}_{0,!}(\mathcal{V}_{\Gamma^0})) = s_0^{(D)}(E\mathbb{J}_{0,!}(\mathcal{V}_{\Gamma^0}))$, which implies the second statement of Corollary 4.15. $\square$

4.11. Fourier transform of quiver $\mathcal{D}_X$ -modules.

4.11.1. Let $z_1, \dots, z_N$ be linear coordinates on $X = \mathbb{C}^N$ and $\xi_1, \dots, \xi_N$ the dual coordinates on the dual space X^* . The assignment

$$\xi_i \mapsto -\frac{\partial}{\partial z_i} , \quad \frac{\partial}{\partial \xi_i} \mapsto z_i$$

for $i = 1, \dots, N$, defines an isomorphism of the rings D_{X^*} and D_X called *the Fourier transform*. The Fourier transform defines a functor from the category $\mathcal{M}_X$ to the category $\mathcal{M}_{X^*}$.

Let $\mathcal{C} = \{H_j\}, j \in J(\mathcal{C})$, be a central arrangement of hyperplanes in the vector space $X = \mathbb{C}^N$. Let Γ be the graph of the arrangement, $\mathcal{V}$ a quiver of Γ , and $E\mathcal{V}$ the associated quiver $\mathcal{D}_X$ -module. In this section we shall discuss how to describe combinatorially the Fourier transform of $E\mathcal{V}$. It turns out that the Fourier transform of $E\mathcal{V}$ is the quiver $\mathcal{D}_{X^*}$ -module associated with a suitable line arrangement in X^* .

First we shall describe the line arrangement. Then we shall introduce quivers associated with the line arrangement. Then we shall describe the Fourier transform of $E\mathcal{V}$.

4.11.2. Let $\mathcal{C} = \{H_j\}, j \in J(\mathcal{C})$, be a central arrangement of hyperplanes in $X = \mathbb{C}^N$. Then the one-dimensional subspaces $H_j^\perp \in X^*$, $j \in J(\mathcal{C})$, form an arrangement of lines in X^* , denoted by $\mathcal{C}^\perp$.

The arrangement $\mathcal{C}^\perp$ defines on X^* the structure of a stratified space. The strata of X^* are labeled by vertices of the graph Γ of the arrangement $\mathcal{C}$. For $\alpha \in I(\Gamma)$, the closed stratum $\overline{X}_\alpha^* \subset X^*$ is defined to be the subspace $\overline{X}_\alpha^\perp$. Equivalently, if $\overline{X}_\alpha \subset X$ is the intersection of hyperplanes $H_j, j \in J_\alpha$, then $\overline{X}_\alpha^*$ is the sum of the one-dimensional subspaces $H_j^\perp, j \in J_\alpha$.

Clearly, the adjacencies of strata in X^* are described by edges of Γ . Thus the adjacency graph of the stratified space X^* is naturally isomorphic to the adjacency graph Γ of the stratified space X .

4.11.3. Having a quiver $\mathcal{V} = \{V_\alpha, A_{\alpha,\beta}\}$ of the graph Γ of the arrangement $\mathcal{C}$, we shall construct a $\mathcal{D}_{X^*}$ -module denoted by $\tilde{E}\mathcal{V}$ and called *the quiver $\mathcal{D}_{X^*}$ -module associated with the line arrangement $\mathcal{C}^\perp$ and quiver $\mathcal{V}$* .

Remark. All our previous constructions were associated with $\mathcal{D}$ modules on X . This is the first construction which considers $\mathcal{D}$ -modules on the dual space X^* .

The construction of the $\mathcal{D}_{X^*}$ -module $\tilde{E}\mathcal{V}$ is analogous to the construction of the $\mathcal{D}_X$ module $E\mathcal{V}$ in Section 4.2. Namely, the $\mathcal{D}_{X^*}$ -module $\tilde{E}\mathcal{V}$ is the quotient of the free sheaf $\mathcal{R}^0 \tilde{E}\mathcal{V} = \mathcal{D}_{X^*} \otimes_{\mathbb{C}} (\oplus_{\alpha \in I(\Gamma)} \overline{\Omega}_\alpha \otimes V_\alpha)$ of $\mathcal{D}_{X^*}$ -modules over its subsheaf $\mathcal{R}' \tilde{E}\mathcal{V}$ of coherent $\mathcal{D}_{X^*}$ -modules, whose global sections are the following global sections of $\mathcal{R}^0 \tilde{E}\mathcal{V}$:

(i) the sections

$$u \xi \otimes \overline{\omega}_\alpha \otimes v_\alpha - \sum_{\beta, \alpha \succ \beta} \langle \xi, f_{\beta,\alpha} \rangle u \otimes \pi_{\beta,\alpha}(\overline{\omega}_\alpha) \otimes A_{\beta,\alpha}(v_\alpha) ,$$

where $\alpha \in I(\Gamma)$, $u \otimes \overline{\omega}_\alpha \otimes v_\alpha \in \mathcal{D}_{X^*} \otimes_{\mathbb{C}} \overline{\Omega}_\alpha \otimes V_\alpha$ and $\xi \in T_\alpha$,

(ii) the sections

$$u f \otimes \overline{\omega}_\alpha \otimes v_\alpha - \sum_{\beta, \beta \succ \alpha} \langle \xi_{\beta,\alpha}, f \rangle u \otimes \pi_{\beta,\alpha}(\overline{\omega}_\alpha) \otimes A_{\beta,\alpha}(v_\alpha) ,$$

where $\alpha \in I(\Gamma)$, $u \otimes \overline{\omega}_\alpha \otimes v_\alpha \in \mathcal{D}_{X^*} \otimes_{\mathbb{C}} \overline{\Omega}_\alpha \otimes V_\alpha$ and $f \in F_\alpha$.

Here $\overline{\Omega}_\alpha$ denotes the one-dimensional space of the top degree holomorphic differential forms on X_α^* , invariant with respect to translations along X_α^* , and $\{f_{\beta,\alpha}, \xi_{\alpha,\beta} \mid \alpha, \beta \in I(\Gamma), \alpha \succ \beta\}$ is an edge framing of $\mathcal{C}^\perp$.

It is easy to see that $\tilde{E}\mathcal{V}$ does not depend on the choice of the edge framing.

4.11.4. Given a quiver $\mathcal{V}$, denote by $\tilde{\mathcal{V}}$ the quiver $\tilde{\mathcal{V}} = \{W_\alpha, B_{\alpha,\beta}\}$, where $W_\alpha = V_\alpha$ and $B_{\alpha,\beta} = \varepsilon(\beta, \alpha) A_{\alpha,\beta}$.

Proposition 4.16. *The Fourier transform of the quiver $\mathcal{D}_X$ -module $E\mathcal{V}$ is isomorphic to the quiver $\mathcal{D}_{X^*}$ -module $\tilde{E}\tilde{\mathcal{V}}$.*

The proof is by easy calculations. The choice of the isomorphism depends on the choice of a top degree differential holomorphic form on X^* , invariant with respect to translations.

Remark. The constructions of Section 4.11 show that quiver $\mathcal{D}$ -modules can be defined not only for hyperplane arrangements but for central line arrangements as well. It is easy to see that the quiver $\mathcal{D}$ -modules of line arrangements are all holonomic with regular singularities. They admit an explicit free resolution and have all nice properties of quiver $\mathcal{D}$ -modules of hyperplane arrangements.

5. QUIVER PERVERSE SHEAVES

5.1. Perverse sheaves over analytic varieties. Let Y be a smooth analytic variety over $\mathbb{C}$ of dimension N . Let $\mathcal{S}$ be a stratification of Y by smooth subvarieties, satisfying equisingularity conditions. It means that Y is presented as a finite disjoint union of strata, such that the closure of each stratum is a union of strata, and for any stratum S and any

points $x, y \in S$, there exists a diffeomorphism ϕ of Y , preserving the stratification and such that $\phi(x) = y$.

Denote by Y^k , $k = 0, 1, \dots$, the closure of the union of all strata of codimension k . The subspaces Y^k form a filtration:

$$Y = Y^0 \supset Y^1 \supset \dots \supset Y^N \supset Y^{N+1} = Y^{N+2} = \dots = \emptyset .$$

A complex $\mathcal{F}$ of sheaves of vector spaces with differential of degree one is called a perverse sheaf with respect to the stratification if it satisfies the following conditions on supports of its cohomology groups [GMc] :

$$(5.1) \quad \text{supp } H^{-N \pm k}(\mathcal{F}) \subset Y^k$$

for all $k = 0, 1, \dots$. A complex of sheaves $\mathcal{F} \in D^b(Y, \mathbb{C})$ of vector spaces over Y is called a perverse sheaf if it is perverse with respect to some stratification of Y , satisfying the conditions above.

Any continuous map $j : Y_1 \rightarrow Y_2$ of topological spaces determines the functor $j^\bullet : Sh(Y_2, \mathbb{C}) \rightarrow Sh(Y_1, \mathbb{C})$ of inverse image and two functors $j_\bullet : Sh(Y_1, \mathbb{C}) \rightarrow Sh(Y_2, \mathbb{C})$ and $j_! : Sh(Y_1, \mathbb{C}) \rightarrow Sh(Y_2, \mathbb{C})$ of direct image.

For any sheaf $\mathcal{G} \in Sh(Y_2, \mathbb{C})$ and a point $y_1 \in Y_1$, the stalk $j^\bullet \mathcal{G}_{y_1}$ of $j^\bullet \mathcal{G}$ at $y_1 \in Y_1$ is the stalk $\mathcal{G}_{j(y_1)}$. For any sheaf $\mathcal{F} \in Sh(Y_1, \mathbb{C})$ and an open $U \subset Y_2$, the sections $\Gamma(U, j_\bullet \mathcal{F})$ of $j_\bullet \mathcal{F}$ are

$$(5.2) \quad \Gamma(U, j_\bullet \mathcal{F}) = \Gamma(j^{-1}(U), \mathcal{F}) .$$

The sections $\Gamma(U, j_! \mathcal{F})$ are those sections $s \in \Gamma(j^{-1}(U), \mathcal{F})$, for which the map $j : \text{supp}(s) \rightarrow U$ is proper.

Both functors $j_\bullet$ and $j_!$ are left exact. The functor $j^\bullet$ is exact. Denote by $Rj_! : D^b(Y_1, \mathbb{C}) \rightarrow D^b(Y_2, \mathbb{C})$ and $Rj_\bullet : D^b(Y_1, \mathbb{C}) \rightarrow D^b(Y_2, \mathbb{C})$ the corresponding derived functors.

For a topological space Y denote by $\overline{\Delta}_Y : D^b(Y, \mathbb{C}) \rightarrow D^b(Y, \mathbb{C})$ the Poincare-Verdier duality,

$$\overline{\Delta}_Y(\mathcal{F}) = R\text{Hom}(\mathcal{F}, \mathbf{D}_Y) ,$$

where $\mathbf{D}_Y$ is the dualizing complex. For a smooth complex analytic variety Y the dualizing complex is isomorphic in $D^b(Y, \mathbb{C})$ to the shifted constant sheaf:

$$\mathbf{D}_Y = \mathbb{C}_Y[2 \dim_{\mathbb{C}} Y] .$$

For any $\mathcal{F} \in D^b(Y_1, \mathbb{C})$ there is an isomorphism in $D^b(Y_2, \mathbb{C})$:

$$Rj_! \approx \overline{\Delta}_{Y_2} Rj_\bullet \overline{\Delta}_{Y_1} .$$

Equalities (5.2) determine isomorphisms of adjunction

$$\alpha_{\mathcal{G}, \mathcal{F}} : \text{Hom}_{Sh(Y_2, \mathbb{C})}(\mathcal{G}, j_\bullet(\mathcal{F})) \xrightarrow{\sim} \text{Hom}_{Sh(Y_1, \mathbb{C})}(j^\bullet(\mathcal{G}), \mathcal{F}) ,$$

which extend to isomorphisms of adjunction in derived categories,

$$\overline{\alpha}_{\mathcal{G}, \mathcal{F}} : \text{Hom}_{D^b(Y_2, \mathbb{C})}(\mathcal{G}, Rj_\bullet(\mathcal{F})) \xrightarrow{\sim} \text{Hom}_{D^b(Y_1, \mathbb{C})}(j^\bullet(\mathcal{G}), \mathcal{F}) ,$$

with respect to which the functor $Rj_\bullet : D^b(Y_1, \mathbb{C}) \rightarrow D^b(Y_2, \mathbb{C})$ is right adjoint to the functor $j^\bullet : D^b(Y_2, \mathbb{C}) \rightarrow D^b(Y_1, \mathbb{C})$.

If the map $j : Y_1 \rightarrow Y_2$ is an open embedding, then the functor $j_!$ is exact, and its extension to derived categories: $j_! : D^b(Y_1, \mathbb{C}) \rightarrow D^b(Y_2, \mathbb{C})$ is left adjoint to the functor of inverse image $j^* : D^b(Y_2, \mathbb{C}) \rightarrow D^b(Y_1, \mathbb{C})$.

Let $j : Y_1 \rightarrow Y_2$ be an open embedding of a smooth affine variety Y_1 to a smooth algebraic variety Y_2 . Denote by the same symbol $j : Y_1^{an} \rightarrow Y_2^{an}$ the corresponding map of complex analytic varieties. Then the functors Rj_* and $j_!$ map perverse sheaves to perverse sheaves. In this situation the *MacPherson extension* $j_{!*}(\mathcal{F})$ of a perverse sheaf $\mathcal{F}$ over Y_1^{an} is defined as the image of $j_!\mathcal{F}$ in $Rj_*\mathcal{F}$ with respect to the canonical map $s_j : j_!(\mathcal{F}) \rightarrow Rj_*(\mathcal{F})$, determined by the two pairs (j^*, Rj_*) and $(j_!, j^*)$ of adjoint functors, see Section 3.3. The MacPherson extension is a perverse sheaf over Y_2^{an} whose restriction to Y_1^{an} is isomorphic to $\mathcal{F}$ in the category $D^b(Y_1^{an}, \mathbb{C})$.

5.2. De Rham functor. Let Y be a smooth algebraic variety over $\mathbb{C}$. Denote by Y^{an} the same variety considered as an analytic variety. Denote by $\mathcal{O}_Y^{an}$ the sheaf of analytic functions on Y^{an} and by $\mathcal{D}_Y^{an}$ the sheaf of rings of differential operators on Y^{an} with analytic coefficients, $\mathcal{D}_Y^{an} = \mathcal{O}_Y^{an} \otimes_{\mathcal{O}_Y} \mathcal{D}_Y$. We assign to any $\mathcal{D}_Y$ -module M the corresponding $\mathcal{D}_Y^{an}$ -module M^{an} ,

$$M^{an} = \mathcal{O}_Y^{an} \otimes_{\mathcal{O}_Y} M.$$

The sheaf Ω_Y of top exterior forms on Y has a structure of the right $\mathcal{D}_Y$ -module. Differential operators of degree zero (functions) act on the top forms by means of pointwise multiplication and vector fields act by means of minus Lie derivative:

$$(5.3) \quad \omega \cdot f = f\omega, \quad \omega \cdot \xi = -Lie_\xi(\omega).$$

Here $f \in \mathcal{O}_Y$, $\omega \in \Omega_Y$, ξ is a vector field on Y .

Let M be a coherent $\mathcal{D}_Y$ module. Then the assignment $M \rightarrow \Omega_Y^{an} \otimes_{\mathcal{D}_Y^{an}} M^{an}$ defines a functor from the category $\mathcal{M}_Y$ of coherent $\mathcal{D}_Y$ -modules to the category $Sh(Y^{an}, \mathbb{C})$ of sheaves of complex vector spaces over Y^{an} . This functor is right exact.

The left derived functor of this functor is called the de Rham functor:

$$(5.4) \quad DR : D^b(\mathcal{M}_Y) \rightarrow D^b(Y^{an}, \mathbb{C}), \quad DR(M^\bullet) = \Omega_Y^{an} \overset{L}{\otimes}_{\mathcal{D}_Y^{an}} (M^\bullet)^{an}$$

To calculate the de Rham functor $DR(M^\bullet)$, the class of free $\mathcal{D}_Y$ -modules can be used: if $\tilde{M}^\bullet$ is a complex of free $\mathcal{D}_Y$ -modules quasi-isomorphic to $M^\bullet$, then

$$DR(M^\bullet) = \Omega_Y^{an} \otimes_{\mathcal{D}_Y^{an}} (\tilde{M}^\bullet)^{an}.$$

Another possibility is to replace the right $\mathcal{D}_Y$ -module Ω_Y by its free $\mathcal{D}_Y$ -resolution $\tilde{\Omega}_Y^\bullet$, then

$$DR(M^\bullet) = (\tilde{\Omega}_Y^\bullet)^{an} \otimes_{\mathcal{D}_Y^{an}} (M^\bullet)^{an}.$$

A free $\mathcal{D}_Y$ -resolution $\tilde{\Omega}_Y^\bullet$ of the right $\mathcal{D}_Y$ -module Ω_Y looks as follows:

$$(5.5) \quad 0 \rightarrow \mathcal{D}_Y \rightarrow \Omega_Y^1 \otimes_{\mathcal{O}_Y} \mathcal{D}_Y \rightarrow \cdots \rightarrow \Omega_Y^{N-1} \otimes_{\mathcal{O}_Y} \mathcal{D}_Y \rightarrow \Omega_Y^N \otimes_{\mathcal{O}_Y} \mathcal{D}_Y \rightarrow \Omega_Y^N \rightarrow 0$$

where $N = \dim Y$, Ω_Y^k is the sheaf of exterior k -forms over Y . In local coordinates $z_1, \dots, z_N$, the differential is

$$(5.6) \quad d(\omega \otimes D) = d\omega \otimes D + \sum_{i=1}^N (dz_i \wedge \omega) \otimes \frac{\partial}{\partial z_i} D.$$

Here $\omega \in \Omega_Y^k$, $D \in \mathcal{D}_Y$.

Let M be a coherent $\mathcal{D}_Y$ -module. Then, due to (5.5), the complex of sheaves $DR(M) = DR(M^{(\cdot)})$ is isomorphic in $\mathcal{D}^b(Y^{an}, \mathbb{C})$ to the complex

$$(5.7) \quad 0 \rightarrow M^{an} \rightarrow (\Omega_Y^1)^{an} \otimes_{\mathcal{O}_Y^{an}} M^{an} \rightarrow \cdots \rightarrow (\Omega_Y^N)^{an} \otimes_{\mathcal{O}_Y^{an}} M^{an} \rightarrow 0$$

where $\deg((\Omega_Y^{N-k})^{an} \otimes_{\mathcal{O}_Y^{an}} M^{an}) = -k$ and the differential is induced from (5.6).

Example. Let $Y = X^0$ be the complement to the arrangement $\mathcal{C}$ in $X = C^N$. For each quiver $\mathcal{V}_{\Gamma^0} = \{V, A^i\} \in Qui_{\Gamma^0}$, the $\mathcal{D}_{X^0}$ -module $E^0 \mathcal{V}_{\Gamma^0}$ is a finitely generated free $\mathcal{O}_{X^0}$ -module. By the Poincare lemma the cohomology groups of the complex (5.7) in this case are nonzero only in degree $-N$. Thus we have an identification in $D^b(Y^{an}, \mathbb{C})$:

$$(5.8) \quad DR(E^0 \mathcal{V}_{\Gamma^0}) \approx \mathcal{L}\mathcal{V}_{\Gamma^0}[N]^{(\cdot)}$$

where $\mathcal{L}\mathcal{V}_{\Gamma^0}$ is the locally constant sheaf (local system) of flat sections of the corresponding connection (4.38), and $C[N]^\bullet$ means the shift of degree by N , $C[N]^k = C^{N+k}$.

5.3. De Rham complex of a quiver $\mathcal{D}_X$ -module. For any quiver $\mathcal{V} \in Qui$, denote by $\mathcal{QV}$ the tensor product over $\mathcal{D}_X^{an}$ of the right $\mathcal{D}_X^{an}$ -module Ω_X^{an} and the complex of left $\mathcal{D}_X^{an}$ -modules $\mathcal{RE}\mathcal{V}^{an}$:

$$(5.9) \quad \mathcal{QV} = \Omega_X^{an} \otimes_{\mathcal{D}_X^{an}} \mathcal{RE}\mathcal{V}^{an}.$$

Theorem 4.13 implies

Corollary 5.1. *Let $\mathcal{V} \in Qui_\Gamma$ be a quiver of Γ . Then the de Rham complex of the quiver $\mathcal{D}_X$ -module $E\mathcal{V}$ is isomorphic in $D^b(X^{an}, \mathbb{C})$ to the complex of sheaves $\mathcal{QV}$.*

Due to the description in (4.44) and (4.47) of the resolution $\overline{\mathcal{RE}}\mathcal{V}$, we get the following description of the complex $\mathcal{QV}$: $0 \rightarrow \mathcal{Q}^{-N}\mathcal{V} \xrightarrow{d} \cdots \xrightarrow{d} \mathcal{Q}^0\mathcal{V}^0 \rightarrow 0$ with $\deg \mathcal{Q}^r\mathcal{V} = r$:

$$(5.10) \quad \mathcal{Q}^r\mathcal{V} = \bigoplus_{\alpha \in I(\Gamma)} \Omega_X^{an} \otimes \left(\bigwedge^{-r} \mathcal{T}_\alpha \otimes \overline{\Omega}_\alpha \otimes V_\alpha \right)$$

with differential $d = d_1 - d_0$, where

$$(5.11) \quad \begin{aligned} d_1(\omega \otimes t_1 \wedge \cdots \wedge t_r \otimes \overline{\omega}_\alpha \otimes v_\alpha) &= \sum_{i=1}^r (-1)^{i+1} (\omega) \cdot t_i \otimes t_1 \wedge \cdots \wedge \widehat{t_i} \cdots \wedge t_r \otimes \overline{\omega}_\alpha \otimes v_\alpha, \\ d_0(\omega \otimes t_1 \wedge \cdots \wedge t_r \otimes \overline{\omega}_\alpha \otimes v_\alpha) &= \sum_{\beta, (\alpha, \beta) \in E(\Gamma)} \omega \otimes A_{\beta, \alpha}^r(t_1 \wedge \cdots \wedge t_r \otimes \overline{\omega}_\alpha \otimes v_\alpha), \end{aligned}$$

for $\omega \in \Omega_X^{an}$, $t_i \in \mathcal{T}_\alpha$, $\overline{\omega}_\alpha \otimes v_\alpha \in \overline{\Omega}_\alpha \otimes V_\alpha$.

Analogously, Theorem 4.11 implies

Corollary 5.2. *For any Γ^0 quiver $\mathcal{V}_{\Gamma^0} = \{V, A^i\}$, the de Rham complex $DR(E^0 \mathcal{V}_{\Gamma^0})$ is isomorphic in $D^b(X^{0, an}, \mathbb{C})$ to the complex $\mathcal{QV}_{\Gamma^0} = \Omega_{X^0}^{an} \otimes_{\mathcal{D}_{X^0}^{an}} \mathcal{RE}^0 \mathcal{V}_{\Gamma^0}^{an}$, where*

$$\mathcal{Q}^r \mathcal{V}_{\Gamma^0} = \Omega_{X^0}^{an} \otimes \left(\bigwedge^{-r} T_\emptyset \otimes \overline{\Omega}_\emptyset \otimes V \right),$$

and the differential is $d = d_1 - d_0$,

$$\begin{aligned} d_1(\omega \otimes \xi_1 \wedge \cdots \wedge \xi_r \otimes \bar{\omega}_\emptyset \otimes v) &= \sum_{i=1}^r (-1)^{i+1} (\omega) \cdot \xi_i \otimes \xi_1 \wedge \cdots \wedge \widehat{\xi_i} \cdots \wedge \xi_r \otimes \bar{\omega}_\emptyset \otimes v , \\ d_0(\omega \otimes \xi_1 \wedge \cdots \wedge \xi_r \otimes \bar{\omega}_\emptyset \otimes v) &= \sum_{j \in J(C)} \omega \cdot f_j^{-1} \otimes i_{f_j}(\xi_1 \wedge \cdots \wedge \xi_r) \otimes \bar{\omega}_\emptyset \otimes A_j(v) . \end{aligned}$$

Notice that this complex is isomorphic in $D^b(X^{0,an}, \mathbb{C})$ to the complex (5.7).

5.4. Quiver perverse sheaves. It is known [K2, Mb, B] that for any smooth algebraic variety Y , the de Rham functor transforms holonomic $\mathcal{D}_Y$ -modules to perverse sheaves. Moreover, if M is a holonomic $\mathcal{D}_Y$ -module such that its singular support $ss.M$ belongs to the union of conormal bundles of nonsingular subvarieties $Z_i \subset Y$,

$$ss.M \subset \bigcap_i T_{Z_i}^* Y ,$$

then the complex of sheaves $DR(M)$ is a perverse sheaf with respect to any equisingular stratification of Y^{an} , such that each Z_i^{an} is a union of strata.

Corollary 5.1 and Proposition 4.2 imply

Proposition 5.3. *Let $\mathcal{V} \in Qui$ be a quiver of Γ . Then the complex of sheaves $\mathcal{QV}$ defined by (5.9) is a perverse sheaf on X^{an} with respect to the stratification $\{X_\alpha^{an}, \alpha \in I(\Gamma)\}$.*

Analogously, for any Γ^0 -quiver $\mathcal{V}_{\Gamma^0} \in Qui_{\Gamma^0}$, the complex of sheaves $\mathcal{QV}_{\Gamma^0}$ is a perverse sheaf over $X^{0,an}$, isomorphic in $D^b(X^{0,an}, \mathbb{C})$ to the shifted by N locally constant sheaf $\mathcal{LV}_{\Gamma^0}[N]$, see (4.38).

The de Rham functor respects the duality and all of the functors of direct and inverse image. In particular, for a smooth complex algebraic variety Y and $M \in \mathcal{M}_Y^{hol}$, we have [K1, B]

$$(5.12) \quad DR(\Delta_Y(M)) \approx \overline{\Delta}_{Y^{an}}(DR(M)) ,$$

For an open embedding $j : Y_1 \rightarrow Y_2$ of a smooth affine variety Y_1 into a smooth algebraic variety Y_2 , any $M_1 \in \mathcal{M}_{Y_1}^{hol}$ and $M_2 \in \mathcal{M}_{Y_2}$, we have

$$\begin{aligned} (5.13) \quad & j^\bullet(DR(M_2)) \approx DR(j^*(M_2)) , \\ & Rj_\bullet(DR(M_1)) \approx DR(j_*(M_1)) , \\ & j_!(DR(M_1)) \approx DR(j_!^{(D)}(M_1)) , \\ & j_{!*}(DR(M_1)) \approx DR(j_{!*}^{(D)}(M_1)) \end{aligned}$$

Corollary 5.4.

(i) *Let $\mathcal{V}_{\Gamma^0}$ be a level zero quiver. Then we have in $D^b(X^{0,an}, \mathbb{C})$:*

$$\overline{\Delta}_{X^{0,an}}(DR(E^0(\mathcal{V}_{\Gamma^0}))) \approx DR(E^0(\tau \mathcal{V}_{\Gamma^0})) .$$

(ii) *Let $\mathcal{V} = \{V_\alpha, A_{\alpha,\beta}\} \in Qui$ be a quiver of Γ . Then we have in $D^b(X^{an}, \mathbb{C})$:*

$$\overline{\Delta}_{X^{an}}(DR(E(\mathcal{V}))) \approx DR(E(\tau \mathcal{V})) .$$

Proof. Both statements follow from Theorem 4.14 and relation (5.12). □

Corollary 5.5. *Let $\mathcal{V}_{\Gamma^0} = \{V, A^i\}$ be a level zero quiver. Suppose that each operator A^i has a single eigenvalue and $\mathcal{V}_{\Gamma^0}$ has a non-resonant spectrum, or suppose that all operators A^i are close to zero. Then the following perverse sheaves are isomorphic:*

$$\begin{aligned} R(j_0)_* DR(E^0(\mathcal{V}_{\Gamma^0})) &\approx DR(E(\mathbb{J}_{0,*}(\mathcal{V}_{\Gamma^0}))) , \\ (j_0)_! DR(E^0(\mathcal{V}_{\Gamma^0})) &\approx DR(E(\mathbb{J}_{0,!}(\mathcal{V}_{\Gamma^0}))) , \\ (j_0)_! DR(E^0(\mathcal{V}_{\Gamma^0})) &\approx DR(E(\mathbb{J}_{0,!*}(\mathcal{V}_{\Gamma^0}))) . \end{aligned}$$

Proof. The first statement follows from Corollary 4.7 and relation (5.13). The second and third statements follow from Corollary 4.15 and relation (5.13). $\square$

5.5. Cohomology groups of $\mathbb{C}^N$ with coefficients in quiver perverse sheaves.
We consider $X = \mathbb{C}^N$ as an analytic variety. For a quiver $\mathcal{V}$, the complex $\mathcal{QV}$ is a complex of free $\mathcal{O}_X$ -modules. The affine space $X = \mathbb{C}^N$ is a Stein space,

$$H^k(X, \mathcal{O}^{an}) = 0, \quad \text{for all } k \neq 0 .$$

Hence the (hyper)cohomology groups of X with coefficients in the complex $\mathcal{QV}$ of sheaves are isomorphic to the cohomology groups of the complex $\overline{\mathcal{QV}} = \Gamma(X, \mathcal{QV})$ of global sections of $\mathcal{QV}$:

$$(5.14) \quad H^k(X, \mathcal{QV}) \simeq H^k(\overline{\mathcal{QV}}) .$$

Theorem 5.6. *Let $\mathcal{C}$ be a central arrangement (that is all hyperplanes of $\mathcal{C}$ contain $0 \in \mathbb{C}^N$). Let $\mathcal{V} \in \text{Qui}_{\Gamma}$ be a quiver with all linear maps $A_{\alpha,\beta}$ of the quiver close to zero. Then the complex $\overline{\mathcal{QV}}$ of global sections of the complex (5.10)-(5.11) is quasi-isomorphic to the shifted by N quiver complex $C_+(\mathcal{V})$, introduced in (3.1)-(3.2):*

$$\overline{\mathcal{QV}} \stackrel{quis}{\simeq} C_+(\mathcal{V})[N] .$$

Note that $C_+(\mathcal{V})$ is a finite dimensional complex.

The theorem is proved in Section 7.

Theorem 5.6 and the isomorphism in (5.14) imply the following statement.

Corollary 5.7. *Let $\mathcal{C}$ be a central arrangement and $\mathcal{V} \in \text{Qui}_{\Gamma}$ a quiver with all linear maps $A_{\alpha,\beta}$ close to zero. Then for any $k \in \mathbb{Z}$,*

$$H^k(X, DR(E\mathcal{V})) \simeq H^k(X, \mathcal{QV}) \simeq H^{k+N}(C_+(\mathcal{V})) .$$

Let X^0 denote the complement to $\mathcal{C}$ in $\mathbb{C}^N$. For a local system $\mathcal{L}$ on X^0 introduce the intersection cohomology groups $IH^k(X, \mathcal{L})$ as the cohomology groups $H^{k-N}(X, (j_0)_! \mathcal{L}[N])$ of the MacPherson extension of the perverse sheaf $\mathcal{L}[N] \in D^b(X^0, \mathbb{C})$,

$$(5.15) \quad IH^k(X, \mathcal{L}) \stackrel{def}{=} H^{k-N}(X, (j_0)_! \mathcal{L}[N]) .$$

Let $\mathcal{LV}$ denote the local system of flat sections of the connection (4.38).

Corollary 5.8. *Let $\mathcal{V} = \{V, A^i\}$ be a level zero quiver of a central arrangement $\mathcal{C}$. Assume that all operators A^i are close to zero. Then for any $k \in \mathbb{Z}$,*

$$\begin{aligned} H^k(X^0, \mathcal{LV}) &\simeq H^k(C_+(\mathbb{J}_{0,*}\mathcal{V})) , \\ IH^k(X, \mathcal{LV}) &\simeq H^k(C_+(\mathbb{J}_{0,!*}\mathcal{V})) . \end{aligned}$$

Proof. The conditions on operators A^i imply that the quiver $\mathcal{V}$ has a non-resonant spectrum and Corollary 5.5 can be applied to the quiver $\mathbb{J}_{0,!}\mathcal{V}$. By (5.8), $\mathcal{LV} \approx DR(E^0\mathcal{V})[-N]$ and

$$\begin{aligned} H^k(X^0, \mathcal{LV}) &\simeq H^k(X, R(j_0)_*\mathcal{LV}) \simeq H^k(X, DR(\mathbb{J}_{0,*}\mathcal{V})[-N]) \simeq H^k(C_+(\mathbb{J}_{0,*}\mathcal{V})) , \\ IH^k(X, \mathcal{LV}) &\simeq H^{k-N}(X, (j_0)_!*\mathcal{LV}[N]) \simeq H^{k-N}(X, DR(\mathbb{J}_{0,!}\mathcal{V})) \simeq H^k(C_+(\mathbb{J}_{0,!}\mathcal{V})) . \end{aligned}$$

The second isomorphism in both lines is due to Corollary 5.5, the third isomorphism in both lines is due to Corollary 5.7. $\square$

Example. Let $\mathcal{C}$ be a central arrangement in X . Let $a_j, j \in I_1(\mathcal{C})$, and κ be complex numbers. Consider the trivial bundle $\mathbb{C} \times X^0 \rightarrow X^0$ with the connection given by the differential one-form $\frac{1}{\kappa} \sum_i a_i \frac{df_i}{f_i}$. Here $f_i = 0$ is the equation of the hyperplane H_i of $\mathcal{C}$. Let $\mathcal{LV}$ be the local system of flat sections of the connection.

Let $\mathcal{V} = \{\mathbb{C}, A^i\}$ be the one-dimensional level zero quiver of Γ^0 , where $A^i : \mathbb{C} \rightarrow \mathbb{C}$ is the operator of multiplication by a_i/κ .

Let $(\mathcal{A}^\bullet(\mathcal{C}), d_a)$ be the Aomoto complex, defined for $\mathcal{C}$ in Section 2.5. Let $(S_a(\mathcal{F}^\bullet(\mathcal{C})), d_a)$ be its subcomplex of flag forms.

By Corollary 5.8, for $|\kappa| \gg 1$, we have

$$H^\bullet(\mathcal{A}^\bullet(\mathcal{C}), d_a) \simeq H^\bullet(X^0, \mathcal{LV}) , \quad H^\bullet(S_a(\mathcal{F}^\bullet(\mathcal{C})), d_a) \simeq IH^\bullet(X, \mathcal{L}) .$$

Here $H^\bullet(X^0, \mathcal{LV})$ are the cohomology groups of X^0 with coefficients in the local system $\mathcal{LV}$ and $IH^\bullet(X, \mathcal{L})$ are the intersection cohomology groups of X^0 with coefficients in the local system $\mathcal{LV}$.

We see that the calculation of cohomology groups of X with coefficients in the perverse sheaf associated with the quiver MacPherson extension of a level zero quiver $\mathcal{V}_{\Gamma^0}$ gives a topological meaning to the complex of the flag forms. One of the aims of this paper was to give a topological meaning to the complex of flag forms.

6. EQUIVARIANT STRUCTURES

6.1. Arrangement with a group action. Suppose a finite group G acts on $X = \mathbb{C}^N$ by affine transformations $L_X(g) : X \rightarrow X$ preserving an arrangement $\mathcal{C}$. This means that G acts on the set $J(\mathcal{C})$ so that $L_X(g)(H_j) = H_{g(j)}$ for $g \in G, j \in J(\mathcal{C})$.

The action of G preserves the principal open subsets X^n .

The group G naturally acts on the graph Γ of the arrangement and on the truncated graphs Γ^n .

The G -action on X induces a G -action on the ring D_X . For an affine transformation $L : X \rightarrow X$, the associated automorphism $L_{D_X} : D_X \rightarrow D_X$ is defined by the formula $L_{D_X}(f)(z) = f(L^{-1}z)$ for a function f on X , and by the formula $L_{D_X}(\eta)(z) = dL \cdot \eta(L^{-1}z)$ for a vector field η . Here dL is the derivative of L .

For $g \in G$, the transformation $L_{D_X}(g) : D_X \rightarrow D_X$ preserves the space F of affine functions on X and for $\alpha \in I(\Gamma)$, maps the subspace $F_\alpha \subset F$ to $F_{g(\alpha)}$.

The transformation $L_{D_X}(g)$ preserves the space T of vector fields on X with constant coefficients and for any $\alpha \in I(\Gamma)$, maps the subspace $T_\alpha \subset T$ to $T_{g(\alpha)}$.

The G -action on X induces a G -action on the algebra Ω_X^* of differential forms. For an affine transformation $L : X \rightarrow X$, the associated automorphism $L_{\Omega_X^*} : \Omega_X^* \rightarrow \Omega_X^*$ is defined by the formula $L_{\Omega_X^*}(\omega)(z) = ((dL)^t)^{-1}(\omega)(L^{-1}(z))$ for $w \in \Omega_X^1$.

For $g \in G$ and $\alpha \in I(\Gamma)$, the transformation $L_{\Omega_X^*}(g)$ maps $\overline{\Omega}_\alpha$ to $\overline{\Omega}_{g(\alpha)}$.

6.2. Equivariant quivers. An equivariant quiver (or a quiver with a group action) $\mathcal{V}_G = \{V_\alpha, A_{\alpha,\beta}, L_{V_\alpha}(g)\}$ is a quiver $\mathcal{V} = \{V_\alpha, A_{\alpha,\beta}\}$ together with a collection of linear maps $L_{V_\alpha}(g) : V_\alpha \rightarrow V_{g(\alpha)}$, for $\alpha \in I(\Gamma)$ and $g \in G$, such that for $\alpha, \beta \in I(\Gamma)$, we have

$$\begin{aligned} & \bullet L_{V_{g''(\alpha)}}(g') L_{V_\alpha}(g'') = L_{V_\alpha}(g'g''), \\ & \bullet L_{V_\alpha}(g^{-1}) = (L_{V_{g^{-1}(\alpha)}}(g))^{-1}, \\ & \bullet L_{V_\alpha}(g) A_{\alpha,\beta} = A_{g(\alpha),g(\beta)} L_{V_\beta}(g). \end{aligned}$$

In the same manner we define level l G -equivariant quivers.

Denote by $Qui_G = Qui_{\Gamma,G}$ the category of G -equivariant quivers, by $Qui_{\Gamma^l,G}$ the category of level l G -equivariant quivers and by $\mathcal{N}^l : Qui_{\Gamma^l,G} \rightarrow Qui_{\Gamma^l}$ the forgetful functor, which assigns to an equivariant quiver $\mathcal{V}_{\Gamma^l,G}$ the same quiver with no morphisms of group action. In this case we will use notation $\mathcal{V}_{\Gamma^l}$ for the quiver $\mathcal{N}^l(\mathcal{V}_{\Gamma^l,G})$.

Let $V_{\Gamma^l,G} = \{V_\alpha, A_{\alpha,\beta}, A_\alpha^\beta, L_{V_\alpha}(g)\}$ be a level l equivariant quiver. Then for $k < l$, the collection of maps $L_{V_\alpha}(g) : V_\alpha \rightarrow V_\alpha$, for $g \in G$ and $\alpha \in I(\Gamma^k)$, determines a group action on the level k quiver $\mathbb{J}_{l,k}^*(V_{\Gamma^l})$.

This assignment determines a functor $\mathbb{J}_{l,k,G}^* : Qui_{\Gamma^l,G} \rightarrow Qui_{\Gamma^k,G}$. This functor commutes with forgetful functors,

$$\mathcal{N}^k \mathbb{J}_{l,k,G}^* = \mathbb{J}_{l,k}^* \mathcal{N}^l.$$

The functor $\mathbb{J}_{l,k,G}^* : Qui_{\Gamma^l,G} \rightarrow Qui_{\Gamma^k,G}$ admits left and right adjoint functors. Let $\mathbb{J}_{l,k,*,G} : Qui_{\Gamma^k,G} \rightarrow Qui_{\Gamma^l,G}$ be the right adjoint functor to $\mathbb{J}_{l,k,G}^*$. Let $\mathbb{J}_{l,k,!},G : Qui_{\Gamma^k,G} \rightarrow Qui_{\Gamma^l,G}$ be the left adjoint functor to $\mathbb{J}_{l,k,G}^*$. We have

$$\mathcal{N}^l \mathbb{J}_{l,k,*,G} = \mathbb{J}_{l,k,*} \mathcal{N}^k, \quad \mathcal{N}^l \mathbb{J}_{l,k,!},G = \mathbb{J}_{l,k,!} \mathcal{N}^k.$$

6.3. Equivariant quiver $\mathcal{D}$ -modules. Let X be a topological space. Any homeomorphism $a : X \rightarrow X$ determines an exact functor $a^* : Sh(X, \mathbb{C}) \rightarrow Sh(X, \mathbb{C})$ of inverse image of sheaves of vector spaces, which can be described as follows. The fiber $a^* \mathcal{F}_x$ of the sheaf $a^* \mathcal{F}$ at a point $x \in X$ is the fiber $\mathcal{F}_{a(x)}$ of $\mathcal{F}$ at the point $a(x)$, and for any morphism $\varphi : \mathcal{F} \rightarrow \mathcal{G}$ of sheaves over X the map of fibers $a^* \varphi(x) : a^* \mathcal{F}_x \rightarrow a^* \mathcal{G}_x$ coincides with the map $\varphi(a(x)) : \mathcal{F}_{a(x)} \rightarrow \mathcal{G}_{a(x)}$.

Suppose a finite group G acts on X by homeomorphisms $L_X(g)$. For $\mathcal{F} \in Sh(X, \mathbb{C})$ and $g \in G$, denote by $\mathcal{F}^g$ the sheaf $(L_X(g)^{-1})^* \mathcal{F}$. A sheaf $\mathcal{F} \in Sh(X, \mathbb{C})$ is called G -equivariant, if for $g \in G$, a morphism $L_{\mathcal{F}}(g) : \mathcal{F} \rightarrow \mathcal{F}^g$ is given such that

$$\begin{aligned} & \bullet L_{\mathcal{F}^{g_2}}(g_1) L_{\mathcal{F}}(g_2) = L_{\mathcal{F}}(g_1 g_2), \\ & \bullet L_{\mathcal{F}^g}(g^{-1}) L_{\mathcal{F}}(g) = Id_{\mathcal{F}}. \end{aligned}$$

Here the morphism of sheaves $L_{\mathcal{F}^{g_2}}(g_1) : \mathcal{F}^{g_2} \rightarrow (\mathcal{F}^{g_2})^{g_1}$ is defined by the relation $L_{\mathcal{F}^{g_2}}(g_1) = L_X(g_2^{-1})^* (L_{\mathcal{F}}(g_2^{-1} g_1 g_2))$.

In other words, an equivariant sheaf is a sheaf $\mathcal{F} \in Sh(X, \mathbb{C})$ and a collection of linear maps $L_{\{\mathcal{F}, U\}}(g) : \Gamma(U, \mathcal{F}) \rightarrow \Gamma(L_X(g)(U), \mathcal{F})$, attached to any open set $U \subset X$ and an

element $g \in G$, such that

$$\begin{aligned} L_{\{\mathcal{F}, L_X(g_2)(U)\}}(g_1) L_{\{\mathcal{F}, U\}}(g_2) &= L_{\{\mathcal{F}, U\}}(g_1 g_2) , \\ L_{\{\mathcal{F}, L_X(g)(U)\}}(g^{-1}) L_{\{\mathcal{F}, U\}}(g) &= Id_{\Gamma(U, \mathcal{F})} . \end{aligned}$$

One defines similarly a G -equivariant complex of sheaves of vector spaces.

Denote by $Sh(X, G, \mathbb{C})$ the category of equivariant sheaves over X and by $D^b(X, G, \mathbb{C})$ the corresponding bounded derived category.

Assume that a finite group G acts on a smooth complex algebraic variety X by algebraic automorphisms $L_X(g)$. Then the sheaves $\mathcal{O}_X$, Ω_X^* , and $\mathcal{D}_X$ are equivariant sheaves. The morphisms $L_{\{\mathcal{D}_X, U\}}(g)$ and $L_{\{\mathcal{O}_X, U\}}(g)$ are defined by the relations

$$\begin{aligned} L_{\{\mathcal{D}_X, U\}}(g)(f)(x) &= L_{\{\mathcal{O}_X, U\}}(g)(f)(x) = f(L_X(g^{-1})(x)) , \\ L_{\{\mathcal{D}_X, U\}}(g)(\eta)(x) &= dL_X(g)(\eta)(L_X(g^{-1})(x)) , \end{aligned}$$

where f is a function and η is a vector field on U . The morphisms $L_{\{\Omega_X^*, U\}}(g)$ are defined by the relations

$$(6.1) \quad L_{\{\Omega_X^*, U\}}(g)(\omega)(x) = ((dL_X(g))^t)^{-1}(\omega)(L_X(g^{-1})(x))$$

for one-forms ω .

A sheaf M of $\mathcal{D}_X$ -modules is called G -equivariant if it is equipped with a structure of an equivariant sheaf, such that the maps $L_{\{M, U\}}(g)$ respect the structure of the $\mathcal{D}_X$ -module. This means that

$$L_{\{M, U\}}(g)(d \cdot m) = L_{\{\mathcal{D}_X, U\}}(g)(d) \cdot L_{\{M, U\}}(g)(m) ,$$

for any $m \in \Gamma(U, M)$ and $d \in \Gamma(U, \mathcal{D}_X)$. In particular, the space $\Gamma(X, M)$ of global sections is an equivariant module over the ring $D_X = \Gamma(X, \mathcal{D}_X)$. Denote by $\mathcal{M}_{X,G}$ the category of equivariant coherent $\mathcal{D}_X$ -modules.

If X is an affine space and M is a coherent $\mathcal{D}_X$ -module, then the equivariant $\mathcal{D}_X$ -module of its global sections determines M as an equivariant $\mathcal{D}_X$ -module.

The sheaf $\mathcal{O}_X$ is an equivariant left $\mathcal{D}_X$ -module. The sheaf Ω_X of top exterior forms is an equivariant sheaf of right $\mathcal{D}_X$ -modules.

Let $\mathcal{V}_G = \{V_\alpha, A_{\alpha,\beta}, L_{\mathcal{V}}(g)\} \in Qui_{\Gamma,G}$ be a G -equivariant quiver of Γ . Denote by $\mathcal{V} = \mathcal{N}^N(\mathcal{V}_G)$ the same quiver with no group action. We equip the $\mathcal{D}_X$ -module $E\mathcal{V}$ with a structure of a G -equivariant $\mathcal{D}_X$ -module in the following way.

Choose an edge framing $\{f_{\alpha,\beta}, \xi_{\alpha,\beta}\}$ of the arrangement $\mathcal{C}$. Then the $\mathcal{D}_X$ -module $E\mathcal{V}$ is defined as the quotient of the left free $\mathcal{D}_X$ -module $\mathcal{R}^0 E\mathcal{V} = \bigoplus_{\alpha \in I(\Gamma)} \mathcal{D}_X \otimes \overline{\Omega}_\alpha \otimes V_\alpha$ by the submodule $\mathcal{R}^0 E\mathcal{V}'$, whose global sections are given by (4.25) and (4.26). For $g \in G$, define a map $L_{\mathcal{R}^0 E\mathcal{V}}(g) : \mathcal{R}^0 E\mathcal{V} \rightarrow (\mathcal{R}^0 E\mathcal{V})^g$ by the formula

$$(6.2) \quad L_{\{\mathcal{R}^0 E\mathcal{V}, X\}}(g)(d \otimes \overline{\omega}_\alpha \otimes v_\alpha) = L_{D_X}(g)(d) \otimes L_{\overline{\Omega}_\alpha}(g)(\overline{\omega}_\alpha) \otimes L_{V_\alpha}(g)(v_\alpha)$$

for $\alpha \in I(\Gamma)$ and $d \otimes \overline{\omega}_\alpha \otimes v_\alpha \in D_X \otimes \overline{\Omega}_\alpha \otimes V_\alpha$.

Proposition 6.1.

- (i) The maps (6.2) determine a G -equivariant structure on the free $\mathcal{D}_X$ -module $\mathcal{R}^0 E\mathcal{V}$.
- (ii) The subsheaf $\mathcal{R}^0 E\mathcal{V}'$ is invariant with respect to the maps $L_{\mathcal{R}^0 E\mathcal{V}}(g)$.

- (iii) Denote by $L_{EV}(g)$ the maps, induced by the action of the maps $L_{\mathcal{R}^0 EV}(g)$ in the quotient space EV . Then the maps $L_{EV}(g) : EV \rightarrow EV^g$ determine a G -equivariant structure on the quiver $\mathcal{D}_X$ -module EV .

□

By Proposition 6.1, we have a functor $E_G : Qui_G \rightarrow \mathcal{M}_{X,G}$ from the category of equivariant quivers to the category of equivariant $\mathcal{D}_X$ -modules.

For a nonnegative n , let $\mathcal{V}_{\Gamma^n, G} = \{V_\alpha, A_{\alpha, \beta}, A_\alpha^\beta, L_V(g)\} \in Qui_{\Gamma, G}$ be a level n G -equivariant quiver. Denote by $\mathcal{V}_{\Gamma^n} = \mathcal{N}^n(\mathcal{V}_{\Gamma^n, G})$ the same quiver with no group action. We equip the sheaf $E^n \mathcal{V}_{\Gamma^n}$ of $\mathcal{D}_{X^n}$ -modules with a structure of a G -equivariant sheaf of $\mathcal{D}_{X^n}$ -modules as follows.

Let $\mathcal{F}_n = \{f_\alpha \mid \alpha \in I_{n+1}(\Gamma)\}$ be a level n vertex framing and $U_{\mathcal{F}_n}$ the corresponding open subset of X^n , see Section 2.3. The $\mathcal{D}_{X^n}$ -module $E^n \mathcal{V}_{\Gamma^n}$ is defined as the quotient of the free sheaf $\mathcal{R}^0 E^n \mathcal{V}_{\Gamma^n} = \mathcal{D}_{X^n} \otimes_{\mathbb{C}} (\oplus_{\alpha \in I(\Gamma^n)} \overline{\Omega}_\alpha \otimes V_\alpha)$ of $\mathcal{D}_{X^n}$ -modules by its subsheaf $\mathcal{R}' E^n \mathcal{V}_{\Gamma^n}$ of coherent $\mathcal{D}_{X^n}$ -modules, where the sections $\Gamma(U_{\mathcal{F}_n}, \mathcal{R}' E^n \mathcal{V}_{\Gamma^n})$ of the sheaf $\mathcal{R}' E^n \mathcal{V}_{\Gamma^n}$ over $U_{\mathcal{F}_n}$ are given by formulas (4.33)–(4.35).

For $g \in G$, define a morphism $L_{\mathcal{R}^0 E^n \mathcal{V}_{\Gamma^n}}(g) : \mathcal{R}^0 E^n \mathcal{V}_{\Gamma^n} \rightarrow (\mathcal{R}^0 E^n \mathcal{V}_{\Gamma^n})^g$ by the formula

$$(6.3) \quad L_{\{\mathcal{R}^0 E^n \mathcal{V}_{\Gamma^n}, X^n\}}(g)(d \otimes \overline{\omega}_\alpha \otimes v_\alpha) = L_{D_{X^n}}(g)(d) \otimes L_{\Omega_X^*}(g)(\overline{\omega}_\alpha) \otimes L_{V_\alpha}(g)(v_\alpha)$$

for any $\alpha \in I(\Gamma^n)$, and $d \otimes \overline{\omega}_\alpha \otimes v_\alpha \in D_{X^n} \otimes \overline{\Omega}_\alpha \otimes V_\alpha$.

The transformation $L_X(g)$ transforms a level n vertex framing $\mathcal{F}_n$ to the level n vertex framing $\mathcal{F}_n^g = \{f_\alpha^g\}$, where $f_\alpha^g(x) = f_{g(\alpha)}(g^{-1}x)$. The transformation $L_X(g)$ maps the subset $U_{\mathcal{F}_n}$ to the subset $U_{\mathcal{F}_n^g}$. Formula (6.3) induces the maps

$$(6.4) \quad L_{\{\mathcal{R}^0 E^n \mathcal{V}_{\Gamma^n}, U_{\mathcal{F}_n}\}}(g) : \Gamma(U_{\mathcal{F}_n}, \mathcal{R}^0 E^n \mathcal{V}_{\Gamma^n}) \rightarrow \Gamma(U_{\mathcal{F}_n^g}, \mathcal{R}^0 E^n \mathcal{V}_{\Gamma^n}) .$$

Example.

Let $\mathcal{V}_{\Gamma^0, G} = \{V, A^j, L_V(g)\} \in Qui_{\Gamma^0, G}$ be an equivariant quiver of level 0. It means, in particular, that linear maps $A^j : V \rightarrow V$, as well as linear maps $A_g^j = L_V(g)A^j L_V(g)^{-1}$ for each $\alpha \in I_2(\Gamma)$, $j \succ \alpha$, satisfy the relations

$$[A^j, \sum_{i, i \succ \alpha} A^i] = 0, \quad [A_g^j, \sum_{i, i \succ \alpha} A_g^i] = 0 .$$

The associated equivariant $\mathcal{D}_{X^0}$ -module $E_G^0 \mathcal{V}_{\Gamma^0, G}$ is generated by the vectors $\omega_\emptyset \otimes v$, where $\omega_\emptyset = dz_1 \wedge \dots \wedge dz_N$ and $v \in V$, subject to the relations (4.37). The group action is defined by the relation

$$L_{E_G^0 \mathcal{V}_{\Gamma^0, G}}(g)(\omega \otimes v) = L_{\Omega_{X^0}^*}(g)(\omega) \otimes L_V(g)(v) .$$

This module is isomorphic to the $\mathcal{D}_{X^0}$ -module, associated to the flat connection (4.38) in the trivial bundle on X^0 with fiber V . The group G acts in the space of the global sections of the trivial bundle L by the relation

$$(6.5) \quad L_L(g)(f \otimes v) = L_{\mathcal{O}_{X^0}}(g)(f) \otimes \det_G(g) \cdot L_V(g)(v) .$$

Here $f \in \mathcal{O}_{X^0}$, $v \in V$ and $\det_G$ is the one-dimensional representation of the group G in the space $\overline{\Omega}_\emptyset$.

Proposition 6.2.

- (i) The maps (6.4) determine an equivariant structure on the sheaf $\mathcal{R}^0 E^n \mathcal{V}_{\Gamma^n}$ of $\mathcal{D}_{X^n}$ -modules.
- (ii) For $g \in G$, the map $L_{\{\mathcal{R}^0 E^n \mathcal{V}_{\Gamma^n}, U_{\mathcal{F}_n}\}}(g)$ maps the subspace $\Gamma(U_{\mathcal{F}_n}, \mathcal{R}' E^n \mathcal{V}_{\Gamma^n}) \subset \Gamma(U_{\mathcal{F}_n}, \mathcal{R}^0 E^n \mathcal{V}_{\Gamma^n})$ to the subspace $\Gamma(U_{\mathcal{F}_n}^g, \mathcal{R}' E^n \mathcal{V}_{\Gamma^n}) \subset \Gamma(U_{\mathcal{F}_n}^g, \mathcal{R}^0 E^n \mathcal{V}_{\Gamma^n})$.
- (iii) Denote by $L_{E^n \mathcal{V}_{\Gamma^n}}(g)$ the maps, induced by the action of $L_{\mathcal{R}^0 E^n \mathcal{V}_{\Gamma^n}}(g)$ on the quotient sheaf $E^n \mathcal{V}_{\Gamma^n}$. Then the maps $L_{E^n \mathcal{V}_{\Gamma^n}}(g)$ determine an equivariant structure on $E^n \mathcal{V}_{\Gamma^n}$.

□

By Proposition 6.2, we have a functor $E_G^n : \text{Qui}_{\Gamma^n, G} \rightarrow \mathcal{M}_{X^n, G}$ from the category of level n equivariant quivers to the category of equivariant sheaves of $\mathcal{D}_{X^n}$ -modules.

Since the map $j_{l,k} : X^k \rightarrow X^l$ is an embedding of G -invariant open subsets of X , the functor $j_{l,k}^*$ of inverse image of $\mathcal{D}$ -modules admits the canonical extension to the functor $j_{l,k,G}^* : \mathcal{M}_{X^l, G} \rightarrow \mathcal{M}_{X^k, G}$ of inverse image of equivariant $\mathcal{D}$ -modules. We define the functors $j_{l,k,*}, G : \mathcal{M}_{X^k, G} \rightarrow \mathcal{M}_{X^l, G}$ and $j_{l,k,*}, G : \mathcal{M}_{X^l, G} \rightarrow \mathcal{M}_{X^k, G}$ as right and left adjoint respectively to the functor $j_{l,k,G}^*$. We have forgetful functors $\mathcal{N}^n : \mathcal{M}_{X^n, G} \rightarrow \mathcal{M}_{X^n}$ and the functors of direct and inverse image of equivariant $\mathcal{D}$ -modules commute with the forgetful functors.

Equivariant versions of Theorems 4.5 and 4.6 are valid. For any $k < l$ and an equivariant level l quiver $\mathcal{V}_{G^k, G}$, we have an isomorphism

$$E_G^k(j_{l,k,G}^*(\mathcal{V}_{G^k, G})) \simeq j_{l,k,G}^*(E_G^l(\mathcal{V}_{G^k, G})) .$$

For any $k < l$ and an equivariant level k quiver $\mathcal{V}_{G^k, G}$, such that the quiver direct image $\mathbb{J}_{l,k,*}(\mathcal{N}^k(\mathcal{V}_{G^k, G}))$ is strongly non-resonant, we have an isomorphism

$$j_{l,k,*}^G(E_G^k(\mathcal{V}_{G^k, G})) \simeq E_G^l(\mathbb{J}_{l,k,*}, G(\mathcal{V}_{G^k, G})) .$$

Let $\mathcal{V}_G = \{V_\alpha, A_{\alpha, \beta}, L_{\mathcal{V}}(g)\} \in \text{Qui}_{\Gamma, G}$ be an equivariant quiver and $\mathcal{V} = \mathcal{N}^N(\mathcal{V}_G)$. Define an equivariant structure on the free $\mathcal{D}_X$ -module $\mathcal{R}E\mathcal{V} = \bigoplus_{r=0}^N \mathcal{R}^r E\mathcal{V}$, where

$$\mathcal{R}^r E\mathcal{V} = \bigoplus_{\alpha \in I(\Gamma)} \mathcal{D}_X \otimes \bigwedge^{-r} \mathcal{T}_\alpha \otimes \overline{\Omega}_\alpha \otimes V_\alpha ,$$

see (4.44). We define the map $L_{\mathcal{R}^r E\mathcal{V}}(g) : \mathcal{R}^r E\mathcal{V} \rightarrow (\mathcal{R}^r E\mathcal{V})^g$ by the formula

$$\begin{aligned} L_{\{\mathcal{R}^r E\mathcal{V}, X\}}(g)(d \otimes \overline{\tau} \otimes \overline{\omega}_\alpha \otimes v_\alpha) &= \\ &= L_{D_X}(g)(d) \otimes \bigwedge^{-r} L_{D_X}(g)(\overline{\tau}) \otimes L_{\Omega_X^*}(g)(\overline{\omega}_\alpha) \otimes L_{V_\alpha}(g)(v_\alpha) , \end{aligned}$$

for $\alpha \in I(\Gamma^n)$, and $d \otimes \overline{\tau} \otimes \overline{\omega}_\alpha \otimes v_\alpha \in D_X \otimes \bigwedge^{-r} \mathcal{T}_\alpha \otimes \overline{\Omega}_\alpha \otimes V_\alpha$.

Proposition 6.3.

- (i) The maps $L_{\mathcal{R}^r E\mathcal{V}}(g)$ determine an equivariant structure on the $\mathcal{D}_X$ -module $\mathcal{R}^r E\mathcal{V}$ and commute with the differentials d_0 and d_1 defined in (4.47),

$$d_i L_{\mathcal{R}^r E\mathcal{V}}(g) = L_{\mathcal{R}^{r+1} E\mathcal{V}}(g) d_i , \quad i = 1, 2 .$$

- (ii) The maps $L_{\mathcal{R}^r E\mathcal{V}}(g)$ and $L_{E\mathcal{V}}(g)$, determine an equivariant structure on the complexes $\mathcal{R}E\mathcal{V}$ and $\overline{\mathcal{R}}E\mathcal{V}$ of sheaves of $\mathcal{D}_X$ -modules.

□

Denote by $\mathcal{REV}_G$ and $\overline{\mathcal{REV}}_G$ the complexes $\mathcal{REV}$ and $\overline{\mathcal{REV}}$ equipped with the equivariant structures described above.

Proposition 6.3 implies that for any equivariant quiver $\mathcal{V}_G$, the complex $\overline{\mathcal{REV}}_G$ presents a free resolution of the equivariant $\mathcal{D}_X$ -module $E_G(\mathcal{V}_G)$ in the category of equivariant $\mathcal{D}_X$ -modules.

In particular, equivariant counterparts of Theorem 4.14 and Corollary 4.15 are valid. Namely, for any level zero equivariant quiver $\mathcal{V}_{G^0, G} = \{V, A^i, L_V(g)\}$ with non-resonant spectrum the following equivariant $\mathcal{D}_X$ -modules are isomorphic :

$$\begin{aligned} j_{0,! , G}(E_G^0(\mathcal{V}_{\Gamma^0, G})) &\approx E_G(\mathbb{J}_{0,! , G}(\mathcal{V}_{\Gamma^0, G})) , \\ j_{0,! *, G}(E_G^0(\mathcal{V}_{\Gamma^0, G})) &\approx E_G(\mathbb{J}_{0,! *, G}(\mathcal{V}_{\Gamma^0, G})) . \end{aligned}$$

6.4. Equivariant quiver perverse sheaves. Let X be a complex algebraic variety equipped with an algebraic action of a finite group G . Then the sheaf Ω_X^{an} of top exterior analytic forms on X is the right equivariant $\mathcal{D}_X$ -module with respect to the action (5.3) and equivariant structure (6.1). For any $M \in \mathcal{M}_{X, G}$ the diagonal action of G on sections of the sheaf $\Omega_X^{an} \otimes_{\mathcal{D}_X^{an}} (M)^{an}$ equips this sheaf with an equivariant structure. The assignment $M \mapsto \Omega_X^{an} \otimes_{\mathcal{D}_X^{an}} (M)^{an}$ defines a functor from the category $\mathcal{M}_{Y, G}$ to the category $Sh(X^{an}, G, \mathbb{C})$. The left derived functor to this functor is called the equivariant de Rham functor $DR_G : D^b(\mathcal{M}_{X, G}) \rightarrow D^b(X^{an}, G, \mathbb{C})$.

In the remaining part of this subsection, X is the affine space $\mathbb{C}^N$, considered as an analytic variety equipped with an affine action of a finite group G .

Let $\mathcal{V}_G$ be an equivariant quiver of Γ . The diagonal action of G equips the complex

$$\mathcal{QV} = \Omega_X^{an} \otimes_{\mathcal{D}_X^{an}} \mathcal{REV}^{an}$$

with a structure of an equivariant complex of sheaves over X^{an} . We describe this equivariant structure for global sections of the sheaf $\mathcal{QV}$. The graded component $\mathcal{QV}^r$ of the complex $\mathcal{QV}$ was described in (5.10) as

$$\mathcal{QV}^r = \bigoplus_{\alpha \in I(\Gamma)} \Omega_X^{an} \otimes \left(\bigwedge^{-r} \mathcal{T}_\alpha \otimes \overline{\Omega}_\alpha \otimes V_\alpha \right)$$

and the equivariant structure is determined by the maps

$$\begin{aligned} (6.6) \quad L_{\{\mathcal{QV}^r, X\}}(g)(\omega \otimes \overline{\tau} \otimes \overline{\omega}_\alpha \otimes v_\alpha) &= \\ &= L_{\{\Omega_X, X\}}(g)(\omega) \otimes \bigwedge^{-r} L_{D_X}(g)(\overline{\tau}) \otimes L_{\Omega_X^*}(g)(\overline{\omega}_\alpha) \otimes L_{V_\alpha}(g)(v_\alpha) , \end{aligned}$$

for any $\alpha \in I(\Gamma^n)$, and $\omega \otimes \overline{\tau} \otimes \overline{\omega}_\alpha \otimes v_\alpha \in \Gamma(X, \Omega_X) \otimes \bigwedge^{-r} \mathcal{T}_\alpha \otimes \overline{\Omega}_\alpha \otimes V_\alpha$. We have

Theorem 6.4. *Let $\mathcal{V}_G$ be an equivariant quiver. Then the equivariant de Rham complex of the equivariant quiver $\mathcal{D}_X$ -module $E_G \mathcal{V}$ is isomorphic in $D^b(X^{an}, G, \mathbb{C})$ to the perverse sheaf $\mathcal{QV}$ equipped with the equivariant structure (6.6).* □

Denote by Y the quotient space $Y = X/G$ and by $q : X \rightarrow Y$ the quotient map. We regard Y as a space with the trivial action of G . We also use the notation Y^0 for the quotient space $Y^0 = X^0/G$ with the quotient map $q_0 : X^0 \rightarrow Y^0$. The open embedding $Y^0 \rightarrow Y$ is denoted by $j_0 : Y^0 \rightarrow Y$.

Let $\mathcal{F} \in Sh(X, G, \mathbb{C})$ be a G -equivariant sheaf of complex vector spaces over X . The direct image $q_* \mathcal{F}$ of the sheaf $\mathcal{F}$ inherits from $\mathcal{F}$ the G -equivariant structure. We regard the sheaf $q_* \mathcal{F}$ as an element of $Sh(Y, G, \mathbb{C})$.

Denote by $q_*^G \mathcal{F} \in Sh(Y, \mathbb{C})$ the sheaf of G -invariant sections of $q_* \mathcal{F}$.

The functor $q_*^G: Sh(X, G, \mathbb{C}) \rightarrow Sh(Y, \mathbb{C})$ is the composition of two exact functors: direct image functor q_* and the functor of invariants of a finite group G . It sends [BL] equivariant perverse sheaves on X to perverse sheaves on Y . Thus we have

Corollary 6.5. *Let $\mathcal{V}_G$ be an equivariant quiver. Then the complex of sheaves $q_*^G \mathcal{QV}$ is a perverse sheaf on the quotient space Y .*

Denote by $\overline{\mathcal{QV}}^G$ the complex $\Gamma^G(X, \mathcal{QV}) = \Gamma(Y, q_*^G \mathcal{QV})$ of G -invariant global sections of the equivariant complex of sheaves $\mathcal{QV}$. Denote by $\det_G$ the one-dimensional representation of the group G in the space $\overline{\Omega}_0$ of top exterior analytic forms on X , invariant with respect to affine translations.

We have the following equivariant version of Theorem 5.6.

Theorem 6.6. *Let $\mathcal{C}$ be a central arrangement of hyperplanes in $\mathbb{C}^N$. Let G be a finite group of linear transformations of $\mathbb{C}^N$ preserving $\mathcal{C}$. Let $\mathcal{V}_G$ be an equivariant quiver of the graph Γ of the arrangement, such that all linear maps $A_{\alpha, \beta}$ of the quiver are close to zero. Then the complex $\overline{\mathcal{QV}}^G$ is quasi-isomorphic to the subcomplex of G -invariants $((C_+(\mathcal{V}) \otimes \det_G)[N])^G$ of the complex $(C_+(\mathcal{V}) \otimes \det_G)[N]$:*

$$\overline{\mathcal{QV}}^G \stackrel{quis}{\simeq} ((C_+(\mathcal{V}) \otimes \det_G)[N])^G .$$

The theorem is proved in Section 7.

6.5. Cohomology of $\mathbb{C}^N$ with coefficients in equivariant quiver perverse sheaves.

The category $Sh(X, G, \mathbb{C})$ of equivariant sheaves has enough injectives [Gr]. Let $\Gamma^G(X, \cdot) : Sh(X, G, \mathbb{C}) \rightarrow Vect_{\mathbb{C}}$ be the functor of invariant global sections. Following [Gr], define $H^k(X, G, \cdot)$ to be the k -th right derived functor of the functor $\Gamma^G(X, \cdot)$, such that $H^k(X, G, \mathcal{F}) = R^k \Gamma^G(X, \mathcal{F})$ for any complex of equivariant sheaves $\mathcal{F} \in D^b(Sh(X, G, \mathbb{C}))$. Since G is a finite group, for any $\mathcal{F} \in D^b(Sh(X, G, \mathbb{C}))$ the graded space $H^\bullet(X, G, \mathcal{F})$ coincides with the equivariant cohomology group of $\mathcal{F}$ [BL].

Since the functor $q_*^G : Sh(X, G, \mathbb{C}) \rightarrow Sh(Y, \mathbb{C})$ is the composition of two exact functors, we have an equality [Gr] :

$$H^k(X, G, \mathcal{F}) = H^k(Y, q_*^G \mathcal{F}) .$$

The same statement is valid for the space X^0 : $H^k(X^0, G, \mathcal{F}) = H^k(Y^0, (q_0)_*^G \mathcal{F})$.

The affine space X satisfies the equivariant Stein condition, that is, for a finite group G

$$(6.7) \quad H^k(X, G, \mathcal{O}^{an}) = 0, \quad \text{for all } k \neq 0.$$

Statement (6.7) can be proved as follows. In the non-equivariant case to show that $H^k(X, \mathcal{O}^{an})$ is trivial we replace the sheaf $\mathcal{O}^{an}$ (modulo constants) by its Dolbeault resolution $(\Omega^{0,q}, \partial_{\bar{z}})$. Then the $\partial_{\bar{z}}$ -Poincare lemma says that any $\partial_{\bar{z}}$ -closed $(0, q)$ -form is a $\partial_{\bar{z}}$ -coboundary. In the equivariant case we can use the same Dolbeault resolution, since it

consists of equivariant soft sheaves. Statement (6.7) requires that an invariant $\partial_{\bar{z}}$ -closed $(0, q)$ -form ω has to be the $\partial_{\bar{z}}$ -coboundary of an invariant form. By the $\partial_{\bar{z}}$ -Poincare lemma, we may conclude that w is the $\partial_{\bar{z}}$ -coboundary of a possibly non-invariant form y . Then the non-invariance of y is corrected by averaging over G .

Corollary 6.7. *Let $\mathcal{C}$ be a central arrangement of hyperplanes in $\mathbb{C}^N$. Let G be a finite group of linear transformations of $\mathbb{C}^N$ preserving $\mathcal{C}$. Let $\mathcal{V}_G$ be an equivariant quiver with all linear maps $A_{\alpha, \beta}$ close to zero. Then for any $k \in \mathbb{Z}$*

$$H^k(X, G, DR(E\mathcal{V})) \simeq H^k(X, G, \mathcal{QV}) \simeq H^{k+N}((C_+(\mathcal{V}) \otimes \det_G)^G) .$$

□

For an equivariant local system $\mathcal{L}$ on X^0 introduce equivariant intersection cohomology groups $IH_G^k(X, G, \mathcal{L})$ as the equivariant cohomology groups $H^{k-N}(X, G, (j_0)_!^*, G\mathcal{L}[N])$ of the MacPherson extension of the equivariant perverse sheaf $\mathcal{L}[N]$,

$$IH^k(X, G, \mathcal{L}) \stackrel{\text{def}}{=} H^{k-N}(X, G, (j_0)_!^*, G\mathcal{L}[N]) .$$

The functors q_* , $(q^0)_*$, q_*^G , and $(q^0)_*^G$ transform perverse sheaves to perverse sheaves [BBD], so that we have an equality

$$IH^k(X, G, \mathcal{L}) = IH^k(Y, (q^0)_*^G \mathcal{L}) ,$$

where $IH^k(Y, \mathcal{L}')$ means the same as in Section 5.5,

$$IH^k(Y, \mathcal{L}') \stackrel{\text{def}}{=} H^{k-N}(Y, (j_0)_!^* \mathcal{L}'[N]) .$$

for any local system $\mathcal{L}'$ over Y^0 .

Let $\mathcal{LV}$ be the local system of flat sections of the connection (4.38) equipped with the group action (6.5).

Corollary 6.8. *Let $\mathcal{V}_G = \{V, A^i, L_V(g)\}$ be an equivariant level zero quiver of a central arrangement $\mathcal{C}$. Assume that all operators A^i are close to zero. Then for any $k \in \mathbb{Z}$,*

$$\begin{aligned} H^k(Y^0, (q_0)_*^G \mathcal{LV}) &\simeq H^k(X^0, G, \mathcal{LV}) \simeq H^k((C_+(\mathbb{J}_{0,*} \mathcal{V}) \otimes \det_G)^G) , \\ IH^k(Y, (q_0)_*^G \mathcal{LV}) &\simeq IH^k(X, G, \mathcal{LV}) \simeq H^k(C_+(\mathbb{J}_{0,*} \mathcal{V}) \otimes \det_G)^G . \end{aligned}$$

6.6. Application to discriminantal arrangements. In this section we consider the discriminantal arrangements and local systems appearing in hypergeometric solutions of the KZ equations in [SV, V]. We calculate the equivariant intersection cohomology groups of the discriminantal arrangements in terms of representation theory.

In this section we follow notations of [V], chapter 11.

We fix the following data:

- (a) a finite-dimensional complex space $\mathfrak{h}$;
- (b) a non-degenerate bilinear form $(,)$ on $\mathfrak{h}$;
- (c) linearly independent covectors ('simple roots') $\alpha_1, \dots, \alpha_r \in \mathfrak{h}^*$.

Denote by $b : \mathfrak{h} \rightarrow \mathfrak{h}^*$ the isomorphism induced by $(,)$. Transform the form $(,)$ to $\mathfrak{h}^*$, using b . Set $b_{i,j} = (\alpha_i, \alpha_j)$, $h_i = b^{-1}(\alpha_i) \in \mathfrak{h}$.

Denote by $\mathfrak{g}$ the Lie algebra, generated by e_i, f_i , and $\mathfrak{h}$, subject to the relations

$$\begin{aligned} [h, e_i] &= \langle h, \alpha_i \rangle e_i, & [h, f_i] &= -\langle h, \alpha_i \rangle f_i, \\ [e_i, f_j] &= \delta_{i,j} h_i, & [h, h'] &= 0, \end{aligned}$$

for all $i, j = 1, \dots, r$ and $h, h' \in \mathfrak{h}$. Denote by $\mathfrak{n}_+$ and $\mathfrak{n}_-$ the subalgebras of $\mathfrak{g}$, generated by e_i and f_i , respectively.

There is a unique symmetric bilinear form $S(\cdot, \cdot)$ on $\mathfrak{g}$, satisfying the conditions

- (i) the subspaces $\mathfrak{n}_+$, $\mathfrak{h}$, and $\mathfrak{n}_-$ are mutually orthogonal with respect to S ;
- (ii) the restriction of S to $\mathfrak{h}$ coincides with $(\cdot, \cdot)$,
- (iii) $S(e_i, e_j) = S(f_i, f_j) = \delta_{i,j}$; $S([f_i, x], y) = S(x, [e_i, y])$.

Set $\bar{\mathfrak{g}} = \mathfrak{g}/\text{Ker } S$, $\bar{\mathfrak{n}}_{\pm} = \mathfrak{n}_{\pm}/(\mathfrak{n}_{\pm} \cap \text{Ker } S)$. These are Lie algebras. If $b_{i,i} \neq 0$ for all i , then $\bar{\mathfrak{g}}$ is the Kac-Moody algebra associated with the matrix $a_{i,j} = 2b_{i,j}/b_{i,i}$.

For $\Lambda \in \mathfrak{h}^*$, we denote by U_{Λ} the irreducible $\bar{\mathfrak{g}}$ -module with highest weight $\Lambda \in \mathfrak{h}^*$.

For a highest weight $\bar{\mathfrak{g}}$ -module U , we denote by $C_{\bullet}(\bar{\mathfrak{n}}_-, U)$ the standard chain complex of $\bar{\mathfrak{n}}_-$ with coefficients in U . The chain complex $C_{\bullet}(\bar{\mathfrak{n}}_-, U)$ has the canonical structure of an $\mathfrak{h}$ -module. For $\mu \in \mathfrak{h}^*$, we denote by $C_{\bullet}(\bar{\mathfrak{n}}_-, U)_{\mu}$ the subcomplex of chains of weight μ .

Let $\mathcal{C}_{1,N}$ be the arrangement in $X = \mathbb{C}^N$ of hyperplanes

$$\begin{aligned} H_i &: t_i = 0, & 1 \leq i \leq N; \\ H_{i,j} &: t_i - t_j = 0, & 1 \leq i < j \leq N. \end{aligned}$$

Fix an r -tuple $\lambda = (k_1, \dots, k_r) \in \mathbb{Z}_{\geq 0}^r$, $k_1 + \dots + k_r = N$, and a map $\pi_{\lambda} : \{1, \dots, N\} \rightarrow \{1, \dots, r\}$, such that $\#\pi_{\lambda}^{-1}(i) = k_i$. We denote by Σ_{λ} the subgroup of the symmetric group Σ_N , which consists of permutations σ of $\{1, \dots, N\}$, satisfying the condition $\pi_{\lambda}(\sigma(i)) = \pi_{\lambda}(i)$ for all i . We have an isomorphism $\Sigma_{\lambda} \approx \Sigma_{k_1} \times \dots \times \Sigma_{k_r}$. The group Σ_{λ} , as well as the symmetric group Σ_N , acts on X by permutations of coordinates. This action preserves $\mathcal{C}_{1,N}$.

For $\Lambda \in \mathfrak{h}^*$ and $\kappa \in \mathbb{C}$, we denote by $a_{\Lambda, \pi_{\lambda}, \kappa}$ the collection of exponents $a_{\Lambda, \pi_{\lambda}, \kappa} : \mathcal{C}_{1,N} \rightarrow \mathbb{C}$, where

$$a_{\Lambda, \pi_{\lambda}, \kappa}(H_{i,j}) = (\alpha_{\pi_{\lambda}(i)}, \alpha_{\pi_{\lambda}(j)})/\kappa, \quad a_{\Lambda, \pi_{\lambda}, \kappa}(H_i) = -(\alpha_{\pi_{\lambda}(i)}, \Lambda)/\kappa.$$

Let $(\mathcal{A}(\mathcal{C}_{1,N}), d_{a_{\Lambda, \pi_{\lambda}, \kappa}})$ be the Aomoto complex of $\mathcal{C}_{1,N}$, see Section 2.5. Let $(\bar{\mathcal{F}}_{a_{\Lambda, \pi_{\lambda}, \kappa}}, d_{a_{\Lambda, \pi_{\lambda}, \kappa}})$ be its subcomplex of flag forms.

The action of Σ_{λ} on $\mathcal{C}_{1,N}$ induces the action of Σ_{λ} on $(\mathcal{A}(\mathcal{C}_{1,N}), d_{a_{\Lambda, \pi_{\lambda}, \kappa}})$. This action preserves $(\bar{\mathcal{F}}_{a_{\Lambda, \pi_{\lambda}, \kappa}}, d_{a_{\Lambda, \pi_{\lambda}, \kappa}})$.

The following isomorphism of complexes was established in [SV], Corollary 6.13.

Theorem 6.9 ([SV]). *We have*

$$(6.8) \quad C_{\bullet}(\bar{\mathfrak{n}}, U_{\Lambda})_{\Lambda - \bar{\lambda}} \approx \left(\bar{\mathcal{F}}_{a_{\Lambda, \pi_{\lambda}, \kappa}}^{N-\bullet}(\mathcal{C}_{1,N}) \otimes \text{sgn} \right)^{\Sigma_{\lambda}},$$

Here $\bar{\lambda} = k_1 \alpha_1 + \dots + k_r \alpha_r$ and sgn is the restriction to Σ_{λ} of the one-dimensional sign representation of Σ_N .

We rewrite (6.8) in a quiver language. The collection of exponents $a_{\Lambda, \pi_\lambda, \kappa}$ determines the level zero equivariant quiver $\mathcal{V}_{\Sigma_\lambda}(\Lambda, \pi_\lambda, \kappa)$ of the arrangement $\mathcal{C}_{1,N}$ by the rule $\mathcal{V}_{\Sigma_\lambda}(\Lambda, \pi_\lambda, \kappa) = \{V, A^l, L_V(g)\}$. Here $V = \mathbb{C}$. The operator $A^l : \mathbb{C} \rightarrow \mathbb{C}$ is the operator of multiplication by the exponent $a_{\Lambda, \pi_\lambda, \kappa}(H(l))$ of the corresponding hyperplane. All operators $L_V(g)$ are equal to Id_V .

In these notations the flag complex $\overline{\mathcal{F}}_{a_{\Lambda, \pi_\lambda, \kappa}}^\bullet(\mathcal{C}_{1,N})$ coincides with the complex of the direct image $\mathbb{J}_{0,!} \mathcal{V}(\Lambda, \pi_\lambda, \kappa)$ of the quiver $\mathcal{V}(\Lambda, \pi_\lambda, \kappa) = \mathcal{N}^0 \mathcal{V}_{\Sigma_\lambda}(\Lambda, \pi_\lambda, \kappa)$,

$$\overline{\mathcal{F}}_{a_{\Lambda, \pi_\lambda, \kappa}}^\bullet(\mathcal{C}_{1,N}) = C_+(\mathbb{J}_{0,!} \mathcal{V}(\Lambda, \pi_\lambda, \kappa)) .$$

Statement (6.8) is equivalent to the statement

$$(6.9) \quad C_\bullet(\overline{\mathbf{n}}, L_\Lambda)_{\Lambda - \overline{\lambda}} \approx (C_+^{N-\bullet}(\mathbb{J}_{0,!} \mathcal{V}_{\Sigma_\lambda}(\Lambda, \pi_\lambda, \kappa)) \otimes \text{sgn})^{\Sigma_\lambda} .$$

Let $\mathcal{L}_{\Lambda, \pi_\lambda, \kappa}$ be the local system on X^0 , associated to the connection in the trivial line bundle L over X^0 with the connection-form

$$\sum_{1 \leq i < j \leq N} \frac{(\alpha_{\pi_\lambda(i)}, \alpha_{\pi_\lambda(j)})}{\kappa} d \log(t_i - t_j) - \sum_{1 \leq i \leq N} \frac{(\alpha_{\pi_\lambda(i)}, \Lambda_j)}{\kappa} d \log t_i .$$

We identify sections of L with scalar functions. We define the action of Σ_λ on sections by the formula

$$L_L(g)(f(x)) = \text{sgn}(g) \cdot f(L_X(g)^{-1}(x)) .$$

In notations of Section 6.5, we denote by $\overline{\mathcal{L}}_{\Lambda, \overline{\lambda}, \kappa}$ the local system on Y^0 given by the formula

$$\overline{\mathcal{L}}_{\Lambda, \overline{\lambda}, \kappa} = (q_0)_\bullet^{\Sigma_\lambda} \mathcal{L}_{\Lambda, \pi_\lambda, \kappa} .$$

Combining (6.9) and Corollary 6.8, we get

Corollary 6.10. *Let X^0 be the complement to $\mathcal{C}_{1,N}$ in $\mathbb{C}^N$. Then for $|\kappa| \gg 1$ and $k \in \mathbb{Z}$, we have*

$$IH^k(Y, \overline{\mathcal{L}}_{\Lambda, \overline{\lambda}, \kappa}) = IH^k(X, \Sigma_\lambda, \mathcal{L}_{\Lambda, \pi_\lambda, \kappa}) \simeq H_{N-k}(\overline{\mathbf{n}}, L_\Lambda)_{\Lambda - \overline{\lambda}} .$$

Let $\overline{\mathfrak{g}}$ be a simple Lie algebra with Weyl group W and half sum of positive roots ρ . Let U_Λ be the irreducible finite-dimensional $\overline{\mathfrak{g}}$ -module with highest weight Λ . By the Borel–Weil–Bott theorem, we have

$$H_k(\overline{\mathbf{n}}, U_\Lambda) = \bigoplus_{w \in W, l(w)=k} \mathbb{C}_{w(\Lambda+\rho)-\rho}$$

where $\mathbb{C}_\mu$ is a one-dimensional space of weight μ and $l(w)$ is the length of w . The dominant weight $\Lambda + \rho$ is regular, that is, the equality $w(\Lambda + \rho) = \Lambda + \rho$ implies $w = id$. Hence each weight subspace of $H_k(\overline{\mathbf{n}}, U_\Lambda)$ is at most one-dimensional.

To a simple Lie algebra $\overline{\mathfrak{g}}$ and an element $\overline{\lambda} = k_1 \alpha_1 + \dots + k_r \alpha_r$, $k_i \in \mathbb{Z}_{\geq 0}$, of the root lattice we assign the vector $\lambda = (k_1, \dots, k_r)$, the number $N = k_1 + \dots + k_r$, and the arrangement $\mathcal{C}_{1,N}$ in $X = \mathbb{C}^N$, equipped with the action of the product of symmetric groups Σ_λ as at the beginning of this section. An integral dominant weight $\Lambda \in \mathbf{h}^*$ determines the Σ_λ -equivariant local system $\mathcal{L}_{\Lambda, \pi_\lambda, \kappa}$ on the complement X^0 to the arrangement $\mathcal{C}_{1,N}$

and the local system $\overline{\mathcal{L}}_{\Lambda, \overline{\lambda}, \kappa}$ on the quotient space $Y^0 = X^0/\Sigma_\lambda$, which is an open subset of the quotient space $Y = X/\Sigma_\lambda$.

Corollary 6.11. *For $|\kappa| \gg 1$ and $k \in \mathbb{Z}$, the intersection cohomology $IH^k(Y, \overline{\mathcal{L}}_{\Lambda, \overline{\lambda}, \kappa})$ is at most one-dimensional and is nonzero if and only there exists $w \in W$, such that $l(w) = N - k$ and $w(\Lambda + \rho) - \rho = \Lambda - \overline{\lambda}$.*

7. PROOFS

7.1. Proof of Propositions 3.1 and 3.2. We start with the proof of Proposition 3.1. The operators A_β^α and A_β^γ commute if $\alpha \succ \beta$ and $\beta \succ \gamma$. Indeed, using relations (b) and (c) in Section 2.1 we get

$$\begin{aligned} A_\beta^\alpha A_\beta^\gamma &= A_{\beta, \alpha} A_{\alpha, \beta} A_{\beta \gamma} A_{\gamma \beta} = - \sum_{\substack{\beta', \beta' \neq \beta \\ \alpha \succ \beta' \succ \gamma}} A_{\beta, \alpha} A_{\alpha \beta'} A_{\beta' \gamma} A_{\gamma \beta} = \\ &= - \sum_{\substack{\beta', \beta' \neq \beta \\ \alpha \succ \beta' \succ \gamma}} A_{\beta \gamma} A_{\gamma \beta'} A_{\beta' \alpha} A_{\alpha, \beta} = A_{\beta \gamma} A_{\gamma \beta} A_{\beta, \alpha} A_{\alpha, \beta} = A_\beta^\gamma A_\beta^\alpha. \end{aligned}$$

To prove (i) we have to check only that $[S_\beta, A_\beta^\alpha] = 0$ for α such that $\alpha \succ \beta$. This is equivalent to the equality $[\sum_{\substack{\alpha', \alpha' \neq \alpha, \\ \alpha' \succ \beta}} A_\beta^{\alpha'}, A_\beta^\alpha] = 0$. Consider the product $A_\beta^\alpha \cdot \sum_{\substack{\alpha_j, \alpha_j \neq \alpha, \\ \alpha_j \succ \beta}} A_\beta^{\alpha_j}$.

A nonzero contribution to the product is given only by the summands with α_j for which there exists δ_j with $\delta_j \succ \alpha_j$ and $\delta_j \succ \alpha$. We have

$$\begin{aligned} (7.1) \quad A_\beta^\alpha \sum_{\substack{\alpha_j, \alpha_j \succ \beta, \\ \alpha_j \neq \alpha}} A_\beta^{\alpha_j} &= A_{\beta, \alpha} A_{\alpha, \beta} \sum_{\delta_i, \delta_i \succ \alpha} \sum_{\substack{\alpha_j, \alpha_j \neq \alpha, \\ \delta_i \succ \alpha_j \succ \beta}} A_{\beta \alpha_j} A_{\alpha_j \beta} = \\ &= \sum_{\delta_i, \delta_i \succ \alpha} \sum_{\substack{\alpha_j, \alpha_j \neq \alpha, \\ \delta_i \succ \alpha_j \succ \beta}} A_{\beta, \alpha} A_{\alpha \delta_i} A_{\delta_i \alpha_j} A_{\alpha_j \beta} = \sum_{\delta_i, \delta_i \succ \alpha} A_{\beta, \alpha} A_{\alpha \delta_i} A_{\delta_i \alpha} A_{\alpha, \beta}. \end{aligned}$$

We can apply the same procedure to the product $\sum_{\substack{\alpha_j, \alpha_j \neq \alpha, \\ \alpha_j \succ \beta}} A_\beta^{\alpha_j} A_\beta^\alpha$ and get the same result

(7.1). This proves statement (i) of Proposition 3.1.

To prove (ii) we note that for any vertex $\delta \succ \beta$ with $l(\delta) = l(\beta) - 2$ and for any vector $e_\delta \in V_\delta$, the vector $\oplus_{\alpha, \delta \succ \alpha \succ \beta} A_{\alpha, \delta} e_\delta$ belongs to the kernel of the operator T_β due to defining relation (b) of Section 2.1. Fix a vertex α , such that $\alpha \succ \beta$, and a vector $e_\alpha \in V_\alpha$. We have

$$T_\beta(e_\alpha) = A_{\alpha, \beta} A_{\beta, \alpha}(e_\alpha) + \sum_{\substack{\alpha_j, \alpha_j \neq \alpha, \\ \alpha_j \succ \beta}} A_{\alpha_j \beta} A_{\beta, \alpha}(e_\alpha).$$

Due to relation (c) of Section 2.1, in the last sum we have a nonzero contribution only from the summands with α_j such that there exists δ_j satisfying $\delta_j \succ \alpha_j$ and $\delta_j \succ \alpha$. Thus

$$\begin{aligned} T_\beta(e_\alpha) &= A_{\alpha,\beta}A_{\beta,\alpha}(e_\alpha) - \sum_{\delta_i, \delta_i \succ \alpha} \sum_{\substack{\alpha_j, \alpha_j \neq \alpha \\ \delta_i \succ \alpha_j \succ \beta}} A_{\alpha_j \delta_i} A_{\delta_i \alpha}(e_\alpha) = \\ &= A_{\alpha,\beta}A_{\beta,\alpha}(e_\alpha) + \sum_{\delta_i, \delta_i \succ \alpha} A_{\alpha \delta_i} A_{\delta_i \alpha}(e_\alpha) - \sum_{\delta_i, \delta_i \succ \alpha} \sum_{\alpha_j, \delta_i \succ \alpha_j \succ \beta} A_{\alpha_j \delta_i} A_{\delta_i \alpha}(e_\alpha). \end{aligned}$$

The last term belongs to the kernel of T_β . So the action of T_β on e_α in the quotient space $\oplus_{\alpha: \alpha \succ \beta} V_\alpha / \text{Ker } T_\beta$ is the same as the action of $A_\alpha^\beta + T_\alpha^0$ on e_α . This proves statement (ii) of Proposition 3.1. $\square$

Let us prove statement (i) of Proposition 3.2. Since A_β^α and A_β^γ commute if $\alpha \succ \beta$ and $\beta \succ \gamma$, we have to prove that $[\tilde{S}_\beta, A_\beta^\alpha] = 0$ for any $\alpha \in I(\Gamma)$, $\beta \succ \alpha$. Fix a vertex $\alpha \in I(\Gamma)$, such that $\beta \succ \alpha$. We have $\tilde{S}_\beta = \sum_{\gamma, \beta \succ \gamma} A_\beta^\gamma$. Since the arrangement is central, for any γ , $\beta \succ \gamma$, the planes $\overline{X}_\alpha$ and $\overline{X}_\gamma$ intersect in codimension one and we can rewrite $\tilde{S}_\beta$ as

$$\tilde{S}_\beta = \sum_{\delta, \beta \succ \delta, l(\delta)=l(\beta)+2} \tilde{S}_{\beta,\delta},$$

where

$$(7.2) \quad \tilde{S}_{\beta,\delta} = \sum_{\gamma, \beta \succ \gamma \succ \delta} A_\beta^\gamma.$$

The defining relations of quivers (see the relation (v) of Section 3.2) imply that $[A_\beta^\alpha, \tilde{S}_{\beta,\delta}] = 0$ for any such δ . Thus $[A_\beta^\alpha, \tilde{S}_\beta] = 0$ and statement (i) is proved.

Let us prove statement (ii) of Proposition 3.2. Fix $\alpha, \beta \in I(\Gamma)$, such that $\alpha \succ \beta$. For any δ , $\beta \succ \delta$ we have by the second equality of Section 3.2 (iv)

$$(7.3) \quad A_{\alpha,\beta}A_\beta^\delta = (\tilde{S}_{\alpha,\delta} - A_\alpha^\beta)A_{\alpha,\beta}.$$

The summation of (7.3) over all δ , $\beta \succ \delta$ gives

$$(7.4) \quad A_{\alpha,\beta}\tilde{S}_\beta = (\tilde{S}_\alpha - A_\alpha^\beta)A_{\alpha,\beta}.$$

On the other hand, we have, analogously to (7.1),

$$\begin{aligned} A_{\alpha,\beta} \sum_{\substack{\alpha_j, \alpha_j \succ \beta, \\ \alpha_j \neq \alpha}} A_\beta^{\alpha_j} &= A_{\alpha,\beta} \sum_{\delta_i, \delta_i \succ \alpha} \sum_{\substack{\alpha_j, \alpha_j \neq \alpha, \\ \delta_i \succ \alpha_j \succ \beta}} A_{\beta \alpha_j} A_{\alpha_j \beta} = \\ &- \sum_{\delta_i, \delta_i \succ \alpha} \sum_{\substack{\alpha_j, \alpha_j \neq \alpha, \\ \delta_i \succ \alpha_j \succ \beta}} A_{\alpha \delta_i} A_{\delta_i \alpha_j} A_{\alpha_j \beta} = \sum_{\delta_i, \delta_i \succ \alpha} A_{\alpha \delta_i} A_{\delta_i \alpha} A_{\alpha, \beta}, \end{aligned}$$

that is,

$$(7.5) \quad A_{\alpha,\beta}(S_\beta - A_\beta^\alpha) = S_\alpha A_{\alpha,\beta}.$$

Combining (7.4) and (7.5), we get

$$A_{\alpha,\beta}(S_\beta + \tilde{S}_\beta - A_\beta^\alpha) = (S_\alpha + \tilde{S}_\alpha - A_\alpha^\beta)A_{\alpha,\beta},$$

which implies (3.9), since $A_{\alpha,\beta}A_{\beta}^{\alpha} = A_{\alpha}^{\beta}A_{\alpha,\beta}$ by definition of operators A_{α}^{β} and A_{β}^{α} , see (3.3). The proof of (3.9) for $\beta \succ \alpha$ is analogous. The relations (3.9) imply the centrality of the operator S . $\square$

7.2. Proof of Propositions 3.9–3.11.

First we prove Proposition 3.9.

Consider the direct image $\mathbb{J}_{0,*}\mathcal{W}_{\Gamma^0}$ of the level zero quiver $\mathcal{W}_{\Gamma^0} = \{W, B^i\}$. Consider a vertex $\alpha \in I(\Gamma)$ and a flag $F_{\alpha,\dots,\alpha_m=\alpha}$ which ends at α . Let $\beta \in I(\Gamma)$ be a vertex such that $\beta \succ \alpha$. According to (3.17) and (3.18), the vector $A_{\alpha,\beta}A_{\beta,\alpha}(F_{\alpha,\dots,\alpha_m} \otimes u)$ in $(\mathbb{J}_{0,*}\mathcal{W}_{\Gamma^0})_{\alpha}$ is equal to zero if $F_{\alpha_0,\dots,\alpha_m}^{\beta} = 0$ or is equal to

$$(-1)^{((\beta;\alpha_0,\dots,\alpha_m)+m-1)} F_{\beta_0,\dots,\beta_{m-1}\alpha_m} \otimes \sum_{\substack{i \in I_1(\Gamma), \\ i \geq \alpha_{k+1}, i \not\geq \alpha_k}} B^i(u),$$

if $F_{\alpha,\dots,\alpha_m}^{\beta} \neq 0$. Here $F_{\beta_0,\dots,\beta_{m-1}} = F_{\alpha,\dots,\alpha_m}^{\beta}$. Denote by $I_k(\alpha_0, \alpha_1, \dots, \alpha_m)$ the set of all $\beta \in I(\Gamma)$ such that $F_{\alpha,\dots,\alpha_m}^{\beta} \neq 0$ and $(\beta; \alpha_0, \dots, \alpha_m) = k$. Applying relations (2.8), we get

$$(7.6) \quad \sum_{\beta \in I_k(\alpha_0, \dots, \alpha_m)} A_{\alpha,\beta}A_{\beta,\alpha}(F_{\alpha,\dots,\alpha_m=\alpha} \otimes u) = F_{\alpha,\dots,\alpha_m} \otimes \sum_{\substack{i \in I_1(\Gamma), \\ i \geq \alpha_{k+1}, i \not\geq \alpha_k}} B^i(u).$$

Relations (7.6) imply that for any $\alpha \in I(\Gamma)$,

$$(7.7) \quad S_{\alpha} = Id_{\mathcal{F}_{\alpha}} \otimes \sum_{i \in I_1(\Gamma), i \geq \alpha} B^i,$$

as an operator in $\mathcal{F}_{\alpha} \otimes U$. Denote by $\tilde{B}^{\alpha}$ the operator $\tilde{B}^{\alpha} = \sum_{i \in I_1(\Gamma), i \geq \alpha} B^i : W \rightarrow W$. According to Proposition 3.1, the operator S_{α} commutes with any product $A_{\alpha,\beta}A_{\beta,\alpha}$, if β is adjacent to α . Applying this fact to relations (7.6) and (7.7) we get

$$[\tilde{B}^{\alpha}, \sum_{\substack{i \in I_1(\Gamma), \\ i \geq \alpha_{k+1}, i \not\geq \alpha_k}} B^i] = 0,$$

and, in particular, for $k = 0$, we get $[\tilde{B}^{\alpha}, B^{\alpha_1}] = 0$. We can repeat these arguments for any flag ending at α and we conclude that the operator $\tilde{B}^{\alpha}$ commutes with all its own summands B^i . In this situation all eigenvalues of the operator $\tilde{B}^{\alpha}$ are equal to λ_{α} . Indeed, for any eigenvalue λ of $\tilde{B}^{\alpha}$ the corresponding subspace of eigenvectors is invariant with respect to any B^i , $i \geq \alpha$. Taking the trace over this subspace of the equality $\tilde{B}^{\alpha} = \sum_{i \in I_1(\Gamma): i \geq \alpha} B^i$, we get $\lambda = \lambda_{\alpha}$. Due to (7.7) we get statement (i) of the proposition.

To prove (iii), note that due to (3.17) and (3.18) for any $\alpha, \beta \in I(\Gamma)$, such that $\alpha \succ \beta$, the operator $A_{\alpha}^{\beta} = A_{\alpha,\beta}A_{\beta,\alpha} : \mathcal{F}_{\alpha} \otimes U \rightarrow \mathcal{F}_{\alpha} \otimes U$ is equal to

$$(7.8) \quad A_{\alpha}^{\beta} = Id_{\mathcal{F}_{\alpha}} \otimes \sum_{\substack{i \in I_1(\Gamma), \\ i \geq \beta, i \not\geq \alpha}} B^i.$$

Statement (iii) follows now from (i) and (ii) and Proposition 3.1. $\square$

Consider the quiver $\mathbb{J}_{0,!}\mathcal{W}_{\Gamma^0} = \tau_N^{-1}\mathbb{J}_{0,*}\tau_0\mathcal{W}_{\Gamma^0}$. The quiver $\tau_0\mathcal{W}_{\Gamma^0} = \{U, C^i\}$ consists of the vector space $U = W^*$ and operators $C^i = (B^i)^*$, respectively dual to B^i . Each of C^i has a single eigenvalue λ_i . By Proposition 3.9, the operators $S_\alpha, A_\alpha^\beta, T_\alpha, T'_\alpha$, attached to the quiver $\mathbb{J}_{0,*}\tau_0\mathcal{W}_{\Gamma^0}$, have eigenvalues, prescribed by that proposition. The operators $S_\alpha, A_\alpha^\beta, T_\alpha, T'_\alpha$, attached to the quiver $\tau_N^{-1}\mathbb{J}_{0,*}\tau_0\mathcal{W}_{\Gamma^0}$, are dual to the corresponding operators for the quiver $\mathbb{J}_{0,*}\tau_0\mathcal{W}_{\Gamma^0}$ and thus have the same eigenvalues. This proves Proposition 3.10. $\square$

In order to prove Proposition 3.11, it is sufficient to prove it for the quiver $\mathbb{J}_{0,*}\mathcal{W}_{\Gamma^0}$ and then apply the duality arguments.

For $\alpha \in I(\Gamma)$, formula (7.8) implies that

$$\tilde{S}_\alpha = Id_{\mathcal{F}_\alpha} \otimes \sum B^i,$$

where the sum is taken over all $i \in I_1(\Gamma)$, such that the hyperplane H_i does not contain X_α , but its intersection with X_α is non-empty. For a central arrangement, the last condition is satisfied always. Thus

$$(7.9) \quad \tilde{S}_\alpha = Id_{\mathcal{F}_\alpha} \otimes \sum_{i \in I_1(\Gamma), i \not\geq \alpha} B^i.$$

Combining (7.9) with (7.7), we get

$$(7.10) \quad S = Id_{\mathcal{F}} \otimes \sum_{i \in I_1(\Gamma)} B^i.$$

The trace arguments as above show that the operator S has a single eigenvalue $\lambda_\infty = \sum_{j \in J(\mathcal{C})} \lambda_j$, if every operator B^i has a single eigenvalue λ_i . $\square$

7.3. Proof of Proposition 4.1. Choose a scalar product on X invariant with respect to parallel translations. Without loss of generality we assume that for any pair of adjacent strata $\overline{X}_\alpha \supset X_\beta$, the function $f_{\beta,\alpha} \in F_\beta \setminus F_\alpha$ is chosen to be orthogonal to F_α and the vector field $\xi_{\alpha,\beta} \in T_\alpha \setminus T_\beta$ belongs to T'_β . Set $M = \oplus_\alpha M_\alpha$. Due to relations (4.25) and (4.26), the space of global sections of the $\mathcal{D}_X$ -module $E\mathcal{V}$ is isomorphic to a quotient space of M . To prove (4.29), we define inductively the structure of a D_X -module on M .

Define the grading on M . It is given by a grading on each summand M_α , where we set $\deg(v_\alpha \otimes \overline{\omega}_\alpha) = 0$, $\deg \xi = \deg f = 1$ for $\xi \in T'_\alpha$ and $f \in F'_\alpha$.

We will use the increasing Bernstein filtration $BD^\bullet$ of the ring D_X , in which the subspace $BD^{\leq k} \subset D_X$ is spanned by the elements $z_{i_1} \cdots z_{i_l} \frac{\partial}{\partial z_{i_{l+1}}} \cdots \frac{\partial}{\partial z_{i_m}}$,

$$BD^{\leq k} = \mathbb{C} \left\langle z_{i_1} \cdots z_{i_l} \frac{\partial}{\partial z_{i_{l+1}}} \cdots \frac{\partial}{\partial z_{i_m}} \mid m \leq k \right\rangle.$$

Define the action of $BD^{\leq 1}$ on elements of zero degree $M^0 \subset M$. Take a stratum X_α . An element $d \in BD^{\leq 1}$ can be presented as a sum

$$(7.11) \quad d = c + d_\alpha + d'_\alpha,$$

where c is a constant, $d_\alpha \in T_\alpha + F_\alpha$, $d'_\alpha \in T'_\alpha + F'_\alpha$. In the ring D_X , we have

$$(7.12) \quad [d_1, d_2] = 0 \quad \text{if} \quad d_1, d_2 \in T_\alpha + F_\alpha \quad \text{or} \quad d_1, d_2 \in T'_\alpha + F'_\alpha.$$

We use relations (4.25) and (4.26) to define the action. We set

$$(7.13) \quad d_\alpha(\bar{\omega}_\alpha \otimes v_\alpha) = \sum_{\beta, \alpha \succ \beta} \langle d_\alpha, f_{\beta,\alpha} \rangle \pi_{\beta,\alpha}(\bar{\omega}_\alpha) \otimes A_{\beta,\alpha}(v_\alpha) \in \sum_{\beta, \alpha \succ \beta} M_\beta^0,$$

if $d_\alpha \in T_\alpha$, and

$$(7.14) \quad d_\alpha(\bar{\omega}_\alpha \otimes v_\alpha) = \sum_{\beta, \beta \succ \alpha} \langle \xi_{\beta,\alpha}, d_\alpha \rangle \pi_{\beta,\alpha}(\bar{\omega}_\alpha) A_{\beta,\alpha}(v_\alpha) \in \sum_{\beta, \beta \succ \alpha} M_\beta^0,$$

if $d_\alpha \in F_\alpha$. We set

$$(7.15) \quad d'_\alpha(\bar{\omega}_\alpha \otimes v_\alpha) = d'_\alpha \otimes (\bar{\omega}_\alpha \otimes v_\alpha) \in M_\alpha^1,$$

if $d'_\alpha \in T'_\alpha + F'_\alpha$.

As the next step we define the action of $BD^{\leq 1}$ on the subspace of degree one elements $M^1 \subset M$. Again we use decomposition (7.11). Define the action of $d' \in T'_\alpha + F'_\alpha$ on $d' \otimes (\bar{\omega}_\alpha \otimes v_\alpha) \in M_\alpha^1$ by the formula

$$(7.16) \quad d'_\alpha \cdot (d' \otimes (\bar{\omega}_\alpha \otimes v_\alpha)) = (d'_\alpha d') \otimes (\bar{\omega}_\alpha \otimes v_\alpha) \in M_\alpha^2.$$

Define the action of $d_\alpha \in T_\alpha + F_\alpha \subset BD^{\leq 1}$ by the formula

$$(7.17) \quad d_\alpha \cdot (d' \otimes (\bar{\omega}_\alpha \otimes v_\alpha)) = [d_\alpha, d'] \otimes (\bar{\omega}_\alpha \otimes v_\alpha) + d' \cdot d_\alpha(\bar{\omega}_\alpha \otimes v_\alpha),$$

where the first term in the right hand side is in M_α^0 , while the action of $BD^{\leq 1}$ for the second term was already defined.

We claim that formulas (7.13)–(7.17) define also an action of $BD^{\leq 2}$ on M^0 .

To prove this we note first that relations (7.13)–(7.17) define an action of all second degree monomials on M^0 and the only thing to check is to verify that this action respects quadratic relations in $BD^{\leq 2}$. The action of the basic relation

$$(7.18) \quad d_1 d_2 - d_2 d_1 = [d_1, d_2]$$

on M_α^0 can be split into several types depending on the types in (7.11) of the first degree generators d_i .

The case, when both d_i belong to $T'_\alpha + F'_\alpha$, is trivial. In that case d_i commute and their actions commute due to (7.15).

If $d_1 \in T'_\alpha + F'_\alpha$ and $d_2 \in T_\alpha + F_\alpha$, then relation (7.18) is preserved due to (7.17).

The nontrivial case is when both d_i belong to $T_\alpha + F_\alpha$ and should commute.

Suppose first that both $d_i \in T_\alpha$. Then

$$d_1 d_2(\bar{\omega}_\alpha \otimes v_\alpha) = \sum_{\beta, \alpha \succ \beta} \langle d_2, f_{\beta,\alpha} \rangle d_1(\pi_{\beta,\alpha}(\bar{\omega}_\alpha) \otimes A_{\beta,\alpha}(v_\alpha)).$$

We can present each of d_i as

$$d_i = c_i^\beta \xi_{\alpha,\beta} + d'_i,$$

where $\xi_{\alpha,\beta} \in T_\alpha \cap T'_\beta$, $f_{\beta,\alpha} \in F_\beta \cap T_\alpha^\perp$, d'_i belongs to T_β , and the constants c_i^β are given by the formula $c_i^\beta = \frac{\langle d_i, f_{\beta,\alpha} \rangle}{\langle \xi_{\alpha,\beta}, f_{\beta,\alpha} \rangle}$. Then

$$(7.19) \quad \begin{aligned} (d_1 d_2 - d_2 d_1)(\bar{\omega}_\alpha \otimes v_\alpha) &= \sum_{\beta, \alpha \succ \beta} \left(c_1^\beta \langle d_2, f_{\beta,\alpha} \rangle - c_2^\beta \langle d_1, f_{\beta,\alpha} \rangle \right) \pi_{\beta,\alpha}(\bar{\omega}_\alpha) \otimes A_{\beta,\alpha} v_\alpha + \\ &+ \sum_{\beta, \alpha \succ \beta} (\langle d_2, f_{\beta,\alpha} \rangle d'_1 - \langle d_1, f_{\beta,\alpha} \rangle d'_2) \cdot (\pi_{\beta,\alpha}(\bar{\omega}_\alpha) \otimes A_{\beta,\alpha} v_\alpha). \end{aligned}$$

The first summand in the right hand side of (7.19) is zero, since

$$\left(c_1^\beta \langle d_2, f_{\beta,\alpha} \rangle - c_2^\beta \langle d_1, f_{\beta,\alpha} \rangle \right) = 0$$

for each $\beta \in I(\Gamma)$, such that $\alpha \succ \beta$. The second summand can be rewritten as a double sum

$$(7.20) \quad \sum_{\substack{\beta, \alpha \succ \beta, \\ \gamma, \beta \succ \gamma}} C_{\alpha,\beta,\gamma} \otimes A_{\gamma\beta} A_{\beta,\alpha} v_\alpha = \sum_{\substack{\beta, \alpha \succ \beta, \\ \gamma, \beta \succ \gamma}} i_{d_1 \wedge d_2} (df_{\beta,\alpha} \wedge df_{\gamma\beta}) \pi_{\gamma,\beta} \pi_{\beta,\alpha}(\bar{\omega}_\alpha) \otimes A_{\gamma\beta} A_{\beta,\alpha} v_\alpha.$$

For a fixed γ the coefficient $C_{\alpha,\beta,\gamma}$ in (7.20) does not depend on index β . This can be seen from the equality

$$C_{\alpha,\beta,\gamma} = \frac{\bar{\omega}_\alpha}{df_{\beta,\alpha} \wedge df_{\gamma\beta}} i_{d_1 \wedge d_2} (df_{\beta,\alpha} \wedge df_{\gamma\beta}).$$

Then the sum in (7.20) equals zero due to relation (b) of Section 2.1.

The case when both d_i belong to F_α is treated in analogous manner. Let now $d_1 \in F_\alpha$, $d_2 \in T_\alpha$. Then

$$(7.21) \quad \begin{aligned} d_1 d_2 (\bar{\omega}_\alpha \otimes v_\alpha) &= \sum_{\beta, \alpha \succ \beta} \langle d_2, f_{\beta,\alpha} \rangle d_1 (\pi_{\beta,\alpha}(\bar{\omega}_\alpha) \otimes A_{\beta,\alpha} v_\alpha) = \\ &= \sum_{\substack{\beta, \alpha \succ \beta \\ \gamma, \gamma \succ \beta}} \langle d_2, f_{\beta,\alpha} \rangle \langle \xi_{\gamma\beta}, d_1 \rangle \pi_{\gamma,\beta} \pi_{\beta,\alpha}(\bar{\omega}_\alpha) \otimes A_{\gamma\beta} A_{\beta,\alpha} v_\alpha, \end{aligned}$$

since d_1 belongs to $F_\beta \subset F_\alpha$ for each summand in the first line of (7.21), and

$$(7.22) \quad \begin{aligned} d_2 d_1 (\bar{\omega}_\alpha \otimes v_\alpha) &= \sum_{\delta, \delta \succ \alpha} \langle \xi_{\delta\alpha}, d_1 \rangle d_2 (\pi_{\delta,\alpha}(\bar{\omega}_\alpha) \otimes A_{\delta,\alpha} v_\alpha) = \\ &= \sum_{\substack{\delta, \delta \succ \alpha \\ \tilde{\gamma}, \delta \succ \tilde{\gamma}}} \langle d_2, f_{\tilde{\gamma}\delta} \rangle \langle \xi_{\delta\alpha}, d_1 \rangle \pi_{\tilde{\gamma},\delta} \pi_{\delta,\alpha}(\bar{\omega}_\alpha) \otimes A_{\tilde{\gamma}\delta} A_{\delta,\alpha} v_\alpha, \end{aligned}$$

since d_2 belongs to $T_\delta \subset T_\alpha$ for each summand in the first line of (7.22).

We have the equalities of codimensions $l(\gamma) = l(\tilde{\gamma}) = l(\alpha)$, and the summation in the last lines of (7.21) and (7.22) goes over one index γ or $\tilde{\gamma}$ while $\bar{X}_\beta = \bar{X}_\alpha \cap \bar{X}_\gamma$ in (7.21) and X_δ is the unique hyperplane which contains both X_α and $X_{\tilde{\gamma}}$ in codimension 1 for (7.22).

Due to (7.12), we should prove that the difference of the right hand sides of (7.21) and (7.22) is zero. First of all, there are no terms with $\gamma = \alpha$ in the right hand side of (7.21),

since in this case the coefficient $\langle \xi_{\gamma\beta}, d_1 \rangle$ is zero. Analogously, there are no terms in the right hand side of (7.22) with $\tilde{\gamma} = \alpha$ since in this case the coefficient $\langle d_2, f_{\tilde{\gamma},\delta} \rangle$ vanishes.

For a vertex γ' of the graph Γ there are three different possibilities:

- (1) there is a summand in (7.21) with index $\gamma = \gamma'$ and a summand in (7.22) with index $\tilde{\gamma} = \gamma'$;
- (2) there is a summand in (7.21) with index $\gamma = \gamma'$ and there is no summand in (7.22) with $\tilde{\gamma} = \gamma'$;
- (3) there is no summand in (7.21) with index $\gamma = \gamma'$ and there is a summand in (7.22) with $\tilde{\gamma} = \gamma'$.

In case (2) the corresponding term vanishes due to relation (c) in Section 2.11. In case (3), the strata X_α and $X_{\tilde{\gamma}}$ are parallel inside X_δ , so the coefficient $\langle d_2, f_{\gamma',\delta} \rangle$ vanishes as well as the corresponding summand. In case (1) we can apply relation (c) of Section 2.1 if the sum of the coefficients in (7.21) and (7.22) is zero. So the only thing to prove is the equality

$$(7.23) \quad \langle d_2, f_{\beta,\alpha} \rangle \langle \xi_{\gamma\beta}, d_1 \rangle \pi_{\gamma,\beta} \pi_{\beta,\alpha}(\bar{\omega}_\alpha) + \langle d_2, f_{\gamma\delta} \rangle \langle \xi_{\delta\alpha}, d_1 \rangle \pi_{\gamma,\delta} \pi_{\delta,\alpha}(\bar{\omega}_\alpha) = 0,$$

when X_α and X_γ are strata of equal dimension, $\bar{X}_\beta = \bar{X}_\alpha \cap \bar{X}_\gamma$, $\bar{X}_\delta = \bar{X}_\alpha + \bar{X}_\gamma$ if an origin of $\mathbb{C}^N$ is taken inside X_β , $d_1 \in F_\alpha$ and $d_2 \in T_\alpha$. Then we can choose $\xi_{\delta\alpha} = \xi_{\gamma\beta}$ and $f_{\beta,\alpha} = f_{\gamma\delta}$. With such an identification the left hand side of (7.23) is equal to

$$(d_2, f_{\gamma,\delta}) \langle \xi_{\gamma\beta}, d_1 \rangle (\pi_{\gamma,\beta} \pi_{\beta,\alpha}(\bar{\omega}_\alpha) + \pi_{\gamma,\delta} \pi_{\delta,\alpha}(\bar{\omega}_\alpha)).$$

The equality $\bar{\omega}_\gamma = \pi_{\gamma,\beta} \pi_{\beta,\alpha}(\bar{\omega}_\alpha)$ is equivalent to the equality

$$\bar{\omega}_\alpha = df_{\beta,\alpha} \wedge i_{\xi_{\gamma,\beta}} \bar{\omega}_\gamma = df_{\gamma,\delta} \wedge i_{\xi_{\gamma,\beta}} \bar{\omega}_\gamma,$$

and the equality $\bar{\omega}_\gamma = \pi_{\gamma,\delta} \pi_{\delta,\alpha}(\bar{\omega}_\alpha)$ is equivalent to

$$\bar{\omega}_\alpha = i_{\xi_{\delta,\alpha}}(df_{\gamma,\delta} \wedge \bar{\omega}_\gamma) = i_{\xi_{\gamma,\beta}}(df_{\gamma,\delta} \wedge \bar{\omega}_\gamma).$$

The elements $\xi_{\gamma\beta}$ and $f_{\gamma\delta}$ commute in D_X by (7.12). This implies the equality $df_{\gamma,\delta} \wedge i_{\xi_{\gamma,\beta}} \bar{\omega}_\gamma + i_{\xi_{\gamma,\beta}}(df_{\gamma,\delta} \wedge \bar{\omega}_\gamma) = 0$ which proves (7.23).

Suppose $n \geq 0$ and $k, 0 \leq k \leq n$ are given. Suppose an action of $B^{\leq k}$ on M^l is defined, if $l \leq n - k$. Let $d \in BD^{\leq 1}$. For a given stratum X_α and a monomial D of degree n over $\mathbb{C}[T'_\alpha + F'_\alpha]$ define

$$(7.24) \quad d \cdot D \otimes \bar{\omega}_\alpha \otimes v_\alpha = [d, D] \otimes \bar{\omega}_\alpha \otimes v_\alpha + D \cdot d(\bar{\omega}_\alpha \otimes v_\alpha) \quad \text{if} \quad d \in T_\alpha + F_\alpha,$$

$$(7.25) \quad d \cdot D \otimes \bar{\omega}_\alpha \otimes v_\alpha = dD \otimes \bar{\omega}_\alpha \otimes v_\alpha \quad \text{if} \quad d \in T'_\alpha + F'_\alpha.$$

Then the action of all monomials from $B^{\leq k}$ on M^{n-k+1} is defined. We should check that this action preserves the commutation relations in the ring D_X , more precisely, the relations of the type

$$(7.26) \quad d_1 \cdot d_2 \cdot (D \otimes \bar{\omega}_\alpha \otimes v_\alpha) - d_2 \cdot d_1 \cdot (D \otimes \bar{\omega}_\alpha \otimes v_\alpha) = [d_1, d_2](D \otimes \bar{\omega}_\alpha \otimes v_\alpha)$$

where $d_i \in BD^{\leq 1}$ and D is a monomial of degree $n - 1$ over $\mathbb{C}[T'_\alpha + F'_\alpha]$. If both $d_i \in T'_\alpha + F'_\alpha$ then (7.26) is satisfied due to (7.25). Let now $d_1 \in T_\alpha + F_\alpha$, $d_2 \in T'_\alpha + F'_\alpha$. Then by (7.24) and by induction assumptions

$$\begin{aligned} d_1 d_2 (D \otimes \bar{\omega}_\alpha \otimes v_\alpha) &= [d_1, d_2 D] \otimes \bar{\omega}_\alpha \otimes v_\alpha + d_2 D d_1 (\bar{\omega}_\alpha \otimes v_\alpha), \\ d_2 d_1 (D \otimes \bar{\omega}_\alpha \otimes v_\alpha) &= d_2 [d_1, D] \otimes \bar{\omega}_\alpha \otimes v_\alpha + d_2 (D d_1 (\bar{\omega}_\alpha \otimes v_\alpha)), \end{aligned}$$

so we get

$$(d_1 d_2 - d_2 d_1) \cdot (D \otimes \bar{\omega}_\alpha \otimes v_\alpha) = [d_1, d_2] D \otimes \bar{\omega}_\alpha \otimes v_\alpha.$$

Let now $d_1, d_2 \in T_\alpha + F_\alpha$. Then by (7.24)

$$d_1 d_2 (D \otimes \bar{\omega}_\alpha \otimes v_\alpha) = [d_1, [d_2, D]] \otimes \bar{\omega}_\alpha \otimes v_\alpha + [d_2, D] (d_1 (\bar{\omega}_\alpha \otimes v_\alpha)) + d_1 (D d_2 (\bar{\omega}_\alpha \otimes v_\alpha))$$

and

$$(d_1 d_2 - d_2 d_1) \cdot (D \otimes \bar{\omega}_\alpha \otimes v_\alpha) = D (d_1 d_2 - d_2 d_1) (\bar{\omega}_\alpha \otimes v_\alpha) = 0$$

by induction assumptions. This ends the induction step and the proof of the proposition. $\square$

7.4. Proof of Proposition 4.2. The description (4.25) and (4.26) of a quiver $\mathcal{D}_X$ -module imply that the principal filtration (4.31) is good and the map

$$\phi : \bigoplus_{\alpha \in I(\Gamma)} (S(T'_\alpha) \otimes \bar{\Omega}_\alpha \otimes V_\alpha) \otimes \mathbb{C}[\bar{X}_\alpha] \rightarrow \bar{E}\mathcal{V},$$

where $\bar{E}\mathcal{V} = \text{gr } E\mathcal{V}$, is a well defined map of $\mathcal{O}_X$ -modules. Proposition 4.1 implies that this map is an isomorphism.

Consider the product

$$\prod_{\alpha \in I(\Gamma)} \bar{d}_\alpha,$$

where $d_\alpha \in T_\alpha + F_\alpha$ and $\bar{d}_\alpha = \text{gr}(d_\alpha)$. Relations (4.25), (4.26), (7.24) and (7.25) show that this product annihilates $\bar{E}\mathcal{V}$. It means that $E\mathcal{V}$ is a holonomic $\mathcal{D}_X$ module with singular support contained in $\cup_\alpha T_{X_\alpha}^* \mathbb{C}^N$. $\square$

7.5. Proof of Proposition 4.4. Fix a nonzero exterior form $\bar{\omega}_\emptyset \in \bar{\Omega}_\emptyset$. Identify vectors $v \in V$ with $\bar{\omega}_\emptyset \otimes v \in \bar{\Omega}_\emptyset \otimes V$, and $w \in W$ with $\bar{\omega}_\emptyset \otimes w \in \bar{\Omega}_\emptyset \otimes W$.

Let $\Phi : E^0\mathcal{V} \rightarrow E^0\mathcal{W}$ be a morphism of $\mathcal{D}_{X^0}$ -modules. Since X_0 is an affine variety, the morphism is determined by the corresponding map of global sections, denoted by Φ also. Since the $\mathcal{D}_{X^0}$ -module $E^0\mathcal{V}$ is generated by the space V , the map of global sections is determined by its restriction to V , denote $\varphi = \Phi|_V$. The map φ is a regular rational function on X^0 with values in $\text{Hom}(V, W)$. We need to prove that φ is constant.

For $j \in J(\mathcal{C})$, choose affine coordinates $z_1, \dots, z_N$ on X , such that the hyperplane H_j has equation $z_1 = 0$. Take a generic point $(0, \varepsilon_2, \dots, \varepsilon_N)$ in H_j . Expand φ in the Laurent series,

$$(7.27) \quad \varphi = \frac{\varphi_k}{z_1^k} + \frac{\varphi_{k-1}}{z_1^{k-1}} + \dots,$$

where φ_k are functions in $z_2, \dots, z_N$ holomorphic at $(\varepsilon_2, \dots, \varepsilon_N)$. We assume that $\varphi_k(\varepsilon_2, \dots, \varepsilon_N) \neq 0$. For $v \in V$, consider the Laurent expansion for both sides of the equality $\Phi(\partial_{z_1} v) = \partial_{z_1}(\Phi(v))$. We have

$$\Phi(\partial_{z_1} v) = \Phi\left(\frac{A^j(v)}{z_1} + \dots\right) = \frac{1}{z_1} \varphi(A^j(v)) + \dots = \frac{\varphi_k(A^j(v))}{z_1^{k+1}} + \dots,$$

and

$$\partial_{z_1}(\Phi(v)) = \partial_{z_1}(\varphi(v)) = \frac{(B^j - k)(\varphi_k(v))}{z_1^{k+1}} + \dots.$$

We get $\varphi_k(A^j(v)) = (B^j - k)(\varphi_k(v))$. Since eigenvalues of A^j and of B^j are in the same weakly non-resonant set G^j , we conclude that $k = 0$. Thus the rational function φ has no poles and is a polynomial function on X .

Let $\mathbb{C}[X]$ be the ring of polynomials on X with complex coefficients. Denote $\tilde{V} = \mathbb{C}[X] \cdot V$, $\tilde{W} = \mathbb{C}[X] \cdot W$. We proved that $\Phi(\tilde{V}) \subset \tilde{W}$.

Let $O \in X$ be the center of the arrangement. Let $z_1, \dots, z_n$ be affine coordinates on X with center at O . Let η be the associated Euler vector field, $\eta = z_1 \partial_{z_1} + \dots + z_n \partial_{z_n}$. For any $v \in V$, we have $\eta(v) = \sum_{j \in J(C)} A^j(v)$. Let $\tilde{V} = \bigoplus_{\mu} \tilde{V}_{\mu}$, $\tilde{W} = \bigoplus_{\mu} \tilde{W}_{\mu}$ be the decompositions into eigenspaces of η . It is easy to see that the summations are over $\mu \in \tilde{G}_{\emptyset} + \mathbb{Z}_{\geq 0}$. We have $\Phi(\tilde{V}_{\mu}) \subset \tilde{W}_{\mu}$ for any μ . Since the set $\tilde{G}_{\emptyset}$ is weakly non-resonant, we have equalities

$$V = \bigoplus_{\mu \in \tilde{G}_{\emptyset}} \tilde{V}_{\mu}, \quad W = \bigoplus_{\mu \in \tilde{G}_{\emptyset}} \tilde{W}_{\mu},$$

and $\Phi(V) \subset W$. Thus φ is constant. The condition $[\Phi, \xi](v) = 0$ for constant vector fields ξ imply that φ is a morphism of quivers and $\Phi = E^0(\varphi)$. $\square$

7.6. Proof of Proposition 4.3. Part (i) of the proposition follows from the description in Section 4.4 of global sections of a quiver $\mathcal{D}_{X^n}$ -module. Namely, the quiver $\mathcal{D}_{X^n}$ -module $E^n \mathcal{V}_{\Gamma^n}$ has a finite-dimensional subspace $\overline{V} = \bigoplus_{\alpha \in I(\Gamma^n)} \overline{\Omega}_{\alpha} \otimes V_{\alpha}$ of the space of its global sections. If $\varphi : \mathcal{V}_{\Gamma^n} \rightarrow \mathcal{W}_{\Gamma^n}$ is a nonzero morphism of quivers, then the associated morphism $E^n \varphi : E^n \mathcal{V}_{\Gamma^n} \rightarrow E^n \mathcal{W}_{\Gamma^n}$ of $\mathcal{D}_{X^n}$ -modules is nonzero on the space $\overline{V}$. Thus the functor E^n is faithful.

The proof of part (ii) is by induction on n . The proof is based on the Beilinson-Kashiwara-Malgrange gluing construction.

First we will recall the gluing construction, following [K3]. Then we will introduce special open subsets $X^{n,\beta} \subset X^n$, on which the gluing procedure will be performed. For a level n quiver $\mathcal{V}_{\Gamma^n}$, we will restrict the quiver $\mathcal{D}_{X^n}$ -module $E^n \mathcal{V}_{\Gamma^n}$ to an open set $X^{n,\beta}$ and describe, how this $\mathcal{D}_{X^{n,\beta}}$ -module can be obtained by gluing quiver $\mathcal{D}$ -modules of level $n - 1$. We will note that the sheaf axioms imply that the category $\mathcal{M}_{X^n}$ of $\mathcal{D}_{X^n}$ -modules is equivalent to a category of fibered products of $\mathcal{M}_{X^{n,\beta}}$ -modules. In this way we will describe the $\mathcal{D}_{X^n}$ -module $E^n \mathcal{V}_{\Gamma^n}$ by means of quiver $\mathcal{D}$ -modules of level $n - 1$ quivers. With such a description we will perform the induction step.

7.6.1. Kashiwara-Malgrange filtrations and the gluing construction. Recall that a subset $\overline{G} \subset \mathbb{C}$ is called a *non-resonant section* of $\mathbb{C}$, if $\overline{G}$ contains zero and for any $a \in \mathbb{C}$, the intersection $\overline{G} \cap (a + \mathbb{Z})$ consists of a single point.

Let Y be a smooth complex algebraic variety and $Z \subset Y$ a smooth complex algebraic subvariety. We assume that Z is a principal divisor, defined by the equation $f = 0$ for some regular function f . Let $I \subset \mathcal{O}_Y$ be the sheaf of ideals of functions vanishing on Z and $F(\mathcal{D}_Y)$ a corresponding (see Section 4.6 decreasing filtration of the sheaf $\mathcal{D}_Y$:

$$F^k(\mathcal{D}_Y) = \{P \in \mathcal{D}_Y \mid P(I_{\beta}^j) \subset I_{\beta}^{j+k} \text{ for any } j\}.$$

Let θ be a vector field in Y tangent to Z and acting on I/I^2 as the identity. We suppose that θ is presented as a product $\theta = f\xi$, where ξ is a vector field on Y such that $\xi(df) = 1$.

Let M be a $\mathcal{D}_Y$ -module. For any good filtration $F(M)$ of a $\mathcal{D}_Y$ -module M , the quotient spaces $\text{gr } F^k(M) = F^k(M)/F^{k+1}(M)$ have the structure of $\mathcal{D}_Z$ -modules. and the function f and the vector field ξ define the $\mathcal{D}_Z$ -module maps

$$\text{gr } f : \text{gr } F^k(M) \rightarrow \text{gr } F^{k+1}(M) \quad \text{and} \quad \text{gr } \xi : \text{gr } F^k(M) \rightarrow \text{gr } F^{k-1}(M) ,$$

such that $\text{gr } f \cdot \text{gr } \xi = \text{gr } \theta : \text{gr } F^k(M) \rightarrow \text{gr } F^k(M)$.

Suppose a non-resonant section $\overline{G} \subset \mathbb{C}$ is chosen. Let $F_{\overline{G}}(M)$ be the unique good filtration of M , satisfying the condition (v) of Section 4.6. Denote $\Psi_{\overline{G}}^{(k)}(M) = \text{gr } F_{\overline{G}}^k(M)$. Then for all $k \geq 0$, the $\mathcal{D}_Z$ -modules $\Psi_{\overline{G}}^{(k)}(M)$ and the maps $\text{gr } \theta : \Psi_{\overline{G}}^{(k)}(M) \rightarrow \Psi_{\overline{G}}^{(k)}(M)$ depend on the restriction $M|_{Y \setminus Z}$ only. We use this remark in notations and set $\Psi_{\overline{G}}^{(0)}(M|_{Y \setminus Z}) = \Psi_{\overline{G}}^{(0)}(M)$. Denote by a the map $a = \text{gr } \xi|_{\Psi_{\overline{G}}^{(0)}(M)} : \Psi_{\overline{G}}^{(0)}(M) \rightarrow \Psi_{\overline{G}}^{(-1)}(M)$ and by b the map $b = \text{gr } f|_{\Psi_{\overline{G}}^{(-1)}(M)} : \Psi_{\overline{G}}^{(-1)}(M) \rightarrow \Psi_{\overline{G}}^{(0)}(M)$.

The Beilinson-Malgrange-Kashiwara gluing theorem [Be], [K3], [M] states that for a given non-resonant section $\overline{G} \subset \mathbb{C}$ the assignment

$$M \rightarrow (M|_{Y \setminus Z}, \Psi_{\overline{G}}^{(-1)}(M), a, b)$$

establishes an equivalence of the category $\mathcal{M}_Y^{hrs} \subset \mathcal{M}_Y^{hol}$ of holonomic $\mathcal{D}_Y$ -modules with regular singularities and the category of quadruples $(\tilde{M}, N, a, b)$, where $\tilde{M} \in \mathcal{M}_{Y \setminus Z}^{hrs}$, $N \in \mathcal{M}_Z^{hrs}$, $a : \Psi_{\overline{G}}^{(0)}(\tilde{M}) \rightarrow N$ and $b : N \rightarrow \Psi_{\overline{G}}^{(0)}(\tilde{M})$ are $\mathcal{D}_Z$ -module maps such that $ba = \text{gr } \theta : \Psi_{\overline{G}}^{(0)}(j_* \tilde{M}) \rightarrow \Psi_{\overline{G}}^{(0)}(j_* \tilde{M})$.

7.6.2. *The spaces $X^{n,\beta}$.* Take a vertex $\beta \in I(\Gamma^n)$, such that $l(\beta) = n$. Let $X^{n,\beta} \subset X^n$ be the following open subset of X^n ,

$$X^{n,\beta} = X^{n-1} \cup X_\beta = X^n \setminus \left(\bigcup_{\alpha \neq \beta, l(\alpha)=n} X_\alpha \right).$$

The sets $X^{n,\beta}$ cover X^n ,

$$X^n = \bigcup_{\beta, l(\beta)=n} X^{n,\beta},$$

and for any $\beta_1, \beta_2 \in I_n(\Gamma)$ we have

$$X^{n,\beta_1} \cap X^{n,\beta_2} = X^{n-1}.$$

Thus the category $\mathcal{M}_{X^n}$ of $\mathcal{D}_{X^n}$ -modules is the fibered product of the categories $\mathcal{M}_{X^{n,\beta}}$:

$$(7.28) \quad \mathcal{M}_{X^n} \approx \mathcal{M}_{X^{n,\beta_1}} \times_{\mathcal{M}_{X^{n-1}}} \mathcal{M}_{X^{n,\beta_1}} \times_{\mathcal{M}_{X^{n-1}}} \cdots \times_{\mathcal{M}_{X^{n-1}}} \mathcal{M}_{X^{n,\beta_m}},$$

where the product is taken over all $\beta_k \in I_n(\Gamma)$. An object of the category $\mathcal{M}_{X^{n,\beta_1}} \times_{\mathcal{M}_{X^{n-1}}} \cdots \times_{\mathcal{M}_{X^{n-1}}} \mathcal{M}_{X^{n,\beta_m}}$ is a collection $M_1 \in \mathcal{M}_{X^{n,\beta_1}}, \dots, M_m \in \mathcal{M}_{X^{n,\beta_m}}$, such that $M_1|_{X^{n-1}} = M_2|_{X^{n-1}} = \dots = M_m|_{X^{n-1}}$.

Let $\mathcal{F}_{n-1} = \{\bar{f}_\alpha \mid \alpha \in I_n(\Gamma)\}$ be a level $n-1$ vertex framing of the arrangement $\mathcal{C}$. The framing determines an affine open subset $U_{\mathcal{F}_{n-1}}^\beta \subset X^{n,\beta}$ of $X^{n,\beta}$:

$$(7.29) \quad U_{\mathcal{F}_{n-1}}^\beta = X \setminus \bigcup_{\alpha \in I_n(\Gamma), \alpha \neq \beta} \{\bar{f}_\alpha = 0\}.$$

Subsets $U_{\mathcal{F}_{n-1}}^\beta$ form a covering of $X^{n,\beta}$ when $\mathcal{F}_{n-1}$ varies.

7.6.3. Gluing data for Γ^n -quivers. Fix a vertex $\beta \in I_n(\Gamma)$. Let $\bar{f}_\beta \in F_\beta$ be a generic affine function vanishing on X_β . In the notations of Section 7.6.1, put $Y = X^{n,\beta}$ and $Z = Y \cap H$, where $H = \{\bar{f}_\beta = 0\}$.

Intersections $H_j \cap H$, $j \in J(\mathcal{C})$, determine in H an arrangement, denoted by $\mathcal{C}_H$, which equips H with a structure of a stratified space. The space Z coincides with the principal level $n-1$ open subset of H :

$$Z = H^{n-1} = H \setminus \bigcup_{\text{codim}_H H_\alpha \geq n} H_\alpha.$$

The strata of Z are the intersections $H \cap X_\alpha$, where $\alpha \in I(\Gamma)$, such that $l(\alpha) < n$.

The graph of adjacencies $\Gamma^{n,\beta}$ of the stratified space Z is a certain contraction of the graph Γ^{n-1} .

More precisely, let $I_{n-1}^\beta(\Gamma)$ be the set of classes of isomorphisms of elements of $I_{n-1}(\Gamma)$ with respect to the following equivalence relation: $\alpha_1 \approx \alpha_2$ if $\alpha_1 \succ \beta$ and $\alpha_2 \succ \beta$, or $\alpha_1 = \alpha_2$. Thus, the set $I_{n-1}^\beta(\Gamma)$ consists of the vertices $\alpha \in I_{n-1}(\Gamma)$, such that $\alpha \not\succ \beta$ and of the equivalence class $\bar{\beta}$ of the vertices $\alpha \in I_{n-1}(\Gamma)$, such that $\alpha \succ \beta$.

Let $\tilde{I}_n^\beta(\Gamma)$ be the subset of $I_n(\Gamma)$, which consists of the elements $\beta' \in I_n(\Gamma)$, such that the stratum $X_{\beta'}$ is not equal to the stratum X_β and is not parallel to it (not obtained by a parallel translation). Let $I_n^\beta(\Gamma)$ be the set of classes of isomorphisms of elements of $\tilde{I}_n^\beta(\Gamma)$ with respect to the following equivalence relation: $\alpha_1 \approx \alpha_2$ if $\alpha_1 = \alpha_2$ or the intersections $\bar{X}_{\alpha_1} \cap \bar{X}_\beta$ and $\bar{X}_{\alpha_2} \cap \bar{X}_\beta$ coincide and have codimension one in $\bar{X}_\beta$:

$$\bar{X}_{\alpha_1} \cap \bar{X}_\beta = \bar{X}_{\alpha_2} \cap \bar{X}_\beta \quad \text{and} \quad \text{codim}_{\bar{X}_\beta} \bar{X}_{\alpha_1} \cap \bar{X}_\beta = 1.$$

In these notations the set of vertices of $\Gamma^{n,\beta}$ is the union

$$I(\Gamma^{n,\beta}) = I(\Gamma^{n-2}) \cup I_{n-1}^\beta(\Gamma).$$

Define the edges of $\Gamma^{n,\beta}$. The vertices $\alpha_1, \alpha_2 \in I(\Gamma^{n,\beta})$, such that $\alpha_1 \neq \bar{\beta}$, $\alpha_2 \neq \bar{\beta}$ and $\alpha_1 \neq \alpha_2$, are connected by an edge in $\Gamma^{n,\beta}$ if and only if they are connected by an edge in Γ^{n-1} . The vertices $\bar{\beta}$ and $\alpha \in I(\Gamma^{n,\beta})$, $\alpha \neq \beta$ define an edge in $\Gamma^{n,\beta}$ if there exists $\alpha_1 \succ \beta$, such that α_1 and α are connected by an edge in Γ^{n-1} .

The graph $\Gamma^{n,\beta}$ has loops $(\bar{\alpha}_1, \bar{\alpha}_1)^{\bar{\alpha}_2}$, where $\bar{\alpha}_1 \in I_{n-1}^\beta(\Gamma)$, $\bar{\alpha}_2 \in I_n^\beta(\Gamma)$ and there exist representatives α_1 of $\bar{\alpha}_1$, $\alpha_1 \in I_{n-1}(\Gamma)$, and α_2 of $\bar{\alpha}_2$, $\alpha_2 \in \tilde{I}_n^\beta(\Gamma)$, such that $\alpha_1 \succ \alpha_2$.

Let $\mathcal{V}_{\Gamma^{n-1}} = \{V_\alpha, A_{\alpha_1, \alpha_2}, A_{\alpha_2}^{\alpha_1}\} \in \text{Qui}_{\Gamma^{n-1}}$ be a quiver of Γ^{n-1} . Define a new $\Gamma^{n,\beta}$ -quiver $\bar{\psi}^{(0)}(\mathcal{V}_{\Gamma^{n-1}}) = \{W_\alpha, B_{\alpha_1, \alpha_2}, B_{\alpha_1}^{\alpha_2}\}$ in the following way.

Set $W_\alpha = V_\alpha$ for all $\alpha \in I_{n-1}(\Gamma)$, and $W_{\bar{\beta}} = \bigoplus_{\alpha \succ \beta} V_\alpha$.

Set $B_{\alpha_1, \alpha_2} = A_{\alpha_1, \alpha_2} : W_{\alpha_2} \rightarrow W_{\alpha_1}$ if both α_1 and α_2 are different from $\bar{\beta}$. Set $B_{\bar{\beta}, \alpha_2} = \oplus_{\alpha, \alpha \succ \bar{\beta}} A_{\alpha, \alpha_2} : W_{\alpha_2} \rightarrow W_{\bar{\beta}}$ and $B_{\alpha_1, \bar{\beta}} = \oplus_{\alpha, \alpha \succ \bar{\beta}} A_{\alpha_1, \alpha} : W_{\bar{\beta}} \rightarrow W_{\alpha_1}$.

Let $(\alpha_1, \alpha_1)^{\bar{\alpha}_2}$ be a loop of $\Gamma^{n, \beta}$ where $\alpha_1 \in I_{n-1}^\beta(\Gamma)$, $\bar{\alpha}_2 \in I_n^\beta(\Gamma)$ and $\alpha_1 \neq \bar{\beta}$. Define the map $B_{\alpha_1}^{\bar{\alpha}_2} : W_{\alpha_1} \rightarrow W_{\alpha_1}$ by the formula

$$B_{\alpha_1}^{\bar{\alpha}_2} = \sum_{j=1}^{m_{\bar{\alpha}_2}} A_{\alpha_1}^{\tilde{\alpha}_j},$$

where $\tilde{\alpha}_1, \dots, \tilde{\alpha}_{m_{\bar{\alpha}_2}} \in \tilde{I}_n^\beta(\Gamma)$ are all representatives of the equivalence class $\bar{\alpha}_2 \in I_n^\beta(\Gamma)$.

Let $(\bar{\beta}, \bar{\beta})^{\bar{\alpha}_2}$ be a loop of $\Gamma^{n, \beta}$. Let $\tilde{\alpha}_1, \dots, \tilde{\alpha}_{m_{\bar{\alpha}_2}} \in \tilde{I}_n^\beta(\Gamma)$ be all representatives of the equivalence class $\bar{\alpha}_2 \in I_n^\beta(\Gamma)$, and $\alpha_1, \dots, \alpha_m \in I_{n-1}^\beta(\Gamma)$ all representatives of the equivalence class $\bar{\beta}$. Define the map $B_{\bar{\beta}}^{\bar{\alpha}_2} : W_{\bar{\beta}} \rightarrow W_{\bar{\beta}}$ by the formula

$$B_{\bar{\beta}}^{\bar{\alpha}_2} = \sum_{j_1=1}^m \sum_{j_2=1}^{m_{\bar{\alpha}_2}} A_{\alpha_{j_1}}^{\tilde{\alpha}_{j_2}} - \sum_{j_1, j_2=1, \dots, m, j_1 \neq j_2} A_{\alpha_{j_1}, \delta(\alpha_{j_1}, \alpha_{j_2})} A_{\delta(\alpha_{j_1}, \alpha_{j_2}), \alpha_{j_2}}.$$

The vertex $\delta(\alpha_{j_1}, \alpha_{j_2})$ is defined by the following rule. Suppose there exists a vertex $\gamma \in \{\tilde{\alpha}_1, \dots, \tilde{\alpha}_{m_{\bar{\alpha}_2}}\}$, such that $\alpha_{j_1} \succ \gamma$ and $\alpha_{j_2} \succ \gamma$. Then we set $\delta(\alpha_{j_1}, \alpha_{j_2})$ to be the unique vertex of Γ^{n-1} , such that $\delta(\alpha_{j_1}, \alpha_{j_2}) \succ \alpha_{j_1}$ and $\delta(\alpha_{j_1}, \alpha_{j_2}) \succ \alpha_{j_2}$. Otherwise the corresponding term is supposed to be zero.

We define a quiver morphism $\bar{T}_\beta : \bar{\psi}^{(0)}(\mathcal{V}_{\Gamma^{n-1}}) \rightarrow \bar{\psi}^{(0)}(\mathcal{V}_{\Gamma^{n-1}})$. Morphism $\bar{T}_\beta$ is a collection of linear operators $\bar{T}_\beta|_{W_\alpha} : W_\alpha \rightarrow W_\alpha$, $\alpha \in I(\Gamma^{n, \beta})$, commuting with the maps B_{α_1, α_2} and $B_{\alpha_1}^{\alpha_2}$. The operators $\bar{T}_\beta|_{W_\alpha}$ are zero for all α except $\alpha = \bar{\beta}$, and

$$(7.30) \quad \bar{T}_\beta|_{W_{\bar{\beta}}} = \sum_{j=1}^m A_{\alpha_j}^\beta - \sum_{j_1, j_2=1, \dots, m, j_1 \neq j_2} A_{\alpha_{j_1}, \delta(\alpha_{j_1}, \alpha_{j_2})} A_{\delta(\alpha_{j_1}, \alpha_{j_2}), \alpha_{j_2}}.$$

Here $\alpha_1, \dots, \alpha_m \in I_{n-1}^\beta(\Gamma)$ are all representatives of the equivalence class $\bar{\beta}$. The vertex $\delta(\alpha_{j_1}, \alpha_{j_2})$ is the unique vertex (if exists) of Γ^{n-1} , such that $\delta(\alpha_{j_1}, \alpha_{j_2}) \succ \alpha_{j_1}$ and $\delta(\alpha_{j_1}, \alpha_{j_2}) \succ \alpha_{j_2}$.

Let $\mathcal{V}_{\Gamma^n}$ be a quiver of Γ^n . We set $\psi^{(0)}(\mathcal{V}_{\Gamma^n}) = \bar{\psi}^{(0)}(\mathbb{J}_{n, n-1}^* \mathcal{V}_{\Gamma^n})$.

Lemma 7.1. *For any level n quiver $\mathcal{V}_{\Gamma^n}$ the linear map $\bar{T}_\beta|_{W_{\bar{\beta}}}$ coincides with the operator T_β , defined in (3.5).*

Proof. The statement follows from relations (ii), Section 3.2. $\square$

The closure $X' = \overline{X}_\beta$ of the stratum X_β is an affine plane. The arrangement $\mathcal{C}$ induces in X' an arrangement denoted by $\mathcal{C}_\beta$. The stratum X_β is the complement in X' to the union of hyperplanes of the arrangement $\mathcal{C}_\beta$. The hyperplanes $\overline{X}'_\gamma$ of the arrangement $\mathcal{C}_\beta$ are closures of the strata $X_\gamma \subset \overline{X}_\beta$ of codimension one in $\overline{X}_\beta$. They are in one-to-one correspondence with elements of the set I_n^β : to each equivalence class $\{\alpha_i\}$ of I_n^β we attach a hyperplane $\overline{X}_{\alpha_i} \cap \overline{X}_\beta$ in $X' = \overline{X}_\beta$.

Denote by Γ^β the level zero graph of the arrangement $\mathcal{C}_\beta$. We have a map of graphs $i_\beta : \Gamma^\beta \rightarrow \Gamma^{n, \beta}$, which sends the vertex $\emptyset$ of Γ^β to the vertex $\bar{\beta}$ of $\Gamma^{n, \beta}$ and the loop $(\emptyset, \emptyset)^\gamma$

of Γ^β to the loop $(\bar{\beta}, \bar{\beta})^{i_\beta(\gamma)}$ of $\Gamma^{n,\beta}$, where $i_\beta(\gamma) \in I_n^\beta$ is characterized by the condition that $\bar{X}_\alpha \cap \bar{X}_\beta = \bar{X}'_\gamma$ for any representative α of $i_\beta(\gamma)$.

To any Γ^β -quiver $\mathcal{V}' = \{V'_\emptyset, A'^\gamma_\emptyset\}$ we attach the following $\Gamma^{n,\beta}$ -quiver $i_\beta(\mathcal{V}) = \mathcal{U} = \{U_\alpha, C_{\alpha_1, \alpha_2}, C_{\alpha_1}^{\alpha_2}\}$. The spaces U_α of the quiver $\mathcal{U}$ are all zero-dimensional except for $\alpha = \bar{\beta}$. Set $U_{\bar{\beta}} = V'_\emptyset$. All operators $C_{\alpha_1, \alpha_2}, C_{\alpha_1}^{\alpha_2}$ are the zero operators except for operators $C_{\bar{\beta}}^{i_\beta(\gamma)}$. We set $C_{\bar{\beta}}^{i_\beta(\gamma)} = A'^\gamma_\emptyset : U_{\bar{\beta}} \rightarrow U_{\bar{\beta}}$.

Let $\mathcal{V}_{\Gamma^n} = \{V_\alpha, A_{\alpha_1, \alpha_2}, A_{\alpha_2}^{\alpha_1}\} \in \text{Qui}_{\Gamma^n}$ be a quiver of Γ^n . Define a quiver $\bar{\psi}_\beta^{(-1)}(\mathcal{V}_{\Gamma^n}) = \{V'_\emptyset, A'^\gamma_\emptyset\}$ of Γ^β as follows. We set $V'_\emptyset = V_\beta$ and $A'^\gamma_\emptyset = A_\beta^\gamma$. Define a quiver $\psi_\beta^{(-1)}(\mathcal{V}_{\Gamma^n}) = \{V'_\emptyset, A'^\gamma_\emptyset\}$ of $\Gamma^{n,\beta}$ as $i_\beta(\bar{\psi}_\beta^{(-1)}(\mathcal{V}_{\Gamma^n}))$.

Define two morphisms of the constructed quivers: $\bar{a}_\beta : \psi_\beta^{(0)}(\mathcal{V}_{\Gamma^n}) \rightarrow \psi_\beta^{(-1)}(\mathcal{V}_{\Gamma^n})$ and $\bar{b}_\beta : \psi_\beta^{(-1)}(\mathcal{V}_{\Gamma^n}) \rightarrow \psi_\beta^{(0)}(\mathcal{V}_{\Gamma^n})$. To define these morphisms, it is sufficient to define linear maps $\bar{a}_\beta|_{W_{\bar{\beta}}} : W_{\bar{\beta}} \rightarrow U_{\bar{\beta}}$ and $\bar{b}_\beta|_{U_{\bar{\beta}}} : U_{\bar{\beta}} \rightarrow W_{\bar{\beta}}$. We set

$$\bar{a}_\beta(v_\alpha) = A_{\beta, \alpha}(v_\alpha), \quad \bar{b}_\beta(v_\beta) = \sum_{\alpha \in I_{n-1}(\Gamma), \alpha \succ \beta} A_{\alpha, \beta}(v_\beta),$$

for any $v_\beta \in V_\beta$ and $v_\alpha \in V_\alpha$. We have the equality

$$\bar{b}_\beta \bar{a}_\beta = \bar{T}_\beta.$$

Let $Q^{n,\beta}$ be the category of quadruples $(\mathcal{V}_{\Gamma^{n-1}}, \mathcal{U}_{\Gamma^\beta}, a_\beta, b_\beta)$, where $\mathcal{V}_{\Gamma^{n-1}}$ is a quiver of Γ^{n-1} , $\mathcal{U}_{\Gamma^\beta}$ is a quiver of Γ^β , a_β and b_β are morphisms of $\Gamma^{n,\beta}$ -quivers, $a_\beta : \bar{\psi}_\beta^{(0)}(\mathcal{V}_{\Gamma^{n-1}}) \rightarrow i_\beta(\mathcal{U}_{\Gamma^\beta})$ and $b_\beta : i_\beta(\mathcal{U}_{\Gamma^\beta}) \rightarrow \bar{\psi}_\beta^{(0)}(\mathcal{V}_{\Gamma^{n-1}})$, such that the composition $ba : \bar{\psi}_\beta^{(0)}(\mathcal{V}_{\Gamma^{n-1}}) \rightarrow \bar{\psi}_\beta^{(0)}(\mathcal{V}_{\Gamma^{n-1}})$ is equal to $\bar{T}_\beta$, see (7.30). A morphism between objects $(\mathcal{V}_{\Gamma^{n-1}}, \mathcal{U}_{\Gamma^\beta}, a_\beta, b_\beta)$ and $(\mathcal{V}'_{\Gamma^{n-1}}, \mathcal{U}'_{\Gamma^\beta}, a'_\beta, b'_\beta)$ of $Q^{n,\beta}$ is a pair (ξ, η_β) , where $\xi \in \text{Hom}_{\text{Qui}_{\Gamma^{n-1}}}(\mathcal{V}_{\Gamma^{n-1}}, \mathcal{V}'_{\Gamma^{n-1}})$, $\eta_\beta \in \text{Hom}_{\text{Qui}_{\Gamma^\beta}}(\mathcal{U}_{\Gamma^\beta}, \mathcal{U}'_{\Gamma^\beta})$, such that $a'_\beta \bar{\psi}^{(0)}(\xi) = i_\beta(\eta_\beta) a_\beta$, and $\bar{\psi}^{(0)}(\xi) b_\beta = b'_\beta i_\beta(\eta_\beta)$.

Now we assume that the index β is not fixed, but range over the set $I_n(\Gamma)$.

Let Q^n be the category of collections $\{\mathcal{V}_{\Gamma^{n-1}}, \mathcal{U}_{\Gamma^\beta}, a_\beta, b_\beta \mid \beta \in I_n(\Gamma)\}$, where $\mathcal{V}_{\Gamma^{n-1}}$ is a quiver of Γ^{n-1} , $\mathcal{U}_{\Gamma^\beta}$ is a quiver of Γ^β , a_β and b_β are morphisms of $\Gamma^{n,\beta}$ -quivers, $a_\beta : \bar{\psi}_\beta^{(0)}(\mathcal{V}_{\Gamma^{n-1}}) \rightarrow i_\beta(\mathcal{U}_{\Gamma^\beta})$ and $b_\beta : i_\beta(\mathcal{U}_{\Gamma^\beta}) \rightarrow \bar{\psi}_\beta^{(0)}(\mathcal{V}_{\Gamma^{n-1}})$, such that the composition $ba : \bar{\psi}_\beta^{(0)}(\mathcal{V}_{\Gamma^{n-1}}) \rightarrow \bar{\psi}_\beta^{(0)}(\mathcal{V}_{\Gamma^{n-1}})$ is equal to $\bar{T}_\beta$. A morphism between objects $\{\mathcal{V}_{\Gamma^{n-1}}, \mathcal{U}_{\Gamma^\beta}, a_\beta, b_\beta \mid \beta \in I_n(\Gamma)\}$ and $\{\mathcal{V}'_{\Gamma^{n-1}}, \mathcal{U}'_{\Gamma^\beta}, a'_\beta, b'_\beta \mid \beta \in I_n(\Gamma)\}$ is a collection $\{\xi, \eta_\beta \mid \beta \in I_n(\Gamma)\}$, where $\xi \in \text{Hom}_{\text{Qui}_{\Gamma^{n-1}}}(\mathcal{V}_{\Gamma^{n-1}}, \mathcal{V}'_{\Gamma^{n-1}})$, and $\eta_\beta \in \text{Hom}_{\text{Qui}_{\Gamma^\beta}}(\mathcal{U}_{\Gamma^\beta}, \mathcal{U}'_{\Gamma^\beta})$, such that $a'_\beta \bar{\psi}^{(0)}(\xi) = i_\beta(\eta_\beta) a_\beta$, and $\bar{\psi}^{(0)}(\xi) b_\beta = b'_\beta i_\beta(\eta_\beta)$ for any $\beta \in I_n(\Gamma)$.

For any $\beta \in I_n(\Gamma)$, $A \in Q^{n,\beta}$, $A = (\mathcal{V}_{\Gamma^{n-1}}, \mathcal{U}_{\Gamma^\beta}, a_\beta, b_\beta)$, denote by $\nu(A)$ the Γ^{n-1} -quiver $\mathcal{V}_{\Gamma^{n-1}}$. Define the fibered product $\tilde{Q}^n$ of the categories $Q^{n,\beta}$,

$$(7.31) \quad \tilde{Q}^n = Q^{n,\beta_1} \times_{\text{Qui}_{\Gamma^{n-1}}} Q^{n,\beta_2} \times_{\text{Qui}_{\Gamma^{n-1}}} \cdots \times_{\text{Qui}_{\Gamma^{n-1}}} Q^{n,\beta_m},$$

where the product is taken over all $\beta_k \in I_n(\Gamma)$. An object of the category $\tilde{Q}^n$ is the collection of quadruples $A_1 \in Q^{n,\beta_1}, A_2 \in Q^{n,\beta_2}, \dots, A_m \in Q^{n,\beta_m}$, such that $\nu(A_1) = \nu(A_2) = \dots = \nu(A_m)$.

Lemma 7.2. *The categories Q^n and $\tilde{Q}^n$ are equivalent.*

□

Define a functor $\mu : \text{Qui}_{\Gamma^n} \rightarrow Q^n$ as follows. For any $\mathcal{V}_{\Gamma^n} \in \text{Qui}_{\Gamma^n}$ we set

$$\mu(\mathcal{V}_{\Gamma^n}) = \{\mathbb{J}_{n,n-1}^* \mathcal{V}_{G^n}, \overline{\psi}_{\beta}^{(-1)}(\mathcal{V}_{\Gamma^n}), \overline{a}_{\beta}, \overline{b}_{\beta} \mid \beta \in I(\Gamma^n)\}.$$

Lemma 7.3. *The functor $\mu : \text{Qui}_{\Gamma^n} \rightarrow Q^n$ is an equivalence of categories.*

Proof. In order to prove that μ is a functor, one should check the commutativity of certain diagrams of maps between quivers. A direct check shows that the relations (ii)-(v) of Section 3.2 imply the required equalities. The functor μ is an equivalence of categories, since the spaces V_{α} and the maps $A_{\alpha,\beta}$ of the quiver $\mathcal{V}_{\Gamma^n}$ can be reconstructed from $\mu(\mathcal{V}_{\Gamma^n})$.

7.6.4. Gluing data for quiver $\mathcal{D}_{X^{n,\beta}}$ -modules. Fix a vertex $\beta \in I_n(\Gamma)$. Choose a generic constant vector field ξ_{β} such that $\langle \xi_{\beta}, \overline{f}_{\beta} \rangle = 1$ and set, following the notations of the previous section, $f = \overline{f}_{\beta}, \xi = \xi_{\beta}, \theta = \overline{f}_{\beta} \xi_{\beta}$. We apply the construction of Section 7.6.1 to $Y = X^{n,\beta}, Z = Y \cap H, \theta$, and the D_Y -module $E_Y^n \mathcal{V}_{\Gamma^n} = E^n \mathcal{V}_{\Gamma^n}(Y)$, which is the restriction of the D_{X^n} -module $E^n \mathcal{V}_{\Gamma^n}$ to the open subset $Y = X^{n,\beta}$. We denote by i_{β} the closed embedding of X_{β} to Z .

Denote by $\overline{V}_{\pm}$ the following finite-dimensional subspaces of the space of global sections of the D_Y -module $E_Y^n \mathcal{V}_{\Gamma^n}$:

$$\overline{V}_{+} = \bigoplus_{\alpha \in I(\Gamma^{n-1})} \overline{\Omega}_{\alpha} \otimes V_{\alpha}, \quad \overline{V}_{-} = \overline{\Omega}_{\beta} \otimes V_{\beta}.$$

Let $F^k(\mathcal{D}_Y)$ be the filtration of $\mathcal{D}_Y$, associated to the sheaf of ideals, generated by $f = \overline{f}_{\beta}$ as in Section 7.6.1. Set

$$(7.32) \quad F^k(E_Y^n \mathcal{V}_{\Gamma^n}) = \begin{cases} F^k(\mathcal{D}_Y) \cdot \overline{V}_{+}, & k \geq 0 \\ F^{k+1}(\mathcal{D}_Y) \overline{V}_{-} + F^k(\mathcal{D}_Y) \cdot \overline{V}_{+}, & k < 0. \end{cases}$$

Formula (7.32) determines a filtration of the sheaf $E^n \mathcal{V}_{\Gamma^n}$.

Lemma 7.4.

- (i) *The filtration (7.32) is good with respect to $F(\mathcal{D}_Y)$.*
- (ii) *The sheaf $F^0(E_Y^n \mathcal{V}_{\Gamma^n})/F^1(E_Y^n \mathcal{V}_{\Gamma^n})$ of $\mathcal{D}_Z$ -modules is isomorphic to the $\mathcal{D}_Z$ -module $E^{n-1}\psi_{\beta}^{(0)}(\mathcal{V}_{\Gamma^n})$.*
- (iii) *The sheaf $F^{-1}(E_Y^n \mathcal{V}_{\Gamma^n})/F^0(E_Y^n \mathcal{V}_{\Gamma^n})$ of $\mathcal{D}_Z$ -modules is isomorphic to the quiver $\mathcal{D}_Z$ -module $E^{n-1}\psi_{\beta}^{(-1)}(\mathcal{V}_{\Gamma^n})$.*
- (iv) *$\mathcal{D}_Z$ -module maps $\text{gr } \overline{f}_{\beta} : F^{-1}(E_Y^n \mathcal{V}_{\Gamma^n})/F^0(E_Y^n \mathcal{V}_{\Gamma^n}) \rightarrow F^0(E_Y^n \mathcal{V}_{\Gamma^n})/F^1(E_Y^n \mathcal{V}_{\Gamma^n})$ and $\text{gr } \xi_{\beta} : F^0(E_Y^n \mathcal{V}_{\Gamma^n})/F^1(E_Y^n \mathcal{V}_{\Gamma^n}) \rightarrow F^{-1}(E_Y^n \mathcal{V}_{\Gamma^n})/F^0(E_Y^n \mathcal{V}_{\Gamma^n})$ are isomorphic to corresponding $\mathcal{D}_Z$ -module maps $E^{n-1}\overline{b}_{\beta} : E^{n-1}\psi_{\beta}^{(-1)}(\mathcal{V}_{\Gamma^n}) \rightarrow E^{n-1}\psi_{\beta}^{(0)}(\mathcal{V}_{\Gamma^n})$ and $E^{n-1}\overline{a}_{\beta} : E^{n-1}\psi_{\beta}^{(0)}(\mathcal{V}_{\Gamma^n}) \rightarrow E^{n-1}\psi_{\beta}^{(-1)}(\mathcal{V}_{\Gamma^n})$.*

- (v) *The minimal polynomial of the map $\text{gr } \theta_\beta$ coincides with minimal polynomial of the map $\bar{b}_\beta \bar{a}_\beta : \oplus_{\alpha \in I_{n-1}(\Gamma)} V_\alpha \rightarrow \oplus_{\alpha \in I_{n-1}(\Gamma)} V_\alpha$.*

The proof of statements (ii)-(v) is based on Proposition 4.2 and consists of direct calculations with quiver $\mathcal{D}_Y$ -modules on affine open sets $U_{\mathcal{F}_{n-1}}^\beta$. $\square$

Let $\bar{G}_\beta$ be a non-resonant section of $\mathbb{C}$.

Denote by $\widetilde{\mathcal{M}}_{X^n}^{hrs} = \widetilde{\mathcal{M}}_{X^n}^{hol} \cap \mathcal{M}_{X^n}^{hrs}$ the category of holonomic $\mathcal{D}_{X^n}$ -modules with regular singularities, smooth along the strata of X^n .

Denote by $\widetilde{\mathcal{N}}^{n,\beta}(\bar{G}_\beta)$ the category of quadruples (M, N, c_β, d_β) where $M \in \widetilde{\mathcal{M}}_{X^{n-1}}^{hrs}$, $N \in \widetilde{\mathcal{M}}_{X_\beta}^{hrs}$, $c_\beta : \Psi_{\bar{G}_\beta}^{(0)}(M) \rightarrow (i_\beta)_* N$ and $d_\beta : (i_\beta)_* N \rightarrow \Psi_{\bar{G}_\beta}^{(0)}(M)$ are $\mathcal{D}_Z$ -module morphisms such that $d_\beta c_\beta = \text{gr } \theta_\beta : \Psi_{\bar{G}_\beta}^{(0)}(M) \rightarrow \Psi_{\bar{G}_\beta}^{(0)}(M)$ and all eigenvalues of $d_\beta c_\beta$ are in the set $\bar{G}_\beta$. The gluing construction, see Section 7.6.1 says that the functor $H(\bar{G}_\beta) : \widetilde{\mathcal{M}}_{X^{n,\beta}}^{hrs} \rightarrow \widetilde{\mathcal{N}}^{n,\beta}(\bar{G}_\beta)$, attaching to every $M \in \widetilde{\mathcal{M}}_{X^{n,\beta}}^{hrs}$ the quadruple $(M|_{X^{n-1}}, (i_\beta)^* \Psi_{\bar{G}_\beta}^{(-1)}(M), c_\beta = \text{gr } \xi_\beta, d_\beta = \text{gr } \bar{f}_\beta)$, establishes an equivalence of the two categories. For any object $\tilde{N} \in \widetilde{\mathcal{N}}^{n,\beta}(\bar{G}_\beta)$, $\tilde{N} = (M, N, c_\beta, d_\beta)$ denote by $\tilde{\nu}_\beta(\tilde{N})$ the $\mathcal{D}_{X^{n-1}}$ -module M .

Let $G_\beta \subset \bar{G}_\beta$ be a non-resonant set.

Consider the category $Q^{n,\beta}(G_\beta)$ of the quadruples $Q = (\mathcal{V}_{\Gamma^{n-1}}, \mathcal{U}_{\Gamma^\beta}, a_\beta, b_\beta)$ in $Q^{n,\beta}$, such that the eigenvalues of the operator (7.30) are contained in G_β . Then an assignment

$$F^{n,\beta} : Q \mapsto (E^{n-1}(\mathcal{V}_{\Gamma^{n-1}}), E^0(\mathcal{U}_{\Gamma^\beta}), E^{n-1}(a_\beta), E^{n-1}(b_\beta)),$$

defines a functor $F^{n,\beta} : Q^{n,\beta}(G_\beta) \rightarrow \widetilde{\mathcal{N}}^{n,\beta}(G_\beta)$.

Consider the category $Qui_{\Gamma^n}(G_\beta) \subset Qui_{\Gamma^n}$ of level n quivers $\mathcal{V}_{\Gamma^n}$, such that the eigenvalues of operator T_β , defined in (3.5), are contained in the set G_β . By Lemma 7.4, we have a commutative diagram of functors:

$$(7.33) \quad \begin{array}{ccc} Qui_{\Gamma^n}(G_\beta) & \xrightarrow{E^n(X^{n,\beta})} & \widetilde{\mathcal{M}}_{X^{n,\beta}}^{hrs} \\ \downarrow \mu_\beta & & \downarrow H(\bar{G}_\beta) \\ Q^{n,\beta}(G_\beta) & \xrightarrow{F^{n,\beta}} & \widetilde{\mathcal{N}}^{n,\beta}(\bar{G}_\beta). \end{array}$$

The vertical lines in the diagram (7.33) represent equivalences of categories.

7.6.5. The induction. Suppose that for every vertex $\beta \in I_n(\Gamma)$ there are chosen: (a) a non-resonant section $\bar{G}_\beta$ of $\mathbb{C}$; (b) a non-resonant set $G_\beta \subset \bar{G}_\beta$; (c) a generic affine function $\bar{f}_\beta \in F_\beta$ and a generic constant vector field ξ_β such that $\langle \xi_\beta, \bar{f}_\beta \rangle = 1$. For every $\beta \in I_n(\Gamma)$ denote by Z_β the intersection $X^{n,\beta} \cap \{\bar{f}_\beta = 0\}$, by i_β the closed embedding of X_β to Z_β .

Consider the category $\widetilde{\mathcal{N}}^n(\{\bar{G}_\beta\})$, which consists of the collections $\{M, N_\beta, a_\beta, b_\beta \mid \beta \in I_n(\Gamma)\}$, where M is a holonomic $\mathcal{D}_{X^{n-1}}$ -module with regular singularities, N_β is a holonomic $\mathcal{D}_{X_\beta}$ -module with regular singularities, $a : \Psi_{\bar{G}_\beta}^{(0)}(M) \rightarrow (i_\beta)_*(N)$ and $b : (i_\beta)_*(N) \rightarrow \Psi_{\bar{G}_\beta}^{(0)}(M)$ are $\mathcal{D}_{Z_\beta}$ -module maps such that $ba = \text{gr } \theta_\beta : \Psi_{\bar{G}_\beta}^{(0)}(M) \rightarrow \Psi_{\bar{G}_\beta}^{(0)}(M)$ and all eigenvalues of ba are in the set $\bar{G}_\beta$. The category $\widetilde{\mathcal{N}}^n(\{\bar{G}_\beta\})$ is equivalent to the fibered product

of the categories $\tilde{\mathcal{N}}^{n,\beta}(\overline{G}_\beta)$. This fibered product consists of the collections $(\tilde{N}_{\beta_1}, \dots, \tilde{N}_{\beta_m})$, where $\tilde{N}_{\beta_k} \in \tilde{\mathcal{N}}^{n,\beta_k}(\overline{G}_{\beta_k})$ and $\tilde{\nu}_{\beta_1}(\tilde{N}_{\beta_1}) = \dots = \tilde{\nu}_{\beta_m}(\tilde{N}_{\beta_m})$.

Denote by $Qui_{\Gamma^n}(\{G_\beta\}) \subset Qui_{\Gamma^n}$ the category of level n quivers $\mathcal{V}_{\Gamma^n}$, such that the eigenvalues of operator T_β , defined in (3.5), are contained in the set G_β for every $\beta \in I_n(\Gamma)$.

Denote by $Q^n(\{G_\beta\})$ the category of collections $Q = \{\mathcal{V}_{\Gamma^{n-1}}, \mathcal{U}_{\Gamma^\beta}, a_\beta, b_\beta \mid \beta \in I_n(\Gamma)\}$ in Q^n , such that the eigenvalues of the operator $\overline{T}_\beta$ are contained in G_β for every $\beta \in I_n(\Gamma)$.

The assignment

$$F^n : Q \mapsto \{E^{n-1}(\mathcal{V}_{\Gamma^{n-1}}), E^0(\mathcal{U}_{\Gamma^\beta}), E^{n-1}(a_\beta), E^{n-1}(b_\beta) \mid \beta \in I_n(\Gamma)\},$$

defines a functor $F^n : Q^n(\{G_\beta\}) \rightarrow \tilde{\mathcal{N}}^n(\{G_\beta\})$. We have a commutative diagram

$$(7.34) \quad \begin{array}{ccc} Qui_{\Gamma^n}(\{G_\beta\}) & \xrightarrow{E^n} & \tilde{\mathcal{M}}_{X^n}^{hrs} \\ \downarrow \mu & & \downarrow H(\{\overline{G}_\beta\}) \\ Q^n(\{G_\beta\}) & \xrightarrow{F^n} & \tilde{\mathcal{N}}^n(\{\overline{G}_\beta\}). \end{array}$$

The functor $H(\{\overline{G}_\beta\})$ attaches to any $M \in \tilde{\mathcal{M}}_{X^n}^{hrs}$ the collection $(M_{X^{n-1}}, (i_\beta)^* \Psi_{\overline{G}_\beta}^{(-1)}(M), c_\beta = \text{gr } \xi_\beta, d_\beta = \text{gr } \overline{f}_\beta \mid \beta \in I_n(\Gamma))$. This functor is equivalent to the fibered product of functors $H(\overline{G}_\beta)$ and thus is an equivalence of categories. We know also that the functor μ is an equivalence of categories. Thus if the functor F^n is full, then the functor E^n is full as well.

Consider the category $Qui_{\Gamma^n, \{G_\alpha, \tilde{G}_\beta, G_{\gamma, \delta}\}}$ associated with a collection of non-resonant sets $\{G_\alpha\}$, $\alpha \in I(\Gamma^n) \setminus \emptyset$, and a collection of weakly non-resonant sets $\tilde{G}_\beta$, $\beta \in I(\Gamma^n) \setminus \emptyset$, and $\{G_{\gamma, \delta}\}$, where $(\gamma, \delta) \in E(\Gamma^{n+1})$, $\gamma \succ \delta$. For the same collection of sets $\{G_\alpha, \tilde{G}_\beta, G_{\gamma, \delta}\}$ denote by $Q_{\{G_\alpha, \tilde{G}_\beta, G_{\gamma, \delta}\}}^n$ the category of sequences $Q = \{\mathcal{V}_{\Gamma^{n-1}}, \mathcal{U}_{\Gamma^\beta}, a_\beta, b_\beta \mid \beta \in I_n(\Gamma)\}$ in Q^n , such that

- (a) for any $\beta \in I_n(\Gamma)$ the eigenvalues of the operator T_β , defined in (3.5), are contained in the set G_β ,
- (b) the level $n-1$ quiver $\mathcal{V}_{\Gamma^{n-1}}$ is in the category $Qui_{\Gamma^{n-1}, \{G_\alpha, \tilde{G}_\beta, G_{\gamma, \delta}\}}$,
- (c) for any $\beta \in I_n(\Gamma)$ the (level zero) Γ^β -quiver $\mathcal{U}_{\Gamma^\beta} = \{U_\emptyset, C_\emptyset^\gamma\}$ is in the category $Qui_{\Gamma^\beta, \{\tilde{G}_\beta, G_{\beta, \gamma}\}}$, that is, the eigenvalues of the operator $C_\emptyset^\gamma$ lie in the set $G_{\beta, i_\beta(\gamma)}$ and the eigenvalues of the operator $\sum_\gamma A_\emptyset^\gamma$ lie in the set $\tilde{G}_\beta$.

The functor μ establishes an equivalence of categories $Qui_{\Gamma^n, \{G_\alpha, \tilde{G}_\beta, G_{\gamma, \delta}\}}$ and $Q_{\{G_\alpha, \tilde{G}_\beta, G_{\gamma, \delta}\}}^n$.

Suppose that the statement (ii) of Proposition 4.3 is valid for quivers of level less than n . Then the restriction of the functor E^{n-1} to the category $Qui_{\Gamma^{n-1}, \{G_\alpha, \tilde{G}_\beta, G_{\gamma, \delta}\}}$ is full and faithful. Proposition 4.4 implies that the restriction of the functor E^0 to the category $Qui_{\Gamma^\beta, \{\tilde{G}_\beta, G_{\beta, \gamma}\}}$ is full and faithful. Thus the restriction of the functor F^n to the category $Q_{\{G_\alpha, \tilde{G}_\beta, G_{\gamma, \delta}\}}^n$ is full. This proves that the restriction of the functor E^n to the category $Qui_{\Gamma^n, \{G_\alpha, \tilde{G}_\beta, G_{\gamma, \delta}\}}$ is full. $\square$

7.7. Proof of Theorem 4.6. In this section we use the following notion. Let $\tilde{Y}$ be an affine nonsingular variety. Let $z_1, \dots, z_k$ be regular functions on $\tilde{Y}$. Assume that $\tilde{Z} \subset \tilde{Y}$ defined by equations $z_1 = 0, \dots, z_k = 0$ is nonsingular. Let N be a $\mathcal{D}_{\tilde{Y}}$ -module. Set $N^{\tilde{Z}} = \{n \in N \mid z_j^p n = 0 \text{ for } j = 1, \dots, k \text{ and } p \in \mathbb{N}, p \gg 1\}$. Then $N^{\tilde{Z}}$ is the maximal submodule of M with support in $\tilde{Z}$.

Keep the notations of Section 7.6. Let $\mathcal{V}_{\Gamma^{n-1}} = \{V_\alpha, A_{\alpha_1, \alpha_2}, A_{\alpha_1}^{\alpha_2}\}$ be a Γ^{n-1} -quiver. Fix a vertex $\beta \in I_n(\Gamma)$, a level $n-1$ vertex framing $\mathcal{F}_{n-1}$, and a generic affine function $\bar{f}_\beta \in F_\beta$. They define open affine subsets $U_{\mathcal{F}_{n-1}} \subset X^{n-1}$, see (2.5), and $U = U_{\mathcal{F}_{n-1}^\beta} \subset X^{n, \beta}$, see (7.29). Then $X_\beta \subset U$ is a closed nonsingular subvariety of U . Denote by M the restriction to $X^{n, \beta}$ of the sheaf $j_{n, n-1, *}(E^{n-1} \mathcal{V}_{\Gamma^{n-1}})$ of $\mathcal{D}_{X^n}$ -modules and by M_U the restriction of M to U .

Let $H = \{\bar{f}_\beta = 0\}$, $H_U = U \cap H \subset Z = X^{n, \beta} \cap H$, and $\tilde{U} = U \setminus H_U$. Let ξ be a constant vector field on X , such that $\xi(d\bar{f}_\beta) = 1$. Denote by θ the vector field $\theta = f\xi$. Let $\bar{G}_\beta$ be a non-resonant section of $\mathbb{C}$ and $G_\beta \subset \bar{G}_\beta$ a non-resonant set. Let $\Psi_{\bar{G}_\beta}^{(k)}(M)$ be the graded factors of the Kashiwara-Malgrange filtration of M , associated with the ideal, generated by the function $\bar{f}_\beta$, and the set $\bar{G}_\beta$.

Lemma 7.5.

- (i) *The image of the map $ba : \Psi_{\bar{G}_\beta}^{(0)}(M) \rightarrow \Psi_{\bar{G}_\beta}^{(0)}(M)$ has support in $X_\beta \subset H_\beta$,*

$$\text{Im } ba \subset \left(\Psi_{\bar{G}_\beta}^{(0)}(M) \right)^{X_\beta}.$$

- (ii) *The $\mathcal{D}_Z$ -module $\Psi_{\bar{G}_\beta}^{(-1)}(M)$ has support in X_β ,*

$$\Psi_{\bar{G}_\beta}^{(-1)}(M) = \left(\Psi_{\bar{G}_\beta}^{(-1)}(M) \right)^{X_\beta}.$$

- (iii) *The gluing data for $\mathcal{D}_U$ -module M are isomorphic to*

$$\left(M|_{\tilde{U}}, \left(\Psi_{\bar{G}_\beta}^{(0)}(M) \right)^{X_\beta}, a, b \right),$$

where $b : \left(\Psi_{\bar{G}_\beta}^{(0)}(M) \right)^{X_\beta} \rightarrow \Psi_{\bar{G}_\beta}^{(0)}(M)$ is the canonical inclusion and $a : \Psi_{\bar{G}_\beta}^{(0)}(M) \rightarrow \left(\Psi_{\bar{G}_\beta}^{(0)}(M) \right)^{X_\beta}$ is the restriction of the monodromy map ba to $\left(\Psi_{\bar{G}_\beta}^{(0)}(M) \right)^{X_\beta}$.

Proof. The statement (i) follows from statements (ii) and (iv) of Lemma 7.4, Section 7.6.4. Let us prove statement (ii). Note that since $\bar{f}_\beta$ is generic, for any $\alpha \in I(\Gamma^{n-1})$ there exists a vector field $\eta \in T_\alpha$, such that $\langle \eta, \bar{f}_\beta \rangle \neq 0$. This implies that in the open set U any element in $F_{\bar{G}_\beta}^{-1}(M_U)$ can be presented as a fraction $\frac{m}{\bar{f}_\beta}$, where $m \in F_{\bar{G}_\beta}^0(M_U)$. Indeed, for any element $n \in M_U$, we have the inclusion $\bar{f}_\beta^k n \in F_{\bar{G}_\beta}^0(M_U)$ for sufficiently big k , so the elements of $F_{\bar{G}_\beta}^0(M_U)$ generate M_U as a $D_U[\bar{f}_\beta^{-1}]$ -module, that is, generate M_U as a $\mathcal{D}_U$ module in the open set $\bar{f}_\beta \neq 0$. Then any element $n \in F_{\bar{G}_\beta}^{-1}(M_U)$ can be

written as $n = \overline{f}_\beta^{-1} m'$, where $m' \in F_{\overline{G}_\beta}^0(M_U)$, or as $n = \nu m'$, where $m' \in F_{\overline{G}_\beta}^0(M_U)$, and ν is a vector field with constant coefficients, such that $\langle \nu, \overline{f}_\beta \rangle \neq 0$. In the last case we can find a constant field ν' , such that $\langle \nu', \overline{f}_\beta \rangle = \langle \nu, \overline{f}_\beta \rangle$ and the element $n' = \nu' m' \in F_G^{-1}(M_U)$ satisfies the following properties. It contains a smaller number of derivatives in its expression in terms of vectors $\overline{\omega}_\alpha \otimes v_\alpha$ and there is an inclusion $n - n' \in F_{\overline{G}_\beta}^0(M_U)$. By induction we prove that there exists $m \in F_{\overline{G}_\beta}^0(M_U)$, such that $n = f^{-1} m$. Let f' be another generic affine function, vanishing on X_β . By the same arguments we show that there exists an element $\tilde{m} \in F_{\overline{G}_\beta}^0(M_U)$, such that $n = f'^{-k} \tilde{m}$ for some integer $k > 0$. Put $x = \overline{f}_\beta - f'$. The equality

$$n = \frac{m}{\overline{f}_\beta} = \frac{\tilde{m}}{(\overline{f}_\beta + x)^k}$$

implies that $x^k n \in F_{\overline{G}_\beta}^0(M_U)$, that is

$$\overline{x}^k n = 0 \quad \text{in} \quad \Psi_{\overline{G}_\beta}^{-1}(M_U),$$

where $\overline{x} = x|_{H_U}$. This proves statement (ii) of the lemma.

Denote by $\tilde{j}$ the inclusion $\tilde{j} : \tilde{U} \hookrightarrow U$ and by $\widetilde{M}$ the $\mathcal{D}_{\tilde{U}}$ -module $\widetilde{M} = M|_{\tilde{U}} = E^{n-1} \mathcal{V}_{\Gamma^{n-1}}(\tilde{U})$. Let M' be the $\mathcal{D}_U$ -module, given by the gluing data

$$\left(\widetilde{M}, \left(\Psi_{\overline{G}_\beta}^{(0)}(M) \right)^{X_\beta}, a, b \right),$$

where the maps a and b are described in statement (iii) of the lemma. For any $\mathcal{D}_U$ -module N , whose gluing data satisfy the condition that $\Psi_{\overline{G}_\beta}^{(-1)}(N)$ is supported at X_β , we have the equality

$$(7.35) \quad \text{Hom}_{\mathcal{D}_{\tilde{U}}}(\tilde{j}^*(N), \widetilde{M}) \simeq \text{Hom}_{\mathcal{D}_U}((N), M').$$

Moreover, M' is the unique $\mathcal{D}_U$ -module, satisfying conditions (7.35). Denote by j_β the inclusion $j_\beta : U_{\mathcal{F}_{n-1}} \hookrightarrow U$. Since $\overline{f}_\beta$ is generic, we have the equality

$$\text{Hom}_{\mathcal{D}_{\tilde{U}}}(\tilde{j}^*(N), \widetilde{M}) \simeq \text{Hom}_{\mathcal{D}_{U_{\mathcal{F}_{n-1}}}}(j_\beta^*(N), E^{n-1} \mathcal{V}_{\Gamma^{n-1}}|_{U_{\mathcal{F}_{n-1}}}).$$

This equality, relation (7.35) and statement (ii) of lemma imply statement (ii). $\square$

Lemma 7.6. *Let $E^n \mathcal{V}_{\Gamma^n}$ be the $\mathcal{D}_{X^n}$ -module, attached to a Γ^n -quiver $\mathcal{V}_{\Gamma^n}$. Let $\beta \in I(\Gamma^n)$, $l(\beta) = n$, be a vertex such that the operator S_β , defined in (3.4) has no positive integer eigenvalues. Then the $\mathcal{D}_{X^n}$ -module $(E^n \mathcal{V}_{\Gamma^n})^{X_\beta}$ is generated by vectors of the vector space $V'_\beta \otimes \overline{\Omega}_\beta$, where*

$$(7.36) \quad V'_\beta = \bigcap_{\alpha, \alpha \succ \beta} \text{Ker } A_{\alpha, \beta} \subset V_\beta.$$

The proof of the lemma goes in two steps. First, we show that the lemma holds in a special case of a quiver $\mathcal{D}$ -module supported at a flag of a normal crossing arrangement and then show how to reduce the general case to the special.

Denote by $\mathcal{C}_{n,N}$ the arrangement of n hyperplanes $\{z_j = 0\}$, $j = 1, \dots, n$, in the affine space $\mathbb{C}^N$ with coordinates z_k , $k = 1, \dots, N$. Let $\bar{\Gamma}_n$ be the graph of the arrangement $\mathcal{C}_{n,N}$. The vertices of $\bar{\Gamma}_n$ are labeled by subsets $J \subset \{1, \dots, n\}$. We say that a $\bar{\Gamma}_n$ -quiver $\mathcal{V}$ is supported at the flag $\emptyset, \{1\}, \{1, 2\}, \dots, \{1, 2, \dots, n\}$ of vertices of $\bar{\Gamma}_n$ if the vector spaces V_J of the quiver are nonzero only for J equal to one of the sets $J_0 = \emptyset$, $J_1 = \{1\}$, $J_2 = \{1, 2\}$, $\dots$, $J_n = \{1, 2, \dots, n\}$.

Let $\mathcal{V}$ be a $\bar{\Gamma}_n$ -quiver supported at the flag $\{1\}, \{1, 2\}, \dots, \{1, 2, \dots, n\}$. Choose basic exterior forms $\bar{\omega}_{J_k} \in \bar{\Omega}_{J_k}$, $\bar{\omega}_{J_k} = dz_{k+1} \wedge \dots \wedge dz_N$ and denote $\bar{v}_{J_k} = \bar{\omega}_{J_k} \otimes v_{J_k}$. Then the defining relations (4.25) and (4.26) for the quiver $\mathcal{D}_X$ -module $E\mathcal{V}$ look as follows:

$$(7.37) \quad \begin{cases} z_i \bar{v}_{J_k} = 0, & i < k, \\ z_i \bar{v}_{J_k} = \frac{\bar{\omega}_{J_k}}{A_{J_{k-1}, J_k}} v_{J_k}, & i = k, \end{cases} \quad \begin{cases} \partial_{z_i} \bar{v}_{J_k} = 0, & i > k+1, \\ \partial_{z_i} \bar{v}_{J_k} = \frac{\bar{\omega}_{J_k}}{A_{J_{k+1}, J_k}} v_{J_k}, & i = k+1. \end{cases}$$

Denote by $\hat{Z}$ the stratum X_{J_n} , given by equations $z_1 = \dots = z_n = 0$. Suppose $x \in (E\mathcal{V})^{\hat{Z}}$. We can assume that

$$(7.38) \quad x = P(z_{n+1}, \dots, z_N) \partial_{z_1}^{k_1} \dots \partial_{z_n}^{k_n} (\bar{v}_{J_n}),$$

where $v_{J_n} \in V_{\{1, 2, \dots, n\}}$. We claim that $x \in (E\mathcal{V})^{\hat{Z}} \Rightarrow v_{J_n} \in \text{Ker } A_{J_{n-1}, n}$ if the operator $S_{J_n} = A_{J_{n-1}, n} A_{J_{n-1}, n} : V_{J_n} \rightarrow V_{J_n}$ has no positive integer eigenvalues.

Suppose we prove this claim for x , such that $k_1 + \dots + k_n < k$, see (7.38). Let us prove the claim for $k_1 + \dots + k_n = k$. Since $z_i(\bar{v}_{J_n}) = 0$ for $i = 1, \dots, n-1$, the condition $z_i^M x = 0$ for $M > 0$ implies $k_i = 0$ for $i = 1, \dots, n-1$. Let $x = P(z_{n+1}, \dots, z_N) \partial_{z_n}^{k_n} (\bar{v}_{J_n})$. If $k_n = 0$, then the condition $z_n^M x = 0$ implies $x \in \text{Ker } A_{J_{n-1}, n}$ by (7.37). Let $k_n \neq 0$. Then

$$z_n x = P(z_{n+1}, \dots, z_N) \partial_{z_n}^{k_n-1} \overline{(A_{J_{n-1}, n} A_{J_{n-1}, n} - k_n)} v_{J_n}.$$

The condition $z_n^M x = 0$ for some $M > 0$ implies by assumption of induction that $(A_{J_{n-1}, n} A_{J_{n-1}, n} - k_n) v_{J_n} \in \text{Ker } A_{J_{n-1}, n}$, that is

$$(7.39) \quad A_{J_{n-1}, n} (A_{J_{n-1}, n} A_{J_{n-1}, n} - k_n) v_{J_n} = (A_{J_{n-1}, n} A_{J_{n-1}, n} - k_n) A_{J_{n-1}, n} v_{J_n} = 0.$$

The operator $A_{J_{n-1}, n} A_{J_{n-1}, n}$, as well as the operator $S_{J_n} = A_{J_{n-1}, n} A_{J_{n-1}, n}$ has no positive integer eigenvalues, so the operator $(A_{J_{n-1}, n} A_{J_{n-1}, n} - k_n)$ is invertible and (7.39) implies $A_{J_{n-1}, n} v_{J_n} = 0$.

The passage from the general case to the special case of normal crossing, described above, is based on the following observation.

Keep the notations of Section 7.6.1. Let Y be a smooth algebraic complex variety, let $Z \subset Y$ be a smooth subvariety of Y and let $I \subset \mathcal{O}_Y$ be a sheaf of ideals of the functions vanishing on Z . Let M be a holonomic $\mathcal{D}_Y$ -module and let $F(M)$ be a good filtration with respect to I . Note that in general $\cap_n F^n(M) \neq 0$.

Lemma 7.7. *Let m be a nonzero element of a $\mathcal{D}_Y$ -module M annihilated by some power of the ideal I , $I^k m = 0$. Then there exists $n \in \mathbb{Z}$ such that $m \notin F^n(M)$. Thus the symbol $\text{gr}(m)$ is well defined and is a nonzero element of the associated graded quotient $\text{gr}(M)$.*

The claim follows from condition (ii), Section 7.6.1, for good filtrations and the fact that $F^k(\mathcal{D}_Y) \subset F^0(\mathcal{D}_Y) \cdot I^k$ for all $k > 0$. $\square$

Return to the proof of Lemma 7.6 for a general arrangement $\mathcal{C}$. Let $E^n\mathcal{V}_{\Gamma^n}$ be the $\mathcal{D}_{X^n}$ -module, attached to the Γ^n -quiver $\mathcal{V}_{\Gamma^n}$. Let $\beta \in I(\Gamma^n)$, $l(\beta) = n$, be a vertex such that the operator S_β has no positive integer eigenvalues. Let $z_1, \dots, z_N$ be affine coordinates in $X = \mathbb{C}^N$ such that the closure $\overline{X}_\beta$ of the stratum X_β is given by the equations $z_1 = 0, \dots, z_n = 0$, and the functions $z_1, \dots, z_n$ are generic affine functions in F_β .

Denote by $(E^n\mathcal{V}_{\Gamma^n})^0$ the $\mathcal{D}_{X^n}$ -module, generated by vectors of the vector space $V'_\beta \otimes \overline{\Omega}_\beta$, where the space V'_β is defined in (7.36). Clearly, $(E^n\mathcal{V}_{\Gamma^n})^0 \subset (E^n\mathcal{V}_{\Gamma^n})^{X_\beta}$. We will prove that

$$(7.40) \quad (E^n\mathcal{V}_{\Gamma^n})^0 = (E^n\mathcal{V}_{\Gamma^n})^{X_\beta}$$

under the conditions of Lemma 7.6.

For $k = 1, \dots, n$ let I_k be the left ideal of $\mathcal{O}_X$, generated by $z_1, z_2, \dots, z_k$. Let gr_k denote the graded quotient space with respect to a good filtration, determined by the ideal I_k . Then $\text{gr}_{1, \dots, n}(E^n\mathcal{V}_{\Gamma^n}) := \text{gr}_n \text{gr}_{n-1} \cdots \text{gr}_1(E^n\mathcal{V}_{\Gamma^n})$ is a quiver $\mathcal{D}_Y$ -module, where Y denotes the affine space $\mathbb{C}^N$ stratified by intersections of coordinate hyperplanes $\{z_j = 0\}$, $j = 1, \dots, n$, and the corresponding quiver is supported at the flag $\emptyset, \{1\}, \{1, 2\}, \dots, \{1, 2, \dots, n\}$ of vertices of $\overline{\Gamma}_n$, described above. More precisely, $\text{gr}_{1, \dots, n}(E^n\mathcal{V}_{\Gamma^n})$ is a $\overline{\Gamma}^n$ -quiver $\mathcal{D}_Y$ -module $E\tilde{\mathcal{V}}_{\overline{\Gamma}^n}$, such that the space $\tilde{V}_{J_k}$ of the quiver $\tilde{\mathcal{V}}_{\overline{\Gamma}^n}$ is isomorphic to the direct sum

$$\tilde{V}_{J_k} \simeq \bigoplus_{\alpha, \alpha \succ \beta, l(\alpha)=k} V_\alpha,$$

and the maps $A_{J_k, J_{k \pm 1}}$ are the sums of the corresponding maps A_{α_i, α_j} . In particular, $\tilde{V}_{J_k} \approx V_\beta$ (and $Y_{J_n} \simeq \overline{X}_\beta$), and the map S_{J_n} is isomorphic to the map S_β . Thus we conclude that the conditions of Lemma 7.6 imply the equality

$$(7.41) \quad (\text{gr}_{1, \dots, n}(E^n\mathcal{V}_{\Gamma^n}))^0 = (\text{gr}_{1, \dots, n}(E^n\mathcal{V}_{\Gamma^n}))^{\overline{X}_\beta}.$$

By Lemma 7.7, any element $m \in (E^n\mathcal{V}_{\Gamma^n})^{\overline{X}_\beta}$ has a well defined nonzero symbol $\text{gr}_{1, \dots, n}(m)$. Thus relation (7.41) implies relation (7.40). This proves Lemma 7.6. $\square$.

Return to the proof of Theorem 4.6. Since the operator $\bar{T}_\beta$ for the quiver $E^{n-1}\mathcal{V}_{\Gamma^{n-1}}$ is isomorphic to the operator S_β for the quiver $\psi_\beta^{(0)}(\mathcal{V}_{\Gamma^n})$, the $\mathcal{D}_{H_\beta}$ -module $(\Psi_G^{(0)}(M))^{X_\beta}$ is generated by the space $W_\beta \subset \bigoplus_{\alpha \in I_{n-1}(\Gamma), \alpha \succ \beta} V_\alpha$, which consists of the vectors $w = \sum_{\alpha \in I_{n-1}(\Gamma), \alpha \succ \beta} v_\alpha$. Here v_α are vectors in V_α , such that for any $\delta \in I_{n-2}(\Gamma)$, satisfying the condition $\delta > \beta$, the sum $\sum_{\alpha \in I_{n-1}(\Gamma), \alpha \succ \beta} A_{\delta, \alpha} v_\alpha$ equals zero.

Then statement (iii) of Lemma 7.5 implies that the gluing data for the $\mathcal{D}_{X^n}$ -module $M = j_{n, n-1, *} E^{n-1}\mathcal{V}_{\Gamma^{n-1}}$ coincide with the gluing data of the $\mathcal{D}_{X^n}$ -module $E^n \mathbb{J}_{n, n-1, *} \mathcal{V}_{\Gamma^{n-1}}$. This proves statement (i) of Theorem 4.6. $\square$.

Let us prove statement (ii) of Theorem 4.6.

We need the following general property of adjoint functors. Let $\mathcal{A}$ and $\mathcal{B}$ be additive categories. Let $F : \mathcal{A} \rightarrow \mathcal{B}$ and $G : \mathcal{B} \rightarrow \mathcal{A}$ form a pair of left and right adjoint functors. The adjunction is established by a collection of isomorphisms of adjunction,

$$\alpha_{X, Y} : \text{Hom}_{\mathcal{A}}(X, G(Y)) \xrightarrow{\sim} \text{Hom}_{\mathcal{B}}(F(X), Y), \quad X \in \mathcal{A}, \quad Y \in \mathcal{B}.$$

Let $\sigma_{F,G}(X) : X \rightarrow GF(X)$, $\tau_{F,G}(Y) : FG(Y) \rightarrow Y$, be the corresponding adjunction morphisms, see (3.13). One can see that the adjunction morphisms completely determine the isomorphisms of adjunction. Namely, for any $X \in \mathcal{A}$, $Y \in \mathcal{B}$ and morphisms $\varphi \in \text{Hom}_{\mathcal{A}}(X, G(Y))$ and $\psi \in \text{Hom}_{\mathcal{B}}(F(X), Y)$ we have the equalities

$$(7.42) \quad \alpha_{X,Y}(\varphi) = \tau_{F,G}(Y) \cdot F(\varphi), \quad \alpha_{X,Y}^{-1}(\psi) = G(\psi) \cdot \sigma_{F,G}(X).$$

Statement (ii) of Theorem 4.6 relates the isomorphisms of adjunction of two pairs of adjoint functors. The first pair consists of the functors $\mathbb{J}_{l,k}^* : \text{Qui}_{\Gamma^l} \rightarrow \text{Qui}_{\Gamma^k}$ and $\mathbb{J}_{l,k,*} : \text{Qui}_{\Gamma^k} \rightarrow \text{Qui}_{\Gamma^l}$, where $0 \leq k < l \leq N$. The second pair consists of the functors $j_{l,k}^* : \mathcal{M}_{X^l}^{\text{hol}} \rightarrow \mathcal{M}_{X^k}^{\text{hol}}$ and $j_{l,k,*} : \mathcal{M}_{X^k}^{\text{hol}} \rightarrow \mathcal{M}_{X^l}^{\text{hol}}$.

Relation (7.42) says that it is sufficient to prove that the corresponding adjunction morphisms τ commute with the functor E^k ,

$$(7.43) \quad E^k \tau_{\mathbb{J}_{l,k}^*, \mathbb{J}_{l,k,*}}(\mathcal{V}_{\Gamma^k}) = \tau_{j_{l,k}^*, j_{l,k,*}}(E^k \mathcal{V}_{\Gamma^k})$$

for any strongly non-resonant level k quiver $\mathcal{V}_{\Gamma^k}$. By the constructions of Section 3.4, the morphism $\tau_{\mathbb{J}_{l,k}^*, \mathbb{J}_{l,k,*}} : \mathbb{J}_{l,k}^* \mathbb{J}_{l,k,*}(\mathcal{V}_{\Gamma^k}) \rightarrow (\mathcal{V}_{\Gamma^k})$ is realized as the identity map, $\tau_{\mathbb{J}_{l,k}^*, \mathbb{J}_{l,k,*}}|_{V_\alpha} = \text{Id}_{V_\alpha}$, $l(\alpha) \leq k$. On the other hand, for any $M \in \mathcal{M}_{X^k}^{\text{hol}}$ the map $\tau_{j_{l,k}^*, j_{l,k,*}} : j_{l,k}^* j_{l,k,*}(M) \rightarrow M$ is the identification with M of the restriction to the open set X^k of the direct image $j_{l,k,*}(M)$. Relation (7.43) means that the restriction to X^k of both sides of relation (4.41) is equal to $E^k(\mathcal{V}_{\Gamma^k})$. This is true since all constructions of Section 7.7 do not change quiver $\mathcal{D}$ -modules on the open set X^k . $\square$.

7.8. Proof of Theorem 4.11. Introduce a filtration on the complex $(\mathcal{R}E^0 \mathcal{V}_{\Gamma^0}, d)$ as follows. Set $\deg(D \otimes \xi_1 \wedge \dots \wedge \xi_r \otimes \overline{\omega}_\emptyset \otimes v_\emptyset) = k$ if D is a differential operator of degree $k - r$ and denote by $\mathcal{R}E^0 \mathcal{V}_{\Gamma^0}^{(k)}$ the vector subspace of $\mathcal{R}E^0 \mathcal{V}_{\Gamma^0}$, generated by elements of degree less than or equal to k . We have $d(\mathcal{R}E^0 \mathcal{V}_{\Gamma^0}^{(k)}) \subset \mathcal{R}E^0 \mathcal{V}_{\Gamma^0}^{(k)}$,

$$(7.44) \quad 0 \subset \mathcal{R}E^0 \mathcal{V}_{\Gamma^0}^{(0)} \subset \dots \subset \mathcal{R}E^0 \mathcal{V}_{\Gamma^0}^{(k)} \subset \dots \subset \mathcal{R}E^0 \mathcal{V}_{\Gamma^0},$$

and $\bigcup_{k \geq 0} \mathcal{R}E^0 \mathcal{V}_{\Gamma^0}^{(k)} = \mathcal{R}E^0 \mathcal{V}_{\Gamma^0}$. The image of the differential d in the quotient space $\mathcal{R}E^0 \mathcal{V}_{\Gamma^0}^{(k)} / \mathcal{R}E^0 \mathcal{V}_{\Gamma^0}^{(k+1)}$ coincides with the image of the differential d_1 . The quotient complex $\text{gr } \mathcal{R}E^0 \mathcal{V}_{\Gamma^0}^{(k)} = \mathcal{R}E^0 \mathcal{V}_{\Gamma^0}^{(k)} / \mathcal{R}E^0 \mathcal{V}_{\Gamma^0}^{(k+1)}$ is isomorphic to the tensor product of the free $\mathcal{O}(X^0)$ -module $\mathcal{O}(X^0) \otimes \overline{\Omega}_\emptyset \otimes V_\emptyset$ and the k -th graded component of the Koszul complex $(S(\mathbb{C}^N) \otimes \wedge(\mathbb{C}^N), d_{Kos})$, where $d_{Kos} = \sum_{i=1}^N e_i \otimes \frac{\partial}{\partial e_i}$, for a given basis $e_1, \dots, e_N$ of $\mathbb{C}^N$:

$$\text{gr } \mathcal{R}E^0 \mathcal{V}_{\Gamma^0}^{(k)} \approx \mathcal{O}(X^0) \otimes \overline{\Omega}_\emptyset \otimes V_\emptyset \otimes (S(\mathbb{C}^N) \otimes \wedge(\mathbb{C}^N))^{(k)}.$$

The complex $((S(\mathbb{C}^N) \otimes \wedge(\mathbb{C}^N))^{(k)}, d_{Kos})$ is acyclic for $k > 0$ and consists of a one dimensional space $S^0(\mathbb{C}^N) \otimes \wedge^0(\mathbb{C}^N)$ with the zero differential for $k = 0$. As a consequence, the complex $\text{gr } \mathcal{R}E^0 \mathcal{V}_{\Gamma^0}^{(k)}$ is acyclic for $k > 0$, while $\mathcal{R}E^0 \mathcal{V}_{\Gamma^0}^{(0)}$ consists of the vector space $\mathcal{O}(X^0) \otimes \overline{\Omega}_\emptyset \otimes V_\emptyset \otimes S^0(\mathbb{C}^N) \otimes \wedge^0(\mathbb{C}^N)$ with the zero differential. This implies that the inclusion $i_0 : \mathcal{R}E^0 \mathcal{V}_{\Gamma^0}^{(0)} \hookrightarrow \mathcal{R}E^0 \mathcal{V}_{\Gamma^0}$ is a quasi-isomorphism. On the other hand, by Proposition 4.2, (i), the composition ν_{i_0} is an isomorphism of the vector spaces $\mathcal{O}(X^0) \otimes \overline{\Omega}_\emptyset \otimes V_\emptyset$ and

$E^0\mathcal{V}_{\Gamma^0}$. Thus the map ν is also a quasi-isomorphism of the complex $(\mathcal{R}E^0\mathcal{V}_{\Gamma^0}, d)$ and of $E^0\mathcal{V}_{\Gamma^0}$, treated as the complex with zero differential. $\square$

7.9. Proof of Theorem 4.13. Let $D_X^k \subset D_X$ denote the subspace of differential operators of order less than or equal to k .

Fix a quiver $\mathcal{V} = \{V_\alpha, A_{\alpha,\beta}\}$. Let $\alpha \in I(\Gamma)$. For any $v_\alpha \in V_\alpha$ and $\bar{w}_\alpha \in \bar{\Omega}_\alpha$, set $\deg(\bar{w}_\alpha \otimes v_\alpha) = l(\alpha)$. Set also $\deg \xi = 1$ for any $\xi \in T_\alpha$ and $\deg f = 0$ for any $f \in F_\alpha$. Hence $\deg x = p + l(\alpha)$ for any $x \in \bigwedge^p T_\alpha \wedge \bigwedge^q F_\alpha \otimes \bar{\Omega}_\alpha \otimes V_\alpha$. Define $\overline{\mathcal{R}EV}^k$ as the subspace of $\overline{\mathcal{R}EV}$, spanned by linear combinations

$$\sum_{\alpha \in I(\Gamma)} y_\alpha x_\alpha,$$

where $x_\alpha \in \bigwedge^* T_\alpha \otimes \bar{\Omega}_\alpha \otimes V_\alpha$, $y_\alpha \in \mathcal{D}_X$ and $\deg x_\alpha + \deg y_\alpha \leq k$.

Lemma 7.8.

- (i) For any $k \geq 0$, $\overline{\mathcal{R}EV}^k$ is a subcomplex of $\overline{\mathcal{R}EV}$.
- (ii) For any $k, l \geq 0$, there are inclusions $\mathcal{D}_X^k \overline{\mathcal{R}EV}^l \subset \overline{\mathcal{R}EV}^{k+l}$.

$\square$

By Lemma 7.8 we have a filtration

$$0 \subset \overline{\mathcal{R}EV}^0 \subset \overline{\mathcal{R}EV}^1 \subset \dots \subset \overline{\mathcal{R}EV}$$

of the complex $\overline{\mathcal{R}EV}$ by subcomplexes $\overline{\mathcal{R}EV}^k$. We can view this filtration as a pair: a filtration

$$0 \subset \mathcal{R}EV^0 \subset \mathcal{R}EV^1 \subset \dots \subset \mathcal{R}EV$$

of the complex $\mathcal{R}EV$, and a filtration

$$0 \subset EV^0 \subset EV^1 \subset \dots \subset EV$$

of the $\mathcal{D}_X$ -module EV . This filtration coincides with the principal filtration $F_p EV$ of EV , see Section 4.2. We also have the map ν of filtered complexes, $\nu : \mathcal{R}EV \rightarrow EV_{(\cdot)}$. Let $\tilde{\mathcal{R}EV}^k = \mathcal{R}EV^k / \mathcal{R}EV^{k-1}$, and $\tilde{EV}^k = EV^k / EV^{k-1}$. The space $\tilde{\mathcal{R}EV}^k$ is spanned by linear combinations $\sum_{\alpha \in I(\Gamma)} y_\alpha x_\alpha$, where $x_\alpha \in \bigwedge^* T_\alpha \otimes \bar{\Omega}_\alpha \otimes V_\alpha$ is an element of some degree

j , y is the symbol of a differential operator of degree $k - j$ in $\mathcal{D}_X^{k-j} / \mathcal{D}_X^{k-j-1}$.

The differential $\tilde{d}$ in the graded quotient complex $\tilde{\mathcal{R}EV} = \bigoplus_{k \geq 0} \tilde{\mathcal{R}EV}^k$ can be presented as the difference $\tilde{d} = \tilde{d}_1 - \tilde{d}_0$, where

$$\begin{aligned} \tilde{d}_1(y_\alpha \otimes t_1 \wedge \dots \wedge t_r \otimes \bar{w}_\alpha \otimes v_\alpha) &= \sum_{i=1}^r (-1)^{i+1} y_\alpha t_i \otimes t_1 \wedge \dots \wedge \widehat{t_i} \dots \wedge t_r \otimes \bar{w}_\alpha \otimes v_\alpha, \\ \tilde{d}_0(y_\alpha \otimes t_1 \wedge \dots \wedge t_r \otimes \bar{w}_\alpha \otimes v_\alpha) &= \sum_{\beta, \alpha \succ \beta} y_\alpha \otimes A_{\beta,\alpha}^r(t_1 \wedge \dots \wedge t_r \otimes \bar{w}_\alpha \otimes v_\alpha). \end{aligned}$$

By Proposition 4.1, the space EV can be identified with the space M ,

$$M = \bigoplus_{\alpha \in I(\Gamma)} S(T'_\alpha) \otimes S(F'_\alpha) \otimes \bar{\Omega}_\alpha \otimes V_\alpha.$$

Here $S(T'_\alpha) = \bigoplus_{j \geq 0} S^j(T'_\alpha)$ and $S(F'_\alpha) \oplus_{j \geq 0} S^j(F'_\alpha)$ are the symmetric algebras of the spaces T'_α and F'_α introduced in Section 2.2. The identification map $\varphi : M \rightarrow E\mathcal{V}$,

$$(7.45) \quad \varphi(\xi_{i_1} \cdots \xi_{i_n} \otimes f_{j_1} \cdots f_{j_m} \otimes \bar{\omega}_\alpha \otimes v_\alpha) = \xi_{i_1} \cdots \xi_{i_n} f_{j_1} \cdots f_{j_m} (\bar{\omega}_\alpha \otimes v_\alpha)$$

is an application of commuting elements $\xi_{i_k} \in T'_\alpha \subset \mathcal{D}_X$ and $f_{j_l} \in F'_\alpha \subset \mathcal{D}_X$ to the vectors $\bar{\omega}_\alpha \otimes v_\alpha \in E\mathcal{V}$.

For any $\alpha \in I(\Gamma)$, $\xi \in T'_\alpha$, and $f \in F'_\alpha$, set $\deg \xi = 1$, $\deg f = 0$, and $\deg(\bar{\omega}_\alpha \otimes v_\alpha) = l(\alpha)$. Thus, M is a graded space with graded components

$$M^{(k)} = \bigoplus_{\alpha \in I(\Gamma)} \bigoplus_{j, j+l(\alpha)=k} S^j(T'_\alpha) \otimes S(F'_\alpha) \otimes \bar{\Omega}_\alpha \otimes V_\alpha.$$

Set $M^k = \bigoplus_{j \leq k} M^{(j)}$. The spaces M^k , $k \geq 0$, give an increasing filtration on M : $0 \subset M^0 \subset \dots \subset M^k \subset \dots \subset M$.

Lemma 7.9. *The map (7.45) establishes an isomorphism of the filtered spaces M and $E\mathcal{V}$.*

Proof. We know by Proposition 4.1 that the linear map (7.45) is an isomorphism. So it is sufficient to establish an equality of vector spaces

$$(7.46) \quad \varphi(M^k) = E\mathcal{V}^k.$$

for any $k \geq 0$. We prove (7.46) by induction over k . By definition of M^k and $E\mathcal{V}^k$, we have $\varphi(M^k) \subset E\mathcal{V}^k$ and $\varphi(M^0) = E\mathcal{V}^0$. Suppose we know that $\varphi(M^{k-1}) = E\mathcal{V}^{k-1}$ for some $k \geq 1$. Consider an element $x = y \cdot \bar{\omega}_\alpha \otimes v_\alpha \in E\mathcal{V}^k$, where $y \in \mathcal{D}_X^{k-l(\alpha)}$. We prove by induction on $\dim X_\alpha = N - l(\alpha)$ that $x \in \varphi(M^k)$. Present x as

$$(7.47) \quad x = \sum_i \xi'_i f'_i \xi_i f_i (\bar{\omega}_\alpha \otimes v_\alpha),$$

where $\xi'_i, f'_i, \xi_i, f_i \in \mathcal{D}_X$ are homogeneous elements of the symmetric algebras over $T'_\alpha, F'_\alpha, T_\alpha, F_\alpha$, respectively. The summands in (7.47) with $f_i \in S^+(F_\alpha) = \bigoplus_{j>0} S^j(F_\alpha)$ belong to $E\mathcal{V}^{k-1}$. The summands in (7.47) with $f_i = 1$ but $\xi_i \in S^+(T_\alpha)$ can be presented in the form $\sum_{\beta, l(\beta) > l(\alpha)} y_\beta (\bar{\omega}_\beta \otimes v_\beta)$ and belong to $E\mathcal{V}^k$ according to the assumption of the induction on $N - l(\alpha)$. For vertices α , $l(\alpha) = N$, the summands in (7.47) with $f_i = 1$ but $\xi_i \in S^+(T_\alpha)$ are missing since $T_\alpha = 0$ in this case. The rest of the summands is of the form $\xi'_i f'_i (\bar{\omega}_\alpha \otimes v_\alpha)$ with $\xi'_i \in S(T'_\alpha)$, $f'_i \in S(F'_\alpha)$, and belong to $\varphi(M^k)$ by the definition of φ . $\square$

Let $\tilde{\nu} : \tilde{\mathcal{R}}E\mathcal{V}_0 \rightarrow \tilde{E}\mathcal{V} = \bigoplus_{k \geq 0} \tilde{E}\mathcal{V}^k$ be the map, induced by ν . Identify $\tilde{E}\mathcal{V}^k$ with $M^{(k)}$. The proof of Lemma 7.9 shows that for any $y \in \mathcal{D}_X^l / \mathcal{D}_X^{l-1}$ and $\alpha \in I(\Gamma)$ we can write the image $\tilde{\nu}(y \otimes \bar{\omega}_\alpha \otimes v_\alpha)$ as

$$\tilde{\nu}(y \otimes \bar{\omega}_\alpha \otimes v_\alpha) = \bigoplus_{\beta, \beta \geq \alpha} \tilde{\nu}_{\beta, \alpha}(y \otimes \bar{\omega}_\alpha \otimes v_\alpha),$$

where $\tilde{\nu}_{\beta, \alpha}(y \otimes \bar{\omega}_\alpha \otimes v_\alpha) \in S(T'_\beta) \otimes S(F'_\beta) \otimes \bar{\Omega}_\beta \otimes V_\beta$ and for any $\xi' \in S(T'_\alpha)$, $f' \in S(F'_\alpha)$, $\xi \in S(T_\alpha)$, $f \in S(F_\alpha)$,

$$\tilde{\nu}_{\alpha, \alpha}(\xi' f' \xi f \otimes \bar{\omega}_\alpha \otimes v_\alpha) = \varepsilon(\xi) \varepsilon(f) \xi' \otimes f' \otimes \bar{\omega}_\alpha \otimes v_\alpha.$$

Here

$$(7.48) \quad \varepsilon : S(V) \rightarrow \mathbb{C}$$

is the augmentation map, equal to the projection of the symmetric algebra of a vector space V to its zero graded component $S^0(V) \approx \mathbb{C}$ parallel to $S^+(V) = \bigoplus_{k>0} S^k(V)$.

For a fixed $k \geq 0$ and $l = 0, \dots, N = \dim X$, define $\tilde{\mathcal{R}}EV^{k,l}$ as the vector subspace of the space $\tilde{\mathcal{R}}EV^k$, spanned by the linear combinations

$$\sum_{\alpha, l(\alpha) \geq l} y_\alpha x_\alpha,$$

where $x_\alpha \in \bigwedge^* \mathcal{T}_\alpha \otimes \overline{\Omega}_\alpha \otimes V_\alpha$, y_α is the symbol of a homogeneous differential operator in $\mathcal{D}_X$, such that $\deg x_\alpha + \deg y_\alpha = k$. Define also $\tilde{E}\mathcal{V}^{k,l}$ to be the subspace of the space $\tilde{E}\mathcal{V}^k$ generated by the elements $\xi' \otimes f' \otimes \overline{\omega}_\alpha \otimes v_\alpha$ with $l(\alpha) \geq l$.

Lemma 7.10. *Let $k > 0$ and $l = 0, \dots, N = \dim X$. Then*

- (i) $\tilde{\mathcal{R}}EV^{k,l}$ is a subcomplex of $\tilde{\mathcal{R}}EV^k$,
- (ii) $\tilde{\mathcal{R}}EV^{k,l} \subset \tilde{\mathcal{R}}EV^{k,l'}$ and $\tilde{E}\mathcal{V}^{k,l} \subset \tilde{E}\mathcal{V}^{k,l'}$ if $l > l'$,
- (iii) $\tilde{\nu}(\tilde{\mathcal{R}}EV_0^{k,l}) \subset \tilde{E}\mathcal{V}^{k,l}$.

□

By Lemma 7.10, for any $k > 0$, we have the filtration of complexes,

$$0 \subset \tilde{\mathcal{R}}EV^{k,N} \subset \dots \subset \tilde{\mathcal{R}}EV^{k,0} = \tilde{\mathcal{R}}EV^k,$$

the filtration of the space $\tilde{E}\mathcal{V}^k$,

$$0 \subset \tilde{E}\mathcal{V}^{k,N} \subset \dots \subset \tilde{E}\mathcal{V}^{k,0} = \tilde{E}\mathcal{V}^k$$

and the map $\tilde{\nu}$ of filtered complexes $\tilde{\nu} : \tilde{\mathcal{R}}EV^k \rightarrow \tilde{E}\mathcal{V}_{(\cdot)}^k$. For $l = 0, \dots, N$ let $\hat{\mathcal{R}}EV^{k,l} = \tilde{\mathcal{R}}EV^{k,l} / \tilde{\mathcal{R}}EV^{k,l+1}$ be the quotient complex of $\tilde{\mathcal{R}}EV^k$ and $\hat{E}\mathcal{V}^{k,l} = \tilde{E}\mathcal{V}^{k,l} / \tilde{E}\mathcal{V}^{k,l+1}$ the quotient module of $\tilde{E}\mathcal{V}^k$. The space $\hat{\mathcal{R}}EV^{k,l}$ is spanned by linear combinations $\sum_{\alpha, l(\alpha)=l} y_\alpha x_\alpha$, where $x_\alpha \in \bigwedge^* \mathcal{T}_\alpha \otimes \overline{\Omega}_\alpha \otimes V_\alpha$, y_α is the symbol of a homogeneous differential operator in $\mathcal{D}_X$ such that $\deg x_\alpha + \deg y_\alpha = k$. The space $\hat{E}\mathcal{V}^{k,l}$ is generated by linear combinations $\sum_{\alpha, l(\alpha)=l} \xi'_\alpha \otimes f'_\alpha \otimes \overline{\omega}_\alpha \otimes v_\alpha$, where $\xi'_\alpha \in S^{k-l}(\mathcal{T}'_\alpha)$, $f'_\alpha \in S(F'_\alpha)$.

Let $\hat{d}$ be the differential in the quotient complex $\hat{\mathcal{R}}EV = \bigoplus_{k,l} \hat{\mathcal{R}}EV^{k,l}$ and $\hat{\nu}$ the map $\hat{\mathcal{R}}EV_0 \rightarrow \hat{E}\mathcal{V} = \bigoplus_{k,l} \hat{E}\mathcal{V}^{k,l}$ induced by $\tilde{\nu}$. We have $\hat{d} = \hat{d}_1$, where

$$(7.49) \quad \hat{d}_1(y_\alpha \otimes t_1 \wedge \dots \wedge t_r \otimes \overline{\omega}_\alpha \otimes v_\alpha) = \sum_{i=1}^r (-1)^{i+1} y_\alpha t_i \otimes t_1 \wedge \dots \wedge \hat{t}_i \dots \wedge t_r \otimes \overline{\omega}_\alpha \otimes v_\alpha,$$

and

$$(7.50) \quad \hat{\nu}(\xi' f' \xi f \otimes \overline{\omega}_\alpha \otimes v_\alpha) = \varepsilon(\xi) \varepsilon(f) \xi' \otimes f' \otimes \overline{\omega}_\alpha \otimes v_\alpha.$$

Relations (7.49) and (7.50) show that the complex $\hat{\mathcal{R}}EV$ is the direct sum of its subcomplexes $\hat{\mathcal{R}}EV_\alpha$, generated by the elements of the form $y_\alpha x_\alpha$ with a fixed $\alpha \in I(\Gamma)$.

The vector space $\widehat{E}$ is the direct sum of its subspaces $\widehat{E}_\alpha$, generated by the elements $\xi' \otimes f' \otimes \overline{\omega}_\alpha \otimes v_\alpha$. The complex $\widehat{\mathcal{R}}E\mathcal{V}_\alpha$, the space $\widehat{E}_\alpha$ and the map $\widehat{\nu}$ form the complex

$$\widehat{\mathcal{R}}E\mathcal{V}: \quad 0 \rightarrow \widehat{\mathcal{R}}^{-N}E\mathcal{V} \xrightarrow{\widehat{d}} \widehat{\mathcal{R}}^{-N+1}E\mathcal{V} \xrightarrow{\widehat{d}} \cdots \xrightarrow{\widehat{d}} \widehat{\mathcal{R}}^0E\mathcal{V} \xrightarrow{\widehat{\nu}} \widehat{E}\mathcal{V} \rightarrow 0.$$

This complex is isomorphic to the tensor product of Koszul complexes related to symmetric algebras $S(T_\alpha)$ and $S(F_\alpha)$ with coefficients in the rings $S(T'_\alpha) \otimes S(F'_\alpha)$:

$$\widehat{\mathcal{R}}E\mathcal{V} \approx \bigoplus_{\alpha \in I(\Gamma)} S(T'_\alpha) \otimes S(F'_\alpha) \otimes \text{Kos}_{T_\alpha} \otimes \text{Kos}_{F_\alpha}.$$

Recall that for a vector space V of the dimension N , the Koszul complex Kos_V is the total complex of free $S(V)$ -module resolution of the trivial $S(V)$ -module $\mathbb{C}$:

$$\text{Kos}_V: \quad 0 \rightarrow S(V) \otimes \bigwedge^N (V^*) \xrightarrow{d_{\text{Kos}}} \cdots \xrightarrow{d_{\text{Kos}}} S(V) \otimes \bigwedge^0 (V^*) \xrightarrow{\varepsilon} \mathbb{C} \rightarrow 0,$$

where $d_{\text{Kos}} = \sum_{i=1}^N e_i \otimes \frac{\partial}{\partial e^i}$ with $\{e_i\}$ and $\{e^i\}$, $i = 1, \dots, N$, being dual bases of V and V^* , and ε is the augmentation map $\varepsilon: S(V) \rightarrow \mathbb{C}$, see (7.48).

The complex $\widehat{\mathcal{R}}E\mathcal{V}$ is acyclic, since Koszul complexes are acyclic. Then the complex $\widetilde{\mathcal{R}}E\mathcal{V}$, as well as $\overline{\mathcal{R}}E\mathcal{V}$ is acyclic, since all its graded factors are acyclic. Theorem 4.13 is proved. $\square$

7.10. Proof of Theorem 4.14. We prove statement (ii) of the theorem. Statement (i) is a specialization of (ii) to the open set X^0 .

First we reformulate statement (ii). Let $\mathcal{V}$ be a quiver of Γ . Consider the complex

$$\text{RHom}_{\mathcal{D}_X}(\mathcal{R}E\mathcal{V}, \widetilde{\mathcal{D}}_X)[N],$$

and denote it by $\widetilde{\mathcal{R}}E\mathcal{V}$. The complex has $N + 1$ terms. Write it as

$$\widetilde{\mathcal{R}}E\mathcal{V}: \quad 0 \rightarrow \widetilde{\mathcal{R}}_{-N}E\mathcal{V} \rightarrow \cdots \rightarrow \widetilde{\mathcal{R}}_0E\mathcal{V} \rightarrow 0.$$

Then $\deg \widetilde{\mathcal{R}}_kE\mathcal{V} = k$, and

$$\begin{aligned} \widetilde{\mathcal{R}}_{-r}E\mathcal{V} &= \text{Hom}_{\mathcal{D}_X} \left(\bigoplus_{\alpha \in I(\Gamma)} \mathcal{D}_X \otimes \left(\bigwedge^{N-r} \mathcal{T}_\alpha \otimes \overline{\Omega}_\alpha \otimes V_\alpha \right), \widetilde{\mathcal{D}}_X \right) \approx \\ &\approx \bigoplus_{\alpha \in I(\Gamma)} \widetilde{\mathcal{D}}_X \otimes \left(\bigwedge^{N-r} \mathcal{T}_\alpha \otimes \overline{\Omega}_\alpha \otimes V_\alpha \right)^*. \end{aligned}$$

We have to prove that $\widetilde{\mathcal{R}}E\mathcal{V}$ is quasi-isomorphic to the complex $\mathcal{R}E(\tau\mathcal{V})$.

Choose a vector $\eta \in \bigwedge^N T_\emptyset$, $\eta = c \frac{\partial}{\partial z_1} \wedge \cdots \wedge \frac{\partial}{\partial z_N}$ in local affine coordinates $z_1, \dots, z_N$ of $\mathbb{C}^N$, and define a pairing

$$\left(\bigwedge^r \mathcal{T}_\alpha \otimes \overline{\Omega}_\alpha \otimes V_\alpha \right) \otimes \left(\bigwedge^{N-r} \mathcal{T}_\alpha \otimes \overline{\Omega}_\alpha \otimes V_\alpha^* \right) \rightarrow \mathbb{C}$$

in the following way. Let $\alpha \in I_l(\Gamma)$, $\vec{\xi}_p, \vec{\xi}_{p'} \in \bigwedge^p T_\alpha$, $\vec{f}_q, \vec{f}_{q'} \in \bigwedge^q F_\alpha$, where $p + q = r$, $\overline{\omega}_\alpha, \overline{\omega}'_\alpha \in \overline{\Omega}_\alpha$, $v_\alpha \in V_\alpha$, $v'_\alpha \in V_\alpha^*$. Set

$$\begin{aligned} (7.51) \quad &\langle \vec{\xi}_p \wedge \vec{f}_q \otimes \overline{\omega}_\alpha \otimes v_\alpha, \vec{\xi}_{p'} \wedge \vec{f}_{q'} \otimes \overline{\omega}'_\alpha \otimes v'_\alpha \rangle = \\ &= (-1)^{\frac{p(p-1)+q(q-1)}{2}} \delta_{p', N-l-p} \delta_{q', l-q} \langle \vec{\xi}_p \wedge \vec{\xi}_{p'}, \overline{\omega}_\alpha \rangle \cdot \langle \eta, d\vec{f}_q \wedge \overline{\omega}'_\alpha \wedge d\vec{f}_{q'} \rangle \cdot \langle v_\alpha, v'_\alpha \rangle. \end{aligned}$$

Due to (7.51) we have an isomorphism

$$\tilde{\mathcal{R}}_{-r}E\mathcal{V} \approx \bigoplus_{\alpha \in I(\Gamma)} \tilde{\mathcal{D}}_X \otimes \left(\bigwedge^r \mathcal{T}_\alpha \otimes \overline{\Omega}_\alpha \otimes V_\alpha \right).$$

The differential $\tilde{d} : \tilde{\mathcal{R}}_{-r}E\mathcal{V} \rightarrow \tilde{\mathcal{R}}_{-r+1}E\mathcal{V}$, adjoint to the differential (4.47), takes the form $\tilde{d} = \tilde{d}_1 - \tilde{d}_0$,

$$\begin{aligned} \tilde{d}_1 \left(\tilde{D} \otimes t_1 \wedge \dots \wedge t_r \otimes \overline{\omega}_\alpha \otimes v_\alpha \right) &= \sum_{i=1}^r (-1)^{i+1} t_i^{(I)}(\tilde{D}) \otimes t_1 \wedge \dots \widehat{t_i} \dots \wedge t_r \otimes \overline{\omega}_\alpha \otimes v_\alpha, \\ \tilde{d}_0 \left(\tilde{D} \otimes t_1 \wedge \dots \wedge t_r \otimes \overline{\omega}_\alpha \otimes v_\alpha \right) &= \sum_{\beta, (\alpha, \beta) \in E(\Gamma)} \tilde{D} \otimes A_{\beta, \alpha}^r(t_1 \wedge \dots \wedge t_r \otimes \overline{\omega}_\alpha \otimes v_\alpha), \end{aligned}$$

where $t_i \in \mathcal{T}_\alpha$, $\overline{\omega}_\alpha \otimes v_\alpha \in \overline{\Omega}_\alpha \otimes V_\alpha$, $\tilde{D} \in \tilde{\mathcal{D}}_X$ and $x^{(I)}(\tilde{D})$ means the action of $x \in \mathcal{D}_X$ on the element $\tilde{D} \in \tilde{\mathcal{D}}_X$ with respect to the first $\mathcal{D}_X$ -module structure on $\tilde{\mathcal{D}}_X$, defined in Section 4.10. We note that $\tilde{\mathcal{D}}_X$ is the free $\mathcal{D}_X$ -module of rank one with respect to the second $\mathcal{D}_X$ -module structure. A global section $\tilde{\omega} \in \Omega_X^{-1}$, say $\tilde{\omega} = (dz_1 \wedge \dots \wedge dz_N)^{-1}$, defines an isomorphism of free left $\mathcal{D}_X$ -modules $\phi_{\tilde{\omega}} : \tilde{\mathcal{D}}_X \rightarrow \mathcal{D}_X$,

$$\phi_{\tilde{\omega}}(D \otimes \tilde{\omega}) = D \cdot 1.$$

Applying the isomorphism $\phi_{\tilde{\omega}}$, we get the isomorphism of $\mathcal{D}_X$ -modules,

$$\tilde{\mathcal{R}}_{-r}E\mathcal{V} \approx \bigoplus_{\alpha \in I(\Gamma)} \mathcal{D}_X \otimes \left(\bigwedge^r \mathcal{T}_\alpha \otimes \overline{\Omega}_\alpha \otimes V_\alpha \right).$$

Under this identification the differential in $\tilde{\mathcal{R}}E\mathcal{V}$ becomes $d' = d'_1 - d'_0$,

$$\begin{aligned} d'_D(D \otimes t_1 \wedge \dots \wedge t_r \otimes \overline{\omega}_\alpha \otimes v_\alpha) &= \sum_{i=1}^r (-1)^{i+1} D \cdot t_i^\tau \otimes t_1 \wedge \dots \widehat{t_i} \dots \wedge t_r \otimes \overline{\omega}_\alpha \otimes v_\alpha, \\ d'_0(D \otimes t_1 \wedge \dots \wedge t_r \otimes \overline{\omega}_\alpha \otimes v_\alpha) &= \sum_{\beta, (\alpha, \beta) \in E(\Gamma)} D \otimes A_{\beta, \alpha}^r(t_1 \wedge \dots \wedge t_r \otimes \overline{\omega}_\alpha \otimes v_\alpha), \end{aligned}$$

where $D \in \mathcal{D}_X$ and τ is an anti-isomorphism of $\mathcal{D}_X$, such that $t_i^\tau = t_i$ and $\left(\frac{\partial}{\partial t_i} \right)^\tau = -\frac{\partial}{\partial t_i}$.

Define a linear map

$$\nu : \bigoplus_{\alpha \in I(\Gamma)} \mathcal{D}_X \otimes \left(\bigwedge^r \mathcal{T}_\alpha \otimes \overline{\Omega}_\alpha \otimes V_\alpha \right) \rightarrow \bigoplus_{\alpha \in I(\Gamma)} \mathcal{D}_X \otimes \left(\bigwedge^r \mathcal{T}_\alpha \otimes \overline{\Omega}_\alpha \otimes V_\alpha \right)$$

as multiplication by $(-1)^p$ on each subspace of the form $\mathcal{D}_X \otimes \bigwedge^p \mathcal{T}_\alpha \wedge \bigwedge^q F_\alpha \otimes \overline{\Omega}_\alpha \otimes V_\alpha$. Conjugation of d' with ν transforms the complex $\tilde{\mathcal{R}}E\mathcal{V}$ to the complex $\mathcal{R}E(\tau\mathcal{V})$. The theorem is proved. $\square$

7.11. Proof of Theorem 5.6. Let $\mathcal{C}$ be a central arrangement of hyperplanes in $X = \mathbb{C}^N$. Let $\mathcal{V} \in \mathcal{Q}ui$, $\mathcal{V} = \{V_\alpha, A_{\alpha, \beta}\}$, be a quiver of the arrangement $\mathcal{C}$ and $\Gamma\mathcal{QV}$ the complex of global sections of the de Rham complex $\mathcal{QV}$. The component $\Gamma\mathcal{QV}^r$, $-n \leq r \leq 0$, of this complex is isomorphic to

$$\Gamma\mathcal{QV}^r = \bigoplus_{\alpha \in \Gamma} \Gamma\Omega_X^{an} \otimes \left(\bigwedge^{-r} \mathcal{T}_\alpha \otimes \overline{\Omega}_\alpha \otimes V_\alpha \right),$$

where $\Gamma\Omega_X^{an}$ denotes the space of global holomorphic N -forms $f(z_1, \dots, z_N)dz_1 \wedge \dots \wedge dz_N$ on $\mathbb{C}^N$ with analytic coefficients. The differential d , $d(\Gamma\mathcal{QV}_r) \subset \Gamma\mathcal{QV}_{r+1}$ in the complex $\Gamma\mathcal{QV}$ is given by formula (5.11).

We construct below a new differential $d^* : \Gamma\mathcal{QV} \rightarrow \Gamma\mathcal{QV}$, such that $d^*(\Gamma\mathcal{QV}_r) \subset \Gamma\mathcal{QV}_{r-1}$. We construct the Laplace operator $\Delta = dd^* + d^*d$. Using the Laplace operator, we prove that the complex $\Gamma\mathcal{QV}$ is quasi-isomorphic to the quiver complex $C_+(\mathcal{V})[N]$ under the assumption that the matrix coefficients of the linear maps $A_{\alpha,\beta}$ are small enough.

The definition of the differential d^* depends on the choice of a non-degenerate symmetric bilinear form $(\cdot, \cdot)$ on the vector space $\mathbb{C}^N$.

For any $\alpha \in I(\Gamma)$, let $\{\xi_i\}$ be an orthonormal basis of the space T_α , $(\xi_i, \xi_j) = \delta_{i,j}$, and let $\{\xi^k\}$ be an orthonormal basis of the space T'_α (see Section 2.2), $(\xi^k, \xi^l) = \delta_{k,l}$. We have $(\xi_i, \xi^k) = 0$, since the spaces T_α and T'_α are orthogonal. We denote by $I(T_\alpha)$ the set of the indices of the basis $\{\xi_i\}$ of the space T_α . It contains $\dim T_\alpha$ elements. We denote by $I(F_\alpha)$ the set of the indices of the basis $\{\xi^k\}$ of the space T'_α . It contains $\dim F_\alpha$ elements.

Let f_l , $l \in I(F_\alpha)$, be the elements of F_α , such that $\langle \xi^k, f_l \rangle = \delta_{k,l}$. Let f^j , $j \in I(T_\alpha)$, be the elements of F'_α , such that $\langle \xi_i, f^j \rangle = \delta_{i,j}$. They form an orthonormal basis of the space $\tilde{F} \simeq F_\alpha \oplus F'_\alpha$ of all linear function on $\mathbb{C}^N$.

Let $d_1^* : \Gamma\mathcal{QV} \rightarrow \Gamma\mathcal{QV}$ be the linear map, defined by the formula

$$d_1^*(\omega \otimes \bar{t} \otimes \bar{\omega}_\alpha \otimes v_\alpha) = \sum_{i \in I(T_\alpha)} \omega \cdot f^i \otimes \xi_i \wedge \bar{t} \otimes \bar{\omega}_\alpha \otimes v_\alpha + \sum_{k \in I(F_\alpha)} \omega \cdot \xi^k \otimes f_k \wedge \bar{t} \otimes \bar{\omega}_\alpha \otimes v_\alpha,$$

where $\alpha \in I(\Gamma)$, $\omega \in \Gamma\Omega_X^{an}$, $\bar{t} \in \bigwedge^* T_\alpha$, $\bar{\omega}_\alpha \in \bar{\Omega}_\alpha$, $v_\alpha \in V_\alpha$.

Choose an edge framing $\{f_{\beta,\alpha}, \xi_{\alpha,\beta}\}$ in such a way that for any edge $(\alpha, \beta) \in E(\Gamma)$, $\alpha \succ \beta$, we have $(\xi_{\alpha,\beta}, \xi_\beta) = 0$ for any $\xi_\beta \in T_\beta$, $\langle \xi', f_{\beta,\alpha} \rangle = 0$ for any $\xi' \in T'_\alpha$ and $\langle \xi_{\alpha,\beta}, f_{\beta,\alpha} \rangle = 1$. In particular, the last equality implies that the maps $\pi_{\alpha,\beta}$ and $\pi_{\beta,\alpha}$ are inverse to each other. We call such a framing orthogonal.

The bilinear form $(\cdot, \cdot)$ on $\mathbb{C}^N$ defines for any $\alpha \succ \beta$ orthogonal projections $T_\alpha \rightarrow T_\beta$ and $F_\beta \rightarrow F_\alpha$, which extend to projections $P_{\beta,\alpha} : \bigwedge^* T_\alpha \rightarrow \bigwedge^* T_\beta$ and $P_{\alpha,\beta} : \bigwedge^* F_\beta \rightarrow \bigwedge^* F_\alpha$. For an orthogonal edge framing $\{f_{\beta,\alpha}, \xi_{\alpha,\beta}\}$ we have

$$\begin{aligned} P_{\beta,\alpha}(\xi_\alpha \wedge \bar{\xi}_\beta) &= (\xi_\alpha - \langle \xi_\alpha, f_{\beta,\alpha} \rangle \xi_{\alpha,\beta}) \wedge \bar{\xi}_\beta, \\ P_{\alpha,\beta}(f_\beta \wedge \bar{f}_\alpha) &= (f_\beta - \langle \xi_{\alpha,\beta}, f_\beta \rangle f_{\beta,\alpha}) \wedge \bar{f}_\alpha, \end{aligned}$$

if $\xi_\alpha \in T_\alpha$, $\bar{\xi}_\beta \in \bigwedge^* T_\beta \subset \bigwedge^* T_\alpha$, $f_\beta \in F_\beta$, $\bar{f}_\alpha \in \bigwedge^* F_\beta \subset \bigwedge^* F_\alpha$.

For each edge $(\alpha, \beta) \in E(\Gamma)$ define a linear map

$$d_{0,(\beta,\alpha)}^* : \Gamma\Omega_X^{an} \otimes \bigwedge^* T_\alpha \otimes \bar{\Omega}_\alpha \otimes V_\alpha \rightarrow \Gamma\Omega_X^{an} \otimes \bigwedge^* T_\beta \otimes \bar{\Omega}_\beta \otimes V_\beta,$$

as follows. Let $\omega \in \Gamma\Omega_X^{an}$, $\bar{\xi}_\alpha \in \bigwedge^* T_\alpha$, $\bar{f}_\alpha \in \bigwedge^* F_\alpha$, $\bar{\omega}_\alpha \in \bar{\Omega}_\alpha$, $v_\alpha \in V_\alpha$. We set

$$d_{0,(\beta,\alpha)}^*(\omega \otimes \bar{\xi}_\alpha \wedge \bar{f}_\alpha \otimes \bar{\omega}_\alpha \otimes v_\alpha) = \omega \otimes f_{\beta,\alpha} \wedge P_{\beta,\alpha}(\bar{\xi}_\alpha) \wedge \bar{f}_\alpha \otimes \pi_{\beta,\alpha}(\bar{\omega}_\alpha) \otimes A_{\beta,\alpha}(v_\alpha)$$

if $\alpha \succ \beta$, and

$$d_{0,(\beta,\alpha)}^*(\omega \otimes \bar{\xi}_\alpha \wedge \bar{f}_\alpha \otimes \bar{\omega}_\alpha \otimes v_\alpha) = \omega \otimes \xi_{\beta,\alpha} \wedge \bar{\xi}_\alpha \wedge P_{\beta,\alpha}(\bar{f}_\alpha) \otimes \pi_{\beta,\alpha}(\bar{\omega}_\alpha) \otimes A_{\beta,\alpha}(v_\alpha)$$

if $\beta \succ \alpha$. Let

$$d_0^* = \sum_{(\alpha, \beta) \in E(\Gamma)} d_{0,(\beta, \alpha)}^*.$$

Lemma 7.11. *The maps d_i^* , $i = 0, 1$, form two commuting differentials,*

$$(d_i^*)^2 = 0 \quad \text{for } i = 0, 1, \quad d_0^* d_1^* + d_1^* d_0^* = 0.$$

□

Define $d^* = d_1^* + d_0^*$ and $\Delta = dd^* + d^*d$. We decompose Δ as

$$\Delta = \Delta^{(2)} + \Delta^{(1)} + \Delta^{(0)},$$

where $\Delta^{(2)} = d_1^* d_1 + d_1 d_1^*$, $\Delta^{(1)} = d_0^* d_1 + d_1 d_0^* - d_1^* d_0 - d_0 d_1^*$, $\Delta^{(0)} = -d_0^* d_0 - d_0 d_0^*$.

Denote by $\overline{V}_\alpha$ the following finite-dimensional space:

$$\overline{V}_\alpha = \bigwedge^* \mathcal{T}_\alpha \otimes \overline{\Omega}_\alpha \otimes V_\alpha,$$

and let $\overline{V} = \bigoplus_{\alpha \in I(\Gamma)} \overline{V}_\alpha$.

Lemma 7.12. *For any $\omega \in \Gamma\Omega_X^{an}$, $\overline{v}_\alpha \in \overline{V}_\alpha$, and $\overline{v} \in \overline{V}$, we have*

$$\begin{aligned} (i) \quad \Delta^{(2)}(\omega \otimes \overline{v}_\alpha) &= \omega \cdot \left(\sum_{i \in I(T_\alpha)} f^i \xi_i + \sum_{k \in I(F_\alpha)} \xi^k f_k \right) \otimes \overline{v}_\alpha, \\ (ii) \quad \Delta^{(1)}(\omega \otimes \overline{v}) &= 0, \\ (iii) \quad \Delta^{(0)}(\omega \otimes \overline{v}) &= \omega \otimes \overline{\Delta}^{(0)}(\overline{v}) \quad \text{for some linear operator } \overline{\Delta}^{(0)} : \overline{V} \rightarrow \overline{V}. \end{aligned}$$

□

Let $\alpha \in I(\Gamma)$. For any $v_\alpha \in V_\alpha$ and $\overline{w}_\alpha \in \overline{\Omega}_\alpha$, set $\deg(\overline{w}_\alpha \otimes v_\alpha) = \dim X_\alpha = N - l(\alpha)$. Set $\deg \xi = -1$ for any $\xi \in T_\alpha$ and $\deg f = 1$ for any $f \in F_\alpha$. Thus for any $x \in \bigwedge^p T_\alpha \wedge \bigwedge^q F_\alpha \otimes \overline{\Omega}_\alpha \otimes V_\alpha$, we have $\deg x = p - q + N - l(\alpha)$. This grading gives the decomposition $\overline{V} = \bigoplus_{k=0}^N \overline{V}^k$, where $\overline{V}^k$ consists of elements $v \in \overline{V}$ of degree k . We say that a polynomial differential form $\omega = f(z_1, \dots, z_N) dz_1 \wedge \dots \wedge dz_N$ has degree $k \geq 0$ and write $\deg \omega = k$, if $f(z_1, \dots, z_N)$ is a homogeneous polynomial of degree k variables $z_1, \dots, z_N$. We say that a holomorphic form $\omega = f(z_1, \dots, z_N) dz_1 \wedge \dots \wedge dz_N$ has degree not less than $k \geq 0$ and write $\deg \omega \geq k$, if the holomorphic function $f(z_1, \dots, z_N)$ has zero of order not less than k at the origin $z_1 = z_2 = \dots = z_N = 0$.

Define $\Gamma\mathcal{QV}^k$ as the subspace of the vector space $\Gamma\mathcal{QV}$, spanned by linear combinations

$$\sum_{j=0}^N \omega^{k-j} v^j,$$

where $v^j \in \overline{V}^j$ and $\omega^{k-j} \in \Gamma\Omega_X^{an}$ is a holomorphic form of degree not less than $k - j$.

Lemma 7.13. *For a central arrangement $\mathcal{C}$ and for any $k \geq 0$, $\Gamma\mathcal{QV}^k$ is a subcomplex of $\Gamma\mathcal{QV}$.*

□

By Lemma 7.13 we have a decreasing filtration

$$\Gamma\mathcal{QV} = \Gamma\mathcal{QV}^0 \supset \Gamma\mathcal{QV}^1 \supset \dots \supset \Gamma\mathcal{QV}^k \supset \dots \supset 0$$

of the complex $\Gamma\mathcal{QV}$ by subcomplexes $\Gamma\mathcal{QV}^k$, such that $\cap_{k \geq 0} \Gamma\mathcal{QV}^k = 0$. Let us prove that the subcomplex $\Gamma\mathcal{QV}^1$ is acyclic.

Take an element $y \in \Gamma\mathcal{QV}^1$, such that $dy = 0$. It can be presented as a convergent series

$$(7.52) \quad y = \sum_{j=0}^N \sum_{k \geq 1} \omega^{j,k-j}(z) \otimes v^j,$$

where $v^j \in \overline{V}^j$ and $\omega^{j,k-j}(z)$ is a homogeneous polynomial form of degree $k-j$. The convergence means that for every $j = 0, \dots, N$ and for any $\eta_j \in (\overline{V}^j)^*$ the series

$$\sum_{k \geq 1} \omega^{j,k-j}(z) \langle v^j, \eta_j \rangle$$

converges as a power series in z .

Since the differential d is homogeneous, we have $\sum_{j=0}^N d(\omega^{j,k-j}(z) \otimes v^j) = 0$ for any $k \geq 1$. Lemma 7.12 says that for any $v^j \in \overline{V}^j$ and for any homogeneous polynomial form ω^k , $\deg \omega^k = k$, we have

$$(7.53) \quad \Delta(\omega^k \otimes v^l) = \omega^k \otimes (\overline{\Delta}^{(0)} + k + l)v^l$$

Since the Laplace operator Δ commutes with d and the eigenvalues of $\overline{\Delta}^{(0)}$ are close to zero, (7.53) implies also that $d(\omega^{j,k-j}(z) \otimes v^j) = 0$ for all j and $k \geq 1$.

The space $\overline{V}^j$ is invariant with respect to $\overline{\Delta}^{(0)}$, $\overline{\Delta}^{(0)}(\overline{V}^j) \subset \overline{V}^j$. Let $\overline{V}^j = \oplus_{\lambda} \overline{V}_{\lambda}^j$ be the decomposition of $\overline{V}^j$ in the generalized eigenspaces of the operator $\overline{\Delta}^{(0)}$,

$$\overline{V}_{\lambda}^j = \bigcup_{k=0}^{\infty} \text{Ker}(\overline{\Delta}^{(0)} - \lambda)^k|_{\overline{V}^j}.$$

Present the element $\omega^{j,k-j}(z) \otimes v^j$ as

$$(7.54) \quad \omega^{j,k-j}(z) \otimes v^j = \omega^{j,k-j}(z) \otimes \sum_{\lambda} v_{\lambda}^j,$$

where $v_{\lambda}^j \in \overline{V}_{\lambda}^j$. Relation (7.53) implies that $\omega^{j,k-j}(z) \otimes v_{\lambda}^j \in \bigcup_{m=0}^{\infty} \text{Ker}(\Delta - \lambda - k)^m$ and $d(\omega^{j,k-j}(z) \otimes v_{\lambda}^j) = 0$.

Standard arguments with the Laplace operator say the following. Let an element x be such that $dx = 0$ and let x belong to a finite-dimensional generalized eigenspace of the Laplace operator $\Delta = dd^* + d^*d$ with eigenvalue $\mu \neq 0$. Then x can be presented as the boundary

$$x = dd^* \left(\frac{x}{\mu} - \frac{(\Delta - \mu)x}{\mu^2} + \frac{(\Delta - \mu)^2 x}{\mu^3} - \dots \right),$$

which is a finite sum. This implies that for any $j = 0, \dots, N$, $k \geq 1$, and eigenvalue λ of the operator $\overline{\Delta}^{(0)}|_{\overline{V}^j}$, we have

$$(7.55) \quad \omega^{j,k-j}(z) \otimes v_\lambda^j = dd^* \left(\frac{\omega^{j,k-j}(z) \otimes v^j}{\lambda + k} - \frac{\omega^{j,k-j}(z) \otimes (\overline{\Delta}^{(0)} - \lambda - k)v^j}{(\lambda + k)^2} + \right. \\ \left. + \frac{\omega^{j,k-j}(z) \otimes (\overline{\Delta}^{(0)} - \lambda - k)^2 v^j}{(\lambda + k)^3} - \dots \right)$$

The sum in (7.55) is finite and the number of terms does not depend on k . The substitution of (7.55) to (7.54) and then to (7.52) gives the presentation of the element y as the boundary of a convergent series of homogeneous elements. Convergence follows from the structure of (7.55). This proves that the subcomplex $\Gamma \mathcal{QV}^1 \subset \Gamma \mathcal{QV}$ is acyclic and the complex $\Gamma \mathcal{QV}$ is quasi-isomorphic to the quotient complex $\Gamma \mathcal{QV} / \Gamma \mathcal{QV}^1$, which is isomorphic to the quiver complex $C_+(\mathcal{V})[N]$. This proves Theorem 5.6. $\square$

7.12. Proof of Theorem 6.6. The proof of Theorem 6.6 repeats the arguments of the proof of Theorem 5.6. For the construction of the corresponding Laplace operator we use the symmetric bilinear form on X , invariant with respect to the G -action. Such a form exists, since G is finite. Then the Laplace operator commutes with the G -action and establishes a quasi-isomorphism of G -invariants of the complexes. $\square$

REFERENCES

- [A] M. Aomoto, *Un theoreme du type M-M concernant l'integrales des fonctions multiformes*, J. Math. Pures Appl. **52** (1973), 1–11.
- [Ar] V.I. Arnol'd, *The cohomology ring of the colored braid group*, Math. Notes **5** (1969), 138–140.
- [Be] A. Beilinson, *How to glue perverse sheaves*, Lecture Notes in Math. **1289** (1986), 42–51.
- [BBD] A. Beilinson, J. Bernstein, and P. Deligne, *Fasceau perverse*, Asterisque **100** (1982).
- [Bn] J. Bernstein, *Modules over the ring of differential operators*, Funkcional'nyi Analiz i ego Pril. **5** (1971), No.2, 1–16.
- [BL] J. Bernstein and V. Lunts, *Equivariant Sheaves and Functors*, Lecture Notes in Math. **1578** (1994).
- [B] A. Borel et al., *Algebraic D-modules*, Academic Press (1987).
- [Bs] E. Brieskorn, *Sur les groupes des tresses*, Séminar Bourbaki, Lecture Notes in Math., **317** (1983), 21–44.
- [GM] S.I. Gelfand and Yu.I. Manin, *Methods of homological algebra*. Translated from the 1988 Russian original. Springer-Verlag, Berlin, 1996.
- [GMc] M. Goresky and R. MacPherson, *Intersection homology*, II, Invent. Math. **72:1** (1983), 77–130.
- [Gr] A. Grothendieck, *Sur quelques points d'algebre homologique*, Tohoku Math. Journal **9:2,3** (1957), 119–221.
- [K1] M. Kashiwara, *B-functions and Holonomic Systems*, Inv. Math. **38** (1976), 33–53.
- [K2] M. Kashiwara, *The Riemann-Hilbert Problem for Holonomic Systems*, Publ. RIMS, Kyoto Univ. **20** (1984), 319–365.
- [K3] M. Kashiwara, *Vanishing cycles for holonomic systems*, Lect. Notes in Math., **1016** (1983), 134–142.
- [K4] M. Kashiwara, *D-modules and Microlocal Calculus*, Translations of Mathematical Monographs **217**, AMS, Providence, Rhode Island.
- [KK] M. Kashiwara, T. Kawai, *On holonomic system of microdifferential equations III - system with regular singularities*, Publ. RIMS, Kyoto Univ. **17** (1981), 813–979.

- [Kh] S. Khoroshkin, *$\mathcal{D}$ -modules over the arrangements of hyperplanes*, Comm. Algebra **23** **9** (1995), 3481–3504.
- [KhS1] S. Khoroshkin and V. Schechtman, *Factorizable $\mathcal{D}$ -modules*, Math. Res. Lett. **4** (1997), no. 2-3, 239–257.
- [KhS2] S. Khoroshkin and V. Schechtman, *Non-resonance $\mathcal{D}$ -modules over arrangements of hyperplanes*, preprint math.AG/9801134 (1998).
- [M] B. Malgrange, *Polynomes de Bernstein-Sato et cohomologie evanescente*, Asterisque, **101-102** (1983), 243–267.
- [Mb] Z. Mebkhout, *Une Equivalence de Categories*, Compositio Math. **51** (1984), 51–62.
- [OS] P. Orlik and L. Solomon, *Combinatorics and topology of complements of hyperplanes*, Inv. Math., **56** (1980), 167–189.
- [SV] V. Schechtman and A. Varchenko, *Arrangements of hyperplanes and Lie algebra homology*, Invent. Math. **106** (1991), 139–194.
- [V] A. Varchenko, *Multidimensional hypergeometric functions and representation theory of Lie algebras and quantum groups*, Advanced Series in Mathematical Physics, **21**, World Scientific Publishing Co., Inc., River Edge, NJ, **1995**.